

# Cementing an unwanted relationship

Chia-Hao Chang, Li-Ta Keng, Jen-Chung Ko

Department of Internal Medicine, National Taiwan University Hospital Hsin-Chu Branch, Hsinchu, Taiwan

### Correspondence to

Dr Li-Ta Keng, Department of Internal Medicine, National Taiwan University Hospital Hsin-Chu Branch, No. 25, Lane 442, Section 1, Jingguo Road, Hsinchu 300, Taiwan; ltkeng@gmail.com

Received 22 September 2016

Revised 24 October 2016

Accepted 30 October 2016

Published Online First

16 November 2016

A 93-year-old male with COPD, hypertension and stroke presented to the emergency department with cough and dyspnoea. Two months before this episode, he had suffered a compression fracture and therefore received first and second lumbar vertebroplasty with cement implantation. In the emergency department, his oxygen saturation was 88% while breathing ambient air; his other vital signs were normal. Physical examination revealed crackles over his bilateral lung field.

A laboratory test revealed leucocytosis and a mild elevation of the C reactive protein level. A chest radiograph showed a curvilinear radiopaque foreign body with cement in the lumbar spine and right lower lung consolidation (figure 1). CT confirmed that linear-shaped cements had embolised into the right pulmonary artery and its branches in the right upper lung (figure 2). Transthoracic echocardiography revealed mild pulmonary hypertension without evidence of right heart failure. His symptoms improved after antibiotics had been administered. He lived uneventfully in the following years.

Vertebral augmentation procedures (eg, vertebroplasty and kyphoplasty) are common management strategies for vertebral compression fractures. These procedures are performed only when conventional medical therapy has not provided pain relief or when the pain substantially alters the patient's lifestyle.<sup>1</sup> The procedures involve the percutaneous



**Figure 2** CT confirmed that linear-shaped cements had embolised into the right pulmonary artery and its branches in the right upper lung (arrows).

injection of bone cement into the fractured vertebra under image guidance, and the most frequent complication is cement leakage.<sup>2</sup> Moreover, pulmonary cement embolism is a common complication, and the risk ranges between 3.5% and 23%.<sup>2</sup> This complication is diagnosed on the basis of clinical history combined with the presence of radiopaque cement emboli on the chest radiograph or CT image.<sup>3</sup> No typical image presentation exists for pulmonary cement embolism, because these emboli develop sporadically without specific lobar distribution.<sup>3</sup> Management is dependent on the position of the cement emboli and the symptoms of the patients. Initial heparinisation with bridging to oral anticoagulants for at least 6 months is preferred for symptomatic patients or central cement emboli, whereas observation is preferred for asymptomatic patients with peripherally located cement emboli.<sup>2</sup> Surgical removal of the emboli should be considered in patients with massive emboli or right heart failure.<sup>2</sup>

**Contributors** All authors were involved in literature review, drafting of manuscript and critically reviewing it for intellectual content.

**Competing interests** None.

**Patient consent** Obtained.

**Provenance and peer review** Not commissioned; externally peer reviewed.

### REFERENCES

- Garfin SR, Yuan HA, Reiley MA. New technologies in spine: kyphoplasty and vertebroplasty for the treatment of painful osteoporotic compression fractures. *Spine* 2001;26:1511–5.
- Krueger A, Bliemel C, Zettl R, et al. Management of pulmonary cement embolism after percutaneous vertebroplasty and kyphoplasty: a systematic review of the literature. *Eur Spine J* 2009;18:1257–65.
- Kim YJ, Lee JW, Park KW, et al. Pulmonary cement embolism after percutaneous vertebroplasty in osteoporotic vertebral compression fractures: incidence, characteristics, and risk factors. *Radiology* 2009;251:250–9.



**Figure 1** Chest radiograph showed a curvilinear radiopaque foreign body (arrows) with cement in lumbar spine (arrowheads) and consolidation in the right lower lung.



**To cite:** Chang C-H, Keng L-T, Ko J-C. *Thorax* 2017;**72**:766.