Abstract P149 Table 1 Summary of clinical characteristics of patients with pulmonary fibrosis in Sweden (cohort 1)

Clinical characteristic	Variable	Patients with this characteristic, n (%)	
		At or within 1 year after diagnosis (n=17,244)	At any time (n=17,244)
Comorbidities	Ischaemic heart disease	3638 (21.1)	4338 (25.2)
	Other infections of airways	2664 (15.4)	6960 (40.4)
	Acid-related disorders	2394 (13.9)	3056 (17.7)
	COPD	2249 (13.0)	2845 (16.5)
	Angina	2233 (12.9)	2539 (14.7)
	Diabetes (types 1 and 2)	1471 (8.5)	1901 (11.0)
	Bacterial pneumonia	998 (5.8)	2954 (17.1)
	Myocardial infarction	612 (3.5)	2408 (14.0)
	Sleep apnoea	514 (3.0)	665 (3.9)
	Depression	429 (2.5)	547 (3.2)
	Anxiety	329 (1.9)	450 (2.6)
	Pulmonary hypertension	303 (1.8)	447 (2.6)
	Bronchitis	280 (1.6)	1146 (6.6)
	Peripheral vascular disease	259 (1.5)	313 (1.8)
Concomitant medications	Cardiovascular medication	7965 (46.2)	11 852 (68.7)
	Anti-coagulants/anti-thrombotics	6569 (38.1)	10 493 (60.9)
	Corticosteroids for systemic use	6086 (35.3)	10 307 (59.8)
	Acid-related disorders medication	5889 (34.2)	10 183 (59.1)
	Heart failure medication	5224 (30.3)	9250 (53.6)
	Cholesterol-lowering medication	3653 (21.2)	6451 (37.4)
	Anti-neoplastic and immunomodulating agents	1793 (10.4)	3688 (21.4)
	Oxygen	1218 (7.1)	2447 (14.2)

COPD, chronic obstructive pulmonary disorder

Conclusions Analysis of the first national cohort of patients with pulmonary fibrosis in Sweden has found slightly higher incidence rates of IPF than those of previous studies, though this may be due to identification of patients with IPF using ICD-10 code J84.1 ('other interstitial pulmonary diseases with fibrosis'). Though a diagnosis algorithm was used to refine the population for those with IPF, these criteria might not exclude all patients with non-IPF interstitial lung diseases. The frequencies of clinical characteristics in this population were broadly in line with those previously observed in patient populations with IPF.

P150 NECK AS MEDIASTINAL EXTENSION: DIAGNOSIS OF SARCOIDOSIS BY CORE BIOPSY OF CERVICAL LYMPH NODES

A Fahim, MM Qasim, D Rosewarne. *New Cross Hospital, Wolverhampton, UK*

10.1136/thoraxjnl-2017-210983.292

Background Sarcoidosis is a multi-system granulomatous disease of uncertain aetiology. It is characterised by bilateral hilar lymphadenopathy. The diagnosis is best supported by the

histological evidence of non-caseating granulomas in the affected organ (s). The diagnostic procedures for histological confirmation are invasive and a less invasive approach to diagnostic pathway is warranted.

Objectives The utility of diagnostic value of neck ultrasound was retrospectively evaluated in this study. A histological diagnosis was made by ultrasound guided head and neck core biopsy to confirm clinically and radiologically suspected sarcoidosis.

Methods Following clinical assessment by a respiratory physician, 25 patients were referred for sonographic evaluation of the head and neck by a thoracic radiologist after CT scan in an attempt to avoid the use of more invasive and expensive tests such as endobronchial ultrasound (EBUS) and mediastinoscopy. Typically these patients had obvious mediastinal adenopathy +/-parenchymal lung disease, but not clinically apparent neck nodes. Where no cervical lymph node suitable for biopsy was seen, the parotid glands were evaluated and biopsied if deemed abnormal. Patients with no suitable lymph nodes and normal parotid glands were returned for consideration of other diagnostic techniques.

Results A diagnosis of sarcoidosis was made in all cases where a core biopsy of cervical lymph nodes (figure 1) was attempted (23 out of 25 patients). It is emphasised that the cervical lymph nodes in this series were not particularly enlarged, short axis dimensions being under 10 mm in the majority of cases biopsied, and that these sub-centimetre short axis lymph nodes did not have any specific sonographic appearances to mark them as pathological. Nevertheless histological examination revealed non-caseating granulomas in all cases. In a further two cases, where no neck nodes were seen, a histological diagnosis of sarcoidosis was made from biopsy of diffusely abnormal parotid gland tissue.

Conclusions Given the clear advantages of cervical diagnosis in terms of invasiveness and economy compared to mediastinal alternatives, it is suggested that where the expertise for core biopsy of normal sized cervical lymph nodes is readily available, the technique may be considered as a first line investigation for the diagnosis of sarcoidosis.

P151 THE ROLE OF BRONCHOALVEOLAR LAVAGE AND ITS QUALITY IN THE DIAGNOSIS OF INTERSTITIAL LUNG DISEASE

A Ebraheem, L Macfarlane, R Booton, N Chaudhuri. *University Hospital Of South Manchester NHS Foundation Trust, Manchester, UK*

10.1136/thoraxjnl-2017-210983.293

Background Bronchoalveolar lavage (BAL) is a widely used diagnostic tool for suspected Interstitial Lung Disease (ILD) patients and is advocated in ILD guidelines.¹ We aimed to evaluate the quality of BAL sampling in the diagnosis of ILD compared to BAL guidelines.²

Methods Retrospective cohort study of all BALs performed in ILD at the University Hospital of South Manchester (UHSM) between January 2015 and November 2016. Electronic reports and histopathological cellular analysis were assessed for qualitative outcomes as per BAL guidelines.²

Results 416 patients' coded as BAL sampling. 95 (24%) were performed in the diagnostic strategy for ILD. The mean age was 58.9 (range 18–89). 36 (37.9%) were Female and 59 (62.1%) were male. 13 (13.7%) had lymphocyte count $\geq 15\%$,