in patients without evidence of trapped lung. However, talc pleurodesis is unsuccessful in 10%–40% of cases, with implications in relation to cost and morbidity.¹ Yet ambiguity exists regarding the factors affecting success in MPE talc pleurodesis. **Aim** This study aims to investigate the predictive factors relating to successful talc pleurodesis in MPE and the impact of a positive outcome on patient mortality.

Methodology Retrospective analysis of patients admitted for management of MPE to the Belfast City Hospital between September 2015 – October 2016 was conducted. Demographic and clinical data relating to drain size, volume of fluid drained and grade of the doctor performing talc pleurodesis was collected. Survival at 18 months post procedure was reviewed through electronic patient records. A positive outcome was defined as successful pleurodesis with the lack of recurrence of pleural effusion.

Results Twenty-seven patients were identified (♂/♀: 40.8/59.2%; age 72.4+/-12.3 years). Two thirds (n=18) received pleurodesis with a 44.4% success rate (n=8). Pleurodesis was precluded in one third of patients (n=9) with displaced and blocked drains the predominant causative factor (n=4). In those receiving talc pleurodesis, drain size (12 F vs 18 F) was not a predictive factor of positive outcomes. Similarly, the grade of doctor performing talc pleurodesis did not affect efficacy. However, compared to 18 F drains, 12 F drains were associated with a significant complication rate precluding pleurodesis (p=0.02). Critically, achieving a successful outcome with talc pleurodesis was associated with improved 18 month mortality (p=0.004).

Conclusion Promoting the exclusive utilisation of 18 F drains in the management of MPE could potentially alleviate the propensity for intercostal drain failure to preclude talc pleurodesis, conceivably improving patients' short term mortality.

REFERENCE

1. Santos PS, Marques MA, Cruz C, Monteiro H, Fradinho F. Predictors of talc slurry pleurodesis success in patients with malignant pleural effusions. *Revista Portuguesa de Pneumologia* 2017;23(4):216–220.

P233

THORACIC ULTRASONOGRAPHY AS A PREDICTOR OF PLEURODESIS SUCCESS IN MALIGNANT PLEURAL EFFUSION

JP Corcoran, RJ Hallifax, A Yousuf, RM Mercer, R Asciak, M Hassan, I Psallidas, NM Rahman. *Oxford Centre for Respiratory Medicine, Oxford University Hospitals NHS Foundation Trust, Oxford, UK*

10.1136/thoraxjnl-2017-210983.375

Background Over 50 000 patients with malignant pleural effusion (MPE) are seen annually in the UK. The majority develop recurrent symptomatic disease requiring definitive treatment. MPE is most frequently managed with talc slurry pleurodesis via intercostal chest drain. This involves a lengthy inpatient stay and has a success rate of around 70%, with no means of predicting which patients will suffer pleurodesis failure. Thoracic ultrasound (TUS) is widely used by respiratory physicians, and data from animal and human studies suggest it can identify pleural adhesions (through the absence of normal lung sliding) in a range of conditions. By extension, TUS may allow clinicians to diagnose the presence or absence of adhesions post-pleurodesis in MPE, identifying patients suitable for discharge or needing further intervention.

Method We recruited 18 adult patients with MPE undergoing drainage and talc slurry pleurodesis to a prospective single-centre cohort study. Patients underwent standardised TUS assessment pre- and post-pleurodesis, evaluating pleural sliding and adhesions at nine points (three anterior, three lateral, three posterior) across the affected hemithorax. Lung sliding was graded as per Zhu *et al.*,¹ creating a total pleurodesis score out of 18. Pleurodesis failure was defined as radiological and symptomatic fluid recurrence in the same hemithorax requiring further intervention at any point up to 3 months post-pleurodesis. Patients also completed a questionnaire addressing satisfaction with TUS assessment.

Results 3/18 patients (16.7%) died before 1 month follow-up. Of 15 patients seen at one month, 11 (73.3%) had successful pleurodesis and 4 (26.7%) had failed. No patient had delayed pleurodesis failure between 1 and 3 month follow-up. There was a significant difference observed in the day 1 TUS pleurodesis score between patients who went on to have successful pleurodesis and those who failed during follow-up (table 1). TUS assessment was acceptable to patients, with none considering it either time-consuming or unwilling to have it again if needed.

Conclusion Our data suggest TUS assessment 24 hours post-pleurodesis for MPE predicts success or failure of this intervention, with significant implications for clinical care. A larger randomised study is now underway to further evaluate this hypothesis.

REFERENCE

1. *Chest* 2005;128(2):934–9.

Abstract P233 Table 1 Ultrasonographic pleurodesis score at day 0 (pre-pleurodesis) and day 1 (24 hours post-pleurodesis) in patients being treated for malignant pleural effusion

	Successful pleurodesis n=11/15 (73.3%) patients	Failed pleurodesis n=4/15 (26.7%) patients	p value unpaired t- test
Day 0 pleurodesis score (mean ±SD, total out of 18)	10.89±3.98	6.50±1.29	0.054
	Difference=4.39 (95% CI -0.09 to 8.86)		
Day 1 pleurodesis score (mean±SD, total out of 18)	13.45±2.63	6.75±2.94	0.002
	Difference=6.70 (95% CI 3.08 to 10.33)		
Change from day 0 to 1 (mean±SD)	2.57±3.98	0.25±3.59	0.326
	Difference=2.32 (95% CI -2.59 to 7.23)		

P234

PATIENT AND FLUID CHARACTERISTICS ASSOCIATED WITH NON-DRAINING MALIGNANT PLEURAL EFFUSION

¹EK Mishra, ²A Clive, ³HE Davies, ³AJ Nunn, ⁴RF Miller, ⁵I Psallidas, ¹NA Maskell, ⁵NM Rahman. ¹Norfolk and Norwich University Hospital, Norwich, UK; ²University of Bristol, Bristol, UK; ³Cardiff and Vale University Health Board, Cardiff, UK; ⁴University College London, London, UK; ⁵University of Oxford, Oxford, UK

10.1136/thoraxjnl-2017-210983.376

Introduction TIME3, a randomised controlled trial of intra-pleural urokinase versus placebo for patients with non-draining malignant pleural effusion (MPE), demonstrated that these patients appear to be a distinct subgroup of patients with a poor prognosis (median survival 58 days). The aim of this study was to identify patient and fluid characteristics

associated with this subgroup, to enable further understanding of why patients with non-draining effusion may have poor prognosis.

Methods Baseline demographics and pleural fluid (PF) characteristics of patients enrolled in TIME3 were compared to patients enrolled in TIME2,² a randomised controlled trial of indwelling pleural catheter versus chest drain and pleurodesis for patients with recurrent MPE. Demographic characteristics compared were: age, sex, histological type of cancer and ECOG performance status (PS). Pleural fluid characteristics compared were: total protein, glucose, cytology (positive or negative), pH, lactate dehydrogenase (LDH) and presence of septations on ultrasound. These characteristics were compared using t test for linear variables and chi squared for categorical variables.

Results The median survival was 58 days (IQR 27–123) in TIME3 versus 187 days (IQR 48–358) in TIME2. Patients with non-draining effusions had a significantly higher PF LDH (mean 1900 (SD 3100) versus 660 (SD 840), $p<0.001$) and CRP (mean 117 (SD 80) versus 62 (SD 55), $p<0.001$). Patients in TIME3 were on average 4 years older (mean 71 years in TIME3 versus 67 in TIME2, $p=0.01$) and less likely to be cytology positive (24% versus 51%, $p=0.021$).

Conclusion Non draining MPEs have a higher LDH than those without. There was a large difference in mortality between groups, but despite this no identifiable differences in baseline ECOG, PS or tumour type, despite these variables being associated with a poor prognosis in unselected cohorts of patients with MPE.¹ We postulate that survival in MPE may be associated with septations and the intrapleural inflammatory milieu. Further study of the association between PF LDH, septations and survival is warranted.

REFERENCES

1. Clive, A. O. Predicting survival in malignant pleural effusion: Development and validation of the LENT prognostic score. *Thorax* 2014;69(12):1098–104.
2. Davies HE, M. E. TIME2 randomised controlled trial. *Journal of the American Medical Association* 2012;2383–9.

Abstract P234 Table 1

Variable	TIME2	TIME3	Difference (p)
Number of patients	106	71	-
Mean age (years) (SD)	67 (11)	71 (9.4)	0.013
Male:female (% male)	46:60 (43)	41:30 (58)	0.06
Median time to death (days) (IQR)	187 (48–358)	58 (27–123)	
Type of cancer (%)			
- breast	27 (25)	12 (17)	0.47
- lung	25 (24)	22 (31)	
- mesothelioma	11 (10)	9 (13)	
- other	43 (41)	28 (39)	
ECOG PS 0–2:3–4 (% 0–2)	60:46 (57)	42:29 (59)	0.11
Bloods			
White cell count	9.8 (5.4)	11.0 (5.4)	0.15
CRP	62 (55)	117 (80)	<0.001
Pleural fluid characteristics:			
- Cytology positive: negative (% positive)	54:48 (51)	17:34 (24)	0.021
- pH (SD)	7.4 (0.24)	7.4 (0.34)	1.0
- mean glucose (mmol/L) (SD)	5.4 (2.8)	4.6 (3.6)	0.13
- LDH (U/L) (SD)	660 (840)	1900 (3100)	<0.001
- Total protein (g/dL) (SD)	43 (8.1)	41 (10)	0.18
Septated on ultrasound (yes:no)	Not recorded	59:7	-

P235

ASSESSMENT OF DIAPHRAGM MOTION IN PATIENTS WITH UNILATERAL OR ASYMMETRICAL PLEURAL EFFUSIONS

¹MG Aldik, ¹A Sibly, ²L Telisinghe, ¹C Daneshvar. ¹Department of Respiratory Medicine, Plymouth Hospitals NHS Trust, Plymouth, UK; ²Department of Clinical Research, Faculty of Infectious and Tropical Diseases, London School of Hygiene and Tropical Medicine, London, UK

10.1136/thoraxjnl-2017-210983.377

Introduction The diaphragm is the most important respiratory muscle. In patients with pleural effusions, abnormal diaphragm shape and movement on inspiration may be observed. We aimed to explore the possible significance of these findings in a our pleural service.

Methods Between August 2015 and June 2016, all patients with an index thoracic ultrasound scan reporting a unilateral or significantly asymmetrical bilateral pleural effusion were retrospectively assessed. Patients were examined in the upright position by an RCR level 1 trained physician. A routine standardised reporting form was used to record dimensions, echogenic properties, the presence of septations, pleural nodularity and diaphragm nodularity, shape and movement. Pleural fluid characteristics and final diagnoses were assessed.

Results Of 491 patients assessed, 259 met the inclusion criteria. Thoracentesis was performed in 70%, with exudative defined by Light's criteria being met in 121 (47%). A diagnosis was established in 84% of cases with the commonest aetiology being malignancy (32%). The median [interquartile range] size of the pleural effusions was 8 [5–10] cm in medial depth and 3 [2–4] rib spaces in height, with 118 (46%) being echogenic. Diaphragm assessment revealed that paradoxical movement of the diaphragm was observed in 56 (22% [95% CI 16.6%–26.6%]) of patients while 11 (4.3% [95% CI 1.8%–6.7%]) had an inverted diaphragm. Patients with paradoxical diaphragm movement had larger effusions (median depth 11 cm versus 7 cm; $p<0.001$) and a higher proportion of effusions of malignant aetiology (45% versus 11%; $p<0.001$) when compared with individuals with normal diaphragm movement. In multivariate analysis, adjusting for age, gender, effusion depth and height and diaphragm shape, effusions of malignant aetiology were associated with an increased odds of paradoxical diaphragm movement (adjusted odds ratio 4.47 (95% CI 1.83–10.95; $p<0.001$).

Conclusion Paradoxical diaphragm movement in the context of a unilateral or asymmetrical pleural effusion is frequently observed, is independently associated with malignant pleural effusions and may be a useful point of care clinical sign.

P236

A SYSTEMATIC REVIEW OF INTERVENTIONS TO IMPROVE HEALTH RELATED QUALITY OF LIFE IN MALIGNANT PLEURAL EFFUSION

¹P Sivakumar, ²A Saigal, ²L Ahmed. ¹St Thomas' Hospital, London, UK; ²Whittington Hospital, London, UK

10.1136/thoraxjnl-2017-210983.378

Introduction Malignant pleural effusion (MPE) Results in breathlessness and impairment of health related quality of life (HRQOL). Despite this there is a lack of consensus on the ideal treatment strategy to improve HRQOL.¹