(5.0% vs 13.8%; $p=0.03$). PD-L1 expression in the cohort was high ($\geq 50\%$) in 28.5%, weak ($\geq 1\%$ –50%) in 28.2%, whilst 43.3% of patients were PD-L1 negative. The only statistically significant predictor for PD-L1 expression in multivariate analysis was the presence of brain metastasis at diagnosis (OR 2.02; CI 1.04–3.90). 47 patients (11.4%) were treated with immunotherapy and the response rate was 16.2%. All patients that responded to immunotherapy had high ($\geq 50\%$) expression of PD-L1.

Conclusions This large multicentre study demonstrates for the first time that samples obtained by EBUS-TBNA in routine practice are suitable for PD-L1 testing in patients with NSCLC. The presence of brain metastases at diagnosis predicts high PD-L1 expression in this cohort and this new finding should be tested in future clinical trials.

REFERENCE

1. Reck M, Rodríguez-Abreu D, Robinson AG, *et al.* Pembrolizumab versus Chemotherapy for PD-L1-Positive Non-Small-Cell Lung Cancer. *N Engl J Med* 2016;375 (19):1823–33. doi:10.1056/NEJMoa1606774

S101

A COMPARISON OF THE IMAGING FEATURES OF EARLY STAGE PRIMARY LUNG CANCER IN PATIENTS TREATED WITH SURGERY, SABR AND MICROWAVE ABLATION

¹A Talwar, ¹N Jenko, ¹M Sarim, ²M Enescu, ²P Whybra, ²JMY Willaime, ³LC Pickup, ¹W Hickes, ²M Gooding, ²D Boukerroui, ¹NM Rahman, ³T Kadir, ¹FV Gleeson. ¹Oxford University Hospitals NHS Foundation Trust, Oxford, UK; ²Mirada Medical Ltd, Oxford, UK; ³Optellum Ltd, Oxford, UK

10.1136/thoraxjnl-2017-210983.107

Introduction Stereotactic Ablative Radiotherapy (SABR) and percutaneous microwave ablation (PMWA) are now being performed in patients deemed “medically inoperable” with non-small cell lung cancer (NSCLC). The majority of these patients are treated without ground truth histology, relying on imaging to establish the diagnosis. The purpose of this study was to

investigate whether there were differences in the visible imaging features including CT Texture Analysis (CTTA) between patients referred for surgery, SABR and PMWA, which might suggest differences in underlying diagnosis.

Methods 92 patients with one pulmonary nodule (PN) suspected as T1N0M0 to T2AN0M0 NSCLC on imaging were treated either with SABR (22 patients), PMWA (25) or Video-assisted thorascopic surgery (45) of which 23 had NSCLC (SURG M) and 22 had benign disease (SURG B). Patient characteristics, CT nodule morphology, presence of emphysema and percentage emphysema score, FDG avidity and CT textural features were compared. Twenty texture features (previously used in combination to create a nodule probability of malignancy score between 0–1) were extracted from each automatic contoured region surrounding the PN. The Kruskal-Wallis test was used to compare texture features between the 4 patient groups (SABR, PMWA, SURG M and SURG B).

Results There was no significant difference in nodule morphology, volume at presentation ($p=0.280$) or volume doubling times ($p=0.149$), and presence of emphysema ($p=0.348$) or emphysema score ($p=0.367$) between the 4 groups. There was no statistical difference in CTTA malignancy prediction score between the SABR, PMWA and SURG M groups ($p\geq 0.05$). The probability of malignancy score was significantly lower ($p\text{-value}<0.01$) for SURG B (0.58 mean ± 0.19 sd) vs. SABR (0.79 ± 0.15) treatment groups (figure 1).

Conclusion This is the first study to our knowledge to evaluate the radiological differences between patient groups referred for surgical and non-surgical treatments for NSCLC. On this small study, the Results support the hypothesis that the non-operative patient groups comprise the same proportion of benign and malignant as those in the operative group. The Results also demonstrate the potential clinical utility of CTTA in patient selection when histology is not obtainable. CTTA does not require volumetry detectable growth to detect change, and therefore may be a useful biomarker of malignancy at first diagnosis.

Abstract S101 Figure 1 Summary statistics for probability of malignancy scores.

Fruit flies to footballers

S102 IDENTIFICATION OF ALLERGENS PRESENT IN DROSOPHILA MELANOGASTER USING A SERUM IMMUNOBLOTTING METHOD

M Brian, D Jarvis, P Burney, P Cullinan, M Jones. *Population Health and Occupational Disease, National Heart and Lung Institute, Imperial College London, London, UK*

10.1136/thoraxjnl-2017-210983.108

Background *Drosophila melanogaster*, otherwise known as the fruit fly is commonly used in laboratory animal research. We have reported (Jones, Blair *et al.* 2017) that laboratory workers (n=286) exposed to fruit flies have an overall sensitisation prevalence of 6%, based on measuring allergen-specific IgE by radioallergosorbent test to a whole fruit fly extract. Although IgE binding to fruit fly extract was detected, the allergenic proteins responsible for IgE binding have not been identified.

Objective To identify allergenic proteins from a fruit fly extract

Methods Fruit flies were collected from the workplace, extracted overnight in 0.01 mol/L ammonium carbonate at 4°C, dialysed against distilled water and lyophilised. SDS-PAGE was used to separate 150 µg of the extract according to protein molecular weight. Extracted proteins binding to serum specific IgE were detected with Western blotting, using 50 µl of sera from fruit fly sensitised workers (n=3). An alkaline phosphatase conjugated mouse anti-human IgE secondary antibody and NBT/BCIP chromogenic substrate were used in detection. Images were acquired with a ChemiDoc MP and

Abstract S102 Figure 1 Western blot detection of serum specific IgE binding proteins from 150µg of fruit fly extract. 50 µl of three individual sera, an alkaline phosphatase conjugated mouse anti-human IgE secondary antibody and NBT/BCIP chromogenic substrate were used to detect proteins. Blots were also performed with pooled sera from the three individuals and with no sera (control).

the molecular weight of allergenic proteins determined with a 10–250 kDa prestained protein ladder (ThermoScientific) on ImageLab software (v5.2.1).

Results From the fruit fly extract, six distinct proteins binding to serum specific IgE were observed (figure 1). In three sensitised workers, IgE binding to proteins with molecular weights of ~107 and 76 kDa was observed. For one individual, IgE binding was present to an additional four proteins from the fruit fly extract, with molecular weights of ~183, 54, 28 and 12 kDa. In a control blot, we did not observe any non-specific binding to the fruit fly extract.

Conclusions There are at least six distinct proteins from a fruit fly extract with IgE binding properties of an allergen. Currently, these proteins are not characterised as allergens in protein databases. We will carry out further proteomic testing to characterise these unknown allergens.

S103 OCCUPATIONAL ALLERGY TO FRUIT FLIES (DROSOPHILA MELANOGASTER) IN LABORATORY WORKERS

¹M Jones, ²S Blair, ¹S MacNeill, ¹J Welch, ¹A Hole, ³P Baxter, ¹P Cullinan. ¹Imperial College, London, UK; ²Cambridge University Hospital NHS Foundation Trust, Cambridge, UK; ³University of Cambridge, Cambridge, UK

10.1136/thoraxjnl-2017-210983.109

Introduction and Objectives *Drosophila melanogaster* (the ‘fruit fly’) is commonly used in genetic research, but there is only one earlier report of immunoglobulin E-associated allergy in exposed workers. Four newly identified cases prompted us to examine the extent of this problem in a university laboratory. Our aim was to determine the prevalence and determinants of sensitisation to fruit flies in a population of exposed workers.

Methods In a cross sectional study we surveyed two hundred and eighty six employees working in a department carrying out research involving *D. melanogaster*. Sensitisation was assessed by specific IgE measurement in serum using radioallergosorbent assay (RAST) and examined in relation to work-related symptoms and to estimated exposure to fruit flies.

Results The overall prevalence of specific sensitisation was 6% with a clear relationship to increasing frequency/intensity of exposure (p trend <0.001). Work-related eye/nose, chest or skin symptoms were reported by substantial proportions of participants but for most of these there was no evidence of specific sensitisation to fruit fly. The overall prevalence of any work related symptoms and sensitisation was 2.4%, rising to 7.1% in those working in high exposure groups.

Conclusions We were able to demonstrate, for the first time, a clear exposure-response relationship between fruit fly exposure and specific sensitisation. Facilities housing fruit flies should carefully consider methods to reduce exposure levels in the workplace.

S104 INVESTIGATING THE DIAGNOSTIC PERFORMANCE OF SPECIFIC IMMUNOLOGICAL TESTS IN OCCUPATIONAL ASTHMA

D Fernandes, J Cannon, B Fitzgerald, J Welch, M Jones, P Cullinan, J Szram. *Imperial College London, London, UK*

10.1136/thoraxjnl-2017-210983.110

The detection (or otherwise) of specific IgE sensitisation is an important tool in the investigation of employees with potential