ORIGINAL ARTICLE

High prevalence of *CCDC103* p.His154Pro mutation causing primary ciliary dyskinesia disrupts protein oligomerisation and is associated with normal diagnostic investigations

Amelia Shoemark,¹ Eduardo Moya,² Robert A Hirst,³ Mitali P Patel,⁴ Evelyn A Robson,² Jane Hayward,^{4,5} Juliet Scully,^{4,6} Mahmoud R Fassad,^{4,7} William Lamb,⁴ Miriam Schmidts,^{8,9} Mellisa Dixon,¹ Ramila S Patel-King,¹⁰ Andrew V Rogers,^{1,11} Andrew Rutman,³ Claire L Jackson,^{12,13} Patricia Goggin,^{12,13} Bruna Rubbo,^{12,13} Sarah Ollosson,¹ Siobhán Carr,¹ Woolf Walker,^{12,13} Beryl Adler,¹⁴ Michael R Loebinger,¹¹ Robert Wilson,¹¹ Andrew Bush,^{1,15} Hywel Williams,¹⁶ Christopher Boustred,⁵ Lucy Jenkins,⁵ Eamonn Sheridan,¹⁷ Eddie M K Chung,¹⁸ Christopher M Watson,¹⁷ Thomas Cullup,⁵ Jane S Lucas,^{12,13} Priti Kenia,¹⁹ Christopher O'Callaghan,^{3,20} Stephen M King,^{10,21} Claire Hogg,¹ Hannah M Mitchison⁴

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/thoraxjnl-2017-209999>).

For numbered affiliations see end of article.

Correspondence to

Dr Hannah M Mitchison, PUW32, Experimental and Personalised Medicine Section, Genetics and Genomic Medicine Programme, UCL Great Ormond Street, Institute of Child Health, London WC1N 1EH, UK; h.mitchison@ucl.ac.uk

Received 16 January 2017

Revised 7 June 2017

Accepted 3 July 2017

Published Online First

8 August 2017

ABSTRACT

Rationale Primary ciliary dyskinesia is a genetically heterogeneous inherited condition characterised by progressive lung disease arising from abnormal cilia function. Approximately half of patients have situs inversus. The estimated prevalence of primary ciliary dyskinesia in the UK South Asian population is 1:2265. Early, accurate diagnosis is key to implementing appropriate management but clinical diagnostic tests can be equivocal.

Objectives To determine the importance of genetic screening for primary ciliary dyskinesia in a UK South Asian population with a typical clinical phenotype, where standard testing is inconclusive.

Methods Next-generation sequencing was used to screen 86 South Asian patients who had a clinical history consistent with primary ciliary dyskinesia. The effect of a *CCDC103* p.His154Pro missense variant compared with other dynein arm-associated gene mutations on diagnostic/phenotypic variability was tested. *CCDC103* p.His154Pro variant pathogenicity was assessed by oligomerisation assay.

Results Sixteen of 86 (19%) patients carried a homozygous *CCDC103* p.His154Pro mutation which was found to disrupt protein oligomerisation. Variable diagnostic test results were obtained including normal nasal nitric oxide levels, normal ciliary beat pattern and frequency and a spectrum of partial and normal dynein arm retention. Fifteen (94%) patients or their sibling(s) had situs inversus suggesting *CCDC103* p.His154Pro patients without situs inversus are missed.

Conclusions The *CCDC103* p.His154Pro mutation is more prevalent than previously thought in the South Asian community and causes primary ciliary dyskinesia that can be difficult to diagnose using pathology-based clinical tests. Genetic testing is critical when there is a strong clinical phenotype with inconclusive standard diagnostic tests.

Key messages

What is the key question?

- Can gene sequencing improve diagnosis of the inherited respiratory condition primary ciliary dyskinesia (PCD) in patients with unclear clinical diagnostic investigations?

What is the bottom line?

- *CCDC103* p.His154Pro missense mutations cause up to 20% of PCD cases in South Asian populations but diagnosis can be difficult in this group using standard clinical diagnostic tests because results are often normal; we therefore propose that genetic analysis is an essential part of the diagnostic algorithm to complement standard clinical tests to improve diagnostic accuracy.

Why read on?

- Patients with PCD may be missed using current pathology-based diagnostic protocols therefore genetic screening can provide valuable support in obtaining a definitive diagnosis.

INTRODUCTION

Primary ciliary dyskinesia (PCD; OMIM: 244400) is an inherited disorder affecting motile cilia. Patients usually present with a history of neonatal respiratory distress and suffer from lifelong symptoms of chronic wet cough and rhinitis. Recurrent chest infections ultimately lead to bronchiectasis and a progressive decline in lung function.¹ Approximately half of patients have situs inversus and other situs abnormalities, due to randomisation of situs determination caused by ciliary dysmotility in the embryonic node.² Fertility can also be affected by

► <http://dx.doi.org/10.1136/thoraxjnl-2017-210776>

CrossMark

To cite: Shoemark A, Moya E, Hirst RA, et al. *Thorax* 2018;**73**:157–166.

defective cilia in the fallopian tubes and non-motile sperm tail flagella.

The estimated prevalence of PCD in the UK is 1:15 000, but as high as 1:2265 in the UK South Asian population.¹³ Early diagnosis is important to maintain lung function, and appropriately treat symptoms to reduce morbidity and mortality.⁴ Diagnosis can be complex and requires a combination of tests for cilia functional and ultrastructural defects.^{5,6} PCD is caused by mutations in genes encoding proteins conferring structure and motility to the cilia and governing ciliogenesis. It is genetically heterogeneous with >200 individual mutations in more than 30 genes known to cause PCD. To date, these account for approximately 65% of cases.^{17–11} UK Genetic Testing Network approved tests are offered in two centres (http://www.labs.gosh.nhs.uk/media/764464/ciliopathies_v8.pdf).¹²

CCDC103 mutations were first reported, in 2012, in patients with dynein arm loss and a typical clinical PCD phenotype.¹³ *CCDC103* is an oligomeric coiled-coil domain protein that is found tightly bound to the ciliary axoneme where it is thought to help facilitate ciliary motility by participating in attachment of the dynein arms to the axoneme. The protein was found to stabilise cytoplasmic microtubules against cold depolymerisation in an in vitro assay.¹³ A missense variant previously identified in *CCDC103* to cause a single amino acid change to the protein, p.His154Pro (rs145457535) was previously described as a hypomorphic mutation since mutant p.His154Pro *CCDC103* induced intermediate partially rescued disease phenotypes when expressed in a zebrafish *CCDC103*-null model, suggesting some protein function was retained.¹³ In cilia from p.His154Pro-mutated patients some cilia showed a partial dynein arm defect with the outer dynein arms (ODA) at least partially assembled, compared to loss-of-function *CCDC103* mutations that cause complete ODA loss. In agreement with this, loss-of-function *CCDC103* mutation patients had largely static cilia, while patients carrying a homozygous p.His154Pro mutation exhibit a mixed (static and motile) ciliary beat pattern.¹³

We conducted genetic screening of 86 patients with PCD of South Asian, mostly Pakistani, origin, detecting that a significant proportion (19%) were homozygous for *CCDC103* p.His154Pro variant. Among these, many were at high risk of being undiagnosed without genetic testing, due to normal diagnostic results obtained in PCD investigations. Using electron microscopy and protein biochemistry we have sought to further determine the pathogenic nature of the p.His154Pro mutation.

METHODS

Patient selection

Eighty-six patients of South Asian (primarily Pakistani) descent with clinical signs and symptoms of PCD were identified from the UK National PCD Diagnostic and Management Services at The Royal Brompton Hospital, London, University Hospital Southampton, Birmingham Children's Hospital, Bradford Royal Infirmary and Leicester General Infirmary.¹⁴

Genetic screening

All participants gave written informed consent to take part in this study. The protocol was approved by the London Bloomsbury Research Ethics Committee (08/H0713/82). High throughput screening used next-generation sequencing, either whole exome sequencing (WES) or targeted gene panel sequencing. Sequencing and variant identification methods are published for WES^{12,15} or used custom gene panels (Illumina TruSeq Custom Amplicon or Agilent SureSelect Focused Exome ('clinical exome' proprietary

product) and SureSelectXT custom panel design systems) with a standardised variant calling pipeline.¹⁶ Sanger sequencing was used for variant confirmation and familial segregations. As shown in table 1, of the 16 p.His154Pro cases, this mutation was detected by WES in case numbers 1–6 and number 9, by use of a 'clinical exome' commercial panel in case number 8 and in case numbers 7, 10–16 by use of custom targeted gene panels containing the known PCD genes and other candidate PCD disease genes.

Comparator group

The comparator patient group consisted of 16 of the 86 individuals tested. This group was closely age and gender matched to the *CCDC103* p.His154Pro group. All had a dynein arm defects on electron microscopy and all were proven negative for the *CCDC103* p.His154Pro mutation.

FEV₁ measurements

Spirometry was performed according to American Thoracic Society/European Respiratory Society recommendations.¹⁷ Forced expiratory volume in 1 second (FEV₁, the standard lung airflow measure) z-scores were calculated using the Global Lungs Initiative parameters.¹⁸

Diagnostic tests

Screening and diagnostic testing was performed according to the PCD National Service protocols. Investigations included nasal nitric oxide (nNO), nasal brush biopsies analysed by high speed video imaging (HSVM) for ciliary beat frequency and pattern¹⁹ and quantitative electron microscopy for ciliary ultrastructure.²⁰ Additional detail on the method is provided in an online supplementary file. When results were inconclusive or inconsistent, patients were offered repeat testing.

Protein biochemistry on recombinant *CCDC103* protein

Site-directed mutagenesis (QuikChange kit, Agilent Technologies, UK) was used to generate the p.His154Pro mutation in an N-terminal His₁₀-tagged *Homo sapiens CCDC103* cDNA subcloned into pET16b vector that was synthesised using *Escherichia coli* codon bias.²¹ Following transformation into *E. coli* BL21 (DE3), protein expression was induced by addition of 2 mM isopropyl β-D-thiogalactoside (IPTG) for 2 or more hours. Following sonication, His₁₀-tagged proteins were dissolved in 8 M urea, then very slowly refolded by their dilution into 1 L of 20 mM Tris-Cl pH8.0 150 mM NaCl. Proteins were then purified by Ni²⁺-affinity chromatography as described previously using 20 mM Tris-Cl pH 8.0, 500 mM NaCl, 250 mM imidazole for elution.²¹ Samples concentrated by ultrafiltration through Amicon Ultra-4 ultrafiltration units were subject to gel filtration in a calibrated Superose 6 10/300 column attached to an Äkta-Purifier-10 chromatography workstation.²¹ The mutant protein was very hard to make and only a little was able to be refolded, indeed even the precipitated material was clearly different from the wild-type protein being more 'sticky.' Hence, a lower concentration of the mutant protein is apparent in figure 1. This experiment was done at one concentration (the highest we were able to achieve): for wild-type protein this was ~80 µg/mL and for the p.His154Pro protein this was ~30 µg/mL.

Statistical analysis

The *CCDC103* p.His154Pro and comparator patient groups were closely matched for age and gender. Data were not normally distributed and therefore groups were compared using

Table 1 Clinical history of *CCDC103* p.His154Pro cases summarising diagnostic investigations

Demographics			Symptoms		Diagnostic test results								
ID	Sex	Age (years)	Age diag (years)	Family history	FEV ₁ z-score	Main symptoms	Situs	Nasal NO (nL/min)	CBF (Hz)	Ciliary beat pattern	Electron microscopy (first brushing)	Electron microscopy (repeat brushing)	Genetics: sequencing protocol and segregations
1	M	7	7	No	-3.45	Chronic cough, developmental delay	SI	ND	0	Immotile	IDA +ODA	IDA +ODA	Watson <i>et al</i> ¹²
2	M	4	1	Sib 3	-2.25	Nasal discharge	SI	1	0.95	Mostly immotile with occasional residual movement	IDA +ODA	IDA +ODA	Watson <i>et al</i> ¹²
3	M	12	10	Sib 2	-2.75	Neonatal respiratory distress, nasal discharge	SS	11	2.9	Mostly immotile with occasional residual movement	IDA +ODA	IDA +ODA	Watson <i>et al</i> ¹²
4	M	13	<1	No	ND	Neonatal respiratory distress, Gilbert syndrome, epilepsy	SI	7	5.9	Mostly immotile with occasional residual movement	IDA +ODA	IDA +ODA	Watson <i>et al</i> ¹²
5	M	13	6	Sib 6	-1.52	Recurrent chest infections	SI	87	5.79	Normal ciliary beat pattern reduced frequency	IDA +ODA	IDA +ODA	WES ^P
6	F	15	4	Sib 5	-0.61	Recurrent chest infections, Eustachian tube dysfunction, nasal discharge	SS	43	7.66	Normal ciliary beat pattern reduced frequency	IDA +ODA	IDA +ODA	WES ^P
7	F	18	9	No	-1.58	Recurrent chest infections, nasal discharge, bilateral glue ear	SI	111	10.6	Mixed Sample 1: normal ciliary beat pattern Sample 2: immotile	IDA +ODA	IDA +ODA	Illumina TruSeq custom gene panel ^M
8	M	9	6	No	-0.08	Wet cough, recurrent chest infections, conductive hearing loss	SI	57	8.8	Dyskinesia	IDA	IDA	Agilent SureSelect Focused Exome ^P
9	F	7	7	CHD	0.43	Chronic cough	SI	33	10.9	Normal ciliary beat pattern	Normal	ND	Watson <i>et al</i> ¹²
10	M	10	<1	Sib 11 and 12	-1.32	Neonatal respiratory distress, nasal discharge	SI	26	11.5	Mixed: normal areas, immotile areas, dyskinetic areas	Normal	Normal	Illumina TruSeq custom gene panel ^B
11	F	21	10	Sib 10 and 12	ND	Recurrent chest infections, asthma, persistent collapse of right lower lobe	SI	293	8.5	Mixed: mostly reduced forward and recovery stroke	IDA	Normal	Illumina TruSeq custom gene panel ^B
12	M	13	<1	Sib 10 and 11	-1.24	Neonatal respiratory distress, nasal discharge	SS	214	16.3	Normal ciliary beat pattern	IDA	Normal	Illumina TruSeq custom gene panel ^B
13	F	29	29	No	ND	Bronchiectasis, infertility	SI	239	4.31	Mixed: mostly reduced forward and recovery stroke Static patches	Normal	Insufficient	Agilent SureSelectXT ^B Cillome_651
14	F	1	1	No	ND	Neonatal respiratory distress, nasal discharge, wet cough	SI	ND	16.76	Dyskinesia	Normal	Inconclusive	Agilent SureSelectXT Cillome_651
15	F	21	21	Sib 16	ND	Recurrent chest infections in childhood. No ear, chest or nasal symptoms in adulthood	SI	151	10.9	Normal ciliary beat pattern	IDA	Normal	Agilent SureSelectXT ^M Cillome_651
16	M	18	18	Sib 15	ND	Sensory neural hearing impairment, recurrent chest infections, chronic nasal congestion and rhinitis	SI	276	9.9	Mixed 80% normal ciliary beat pattern 20% immotile cilia on strips	IDA	Normal	Agilent SureSelectXT Cillome_651 ^M

Normal test results are highlighted in bold. Normal range for nNO is above 77 nL/min. Local normal range for CBF is 8.5-16.8 Hz.

Segregation analysis using available parental samples indicated by M (mother is carrier), P (father is carrier) or B (both parents carriers), unaffected siblings have not been tested.

Age diag, age diagnosis confirmed; CBF, ciliary beat frequency; CHD, congenital heart defects; F, female; IDA, inner dynein arms; M, male; ND, not done; NO, nitric oxide; ODA, outer dynein arms; SI, situs inversus; Sib, sibling of; SS, situs solitus; WES, whole exome sequencing.

Figure 1 Comparison of predicted FEV₁ in *CCDC103* p.His154Pro cases versus a comparator group. The *CCDC103* p.His154Pro cases are described in [table 1](#) and the comparator group are South Asian origin *CCDC103* p.His154Pro-negative cases with a confirmed absent dynein arms defect. Mutations carried by the comparator group are in *DNAAF1* (three cases), *DNAAF3* (two cases), *DNAH5* (two cases), *LRRC6* (three cases), *ZMYND10* (two cases), while four cases were excluded from *CCDC103* p.His154Pro but genetically undefined.

non-parametric statistical tests. $p < 0.05$ was considered statistically significant.

RESULTS

A PCD-causing mutation *CCDC103* p.His154Pro is prevalent in UK individuals of South Asian origin with a clinical phenotype of PCD

Next-generation sequencing analysis in 86 patients of South Asian descent with suspected PCD revealed 16 patients (19%) from 11 independent families homozygous for a previously published single base change (NM_001258395.1: c.461A>C) mutation in *CCDC103* that predicts the amino acid substitution p.His154Pro ([table 1](#)). Consistent with their homozygous segregation pattern, all 16 patients were children of consanguineous parents and had dynein arm defects or normal ultrastructure.

Due to the high frequency of this mutation, we examined whole exome sequence data available from 1542 unaffected parents with similar ethnic backgrounds participating in the Born-in-Bradford study, all of UK South Asian, primarily Pakistani heritage.²² This revealed six heterozygous carriers of the *CCDC103* p.His154Pro substitution (E Sheridan, unpublished data). The ExAc database²³ of exome sequencing results from 60 706 unrelated control individuals records an allele frequency for p.His154Pro three times as high in 8256 South Asian individuals (0.003) compared with 33,345 North Europeans (0.001). No p.His154Pro homozygote individuals were identified in the entire Born-in-Bradford or ExAc control cohorts.

Among the 16 *CCDC103* p.His154Pro homozygote PCD patients identified, normal diagnostic test results were apparent as highlighted in bold in [table 1](#). Although PCD was strongly suspected in all cases, 5 of the 16 patients (case 9 and 13–16 in [table 1](#)) did not receive a definitive disease diagnosis until their genotype was confirmed. Remarkably in the case of patient 15, this individual had been deemed to not have PCD and was discharged from respiratory care. She was retested due to her situs inversus and the finding that her brother, who remained under clinical suspicion, had this *CCDC103* mutation. Thirteen of the 16 *CCDC103* p.His154Pro patients (81%) have situs inversus, including in two families the presence of two siblings with situs inversus, and no families without situs inversus. Situs inversus is reported to affect approximately 50% of patients with PCD with dynein arm defects² and affected 56% of the PCD comparator group in this study ([table 2](#)). Seventy per cent of the comparator group had situs inversus in their family among affected individuals.

FEV₁ measurements showed that impact on lung function was similar in the group of 16 p.His154Pro patients compared with the PCD comparator group ([table 1](#), [figure 1](#)). The comparator group of 16 South Asian individuals with PCD all had dynein arm deficiency, but due to different genetic causes since this group comprised three *DNAAF1*,²⁴ two *DNAAF3*,²⁵ two *DNAH5*,²⁶ three *LRRC6*²⁷ and two *ZMYND10*²⁸ cases while four were excluded from *CCDC103* p.His154Pro but genetically undefined. The mean difference for clinical measures between the *CCDC103* p.His154Pro and comparator groups

Table 2 Comparison of mean values of clinical tests obtained from *CCDC103* p.His154Pro cases versus a comparator group

Clinical features	N	<i>CCDC103</i> p.His154Pro	N	Comparator group
Age (mean SD)	16	11.8 (4.8)	16	11.9 (8.8)
Gender (% male)	16	67%	16	63%
Nasal nitric oxide (nL/min)	14	63 (63)	9	12 (7)**
Ciliary beat frequency (Hz)	16	7.5 (4.7)	16	1.3 (3.6)**
% Cross sections with IDA absent on TEM	13	45% (24)	16	63% (30)
% Cross sections with ODA absent on TEM	13	19% (12)	16	89% (20)**
FEV ₁ z-score (median IQR)	10	−1.4 (−0.8, −2.1)	13	−1.8 (−1.5, −2.1)
Situs inversus	16	81%	16	56%

**p<0.005.

IDA, inner dynein arms; ODA, outer dynein arms; TEM, transmission electron microscopy.

The *CCDC103* p.His154Pro cases are described in [table 1](#) and the comparator group are South Asian origin *CCDC103* p.His154Pro-negative cases with a confirmed absent dynein arms defect and genetic results as described in the main text. We only show TEM data for patients in whom more than 100 cilia were counted. Data shown as the mean, with SD shown in brackets unless otherwise stated.

is presented in [table 2](#), showing an equivalent age and gender composition. Individuals from both groups had symptoms typical of PCD as detailed in online supplementary table 1. There was a higher rate of glue ear and positive sputum microbiology and a lower rate of bronchiectasis in the *CCDC103* p.His154Pro group compared with controls, but due to the small number of patients and heterogeneous nature of PCD it is difficult to interpret the significance of these findings.

PCD caused by the *CCDC103* p.His154Pro mutation can be associated with normal nNO results

We proceeded to retrospectively analyse clinical phenotypes of the 16 affected South Asian patients carrying the homozygous p.His154Pro missense mutation in more detail, as presented in [table 1](#). Of the 16 p.His154Pro patients tested, 7 (43%) had normal nNO levels, above 77 nL/min which is a recommended diagnostic cut-off.²⁹ Of these seven patients, most (patients 5, 7, 11, 12 and 13) were subject to repeat measurements at subsequent clinic appointments. Patient 12 displayed persistent values within the normal range up to 5 years from the first measurement. Mean nNO was significantly higher than the low levels consistently detected in the comparator PCD patient group ([table 2](#)).

PCD caused by the *CCDC103* p.His154Pro mutation can be associated with areas of normal ciliary beat frequency on high-speed video light microscopy

High-speed video microscopy revealed that among the 16 homozygous p.His154Pro individuals, 9 (56%) had ciliary beat frequencies within the local normal range of 8.5–16.8 Hz (mean 11.6 Hz) ([table 1](#)). The other seven p.His154Pro patients (patients 1–6 and 13) had a reduced beat frequency with a mean of 3.9 Hz (range 0–7.7 Hz), which is more in keeping with typical results found for patients with PCD and more specifically found for *CCDC103* patients expressing loss-of-function alleles.^{13 30} The comparator group of patients with PCD displayed a reduced ciliary beat frequency with a mean of 1.3 Hz ([table 2](#)), which is similar to the reported reduced ciliary beating seen in published PCD cases associated with other causes of dynein arm loss. In half of the p.His154Pro patients the beat pattern of cilia was completely or largely normal in some strips of epithelium (patients 5–7, 9, 10, 12, 15, 16; [table 1](#)) while static, slow or dyskinetic in others. Some p.His154Pro patients (7, 10, 16) demonstrated a full beat pattern and almost immotile cilia together within the same

sample (online supplementary videos 1–3). Therefore, the motility in p.His154Pro individuals was significantly different from controls but variable, compared with the fully motile cilia of healthy controls (online supplementary video 4) and the static cilia seen in patients from the comparator group (online supplementary video 5).

PCD caused by the *CCDC103* p.His154Pro mutation can be associated with normal ultrastructural appearance by electron microscopy

Seven of the 16 *CCDC103* p.His154Pro individuals (44%) had a defect of the ciliary inner dynein arms (IDA) and ODA demonstrated by transmission electron microscopy (TEM; cases 1–7 in [table 1](#)). The other nine p.His154Pro individuals (cases 8–16) had TEM that either showed an absence of the IDA, or that was considered normal or inconclusive despite extensive interrogation. The spectrum of ultrastructural defects found in p.His154Pro individuals is illustrated with representative examples in [figure 2](#). Overall, the IDAs appeared to be the most affected structure in these cases, with significant retention of ODAs. Quantification of these dynein arm defects is shown in [figure 3](#). Published normal range counts from >200 non-PCD respiratory controls are also shown for comparison ([figure 3](#) grey box). Notably, samples from four *CCDC103* p.His154Pro patients were completely within this normal range, a further two were closer to the normal range than the diagnostic range. Two more, although not quantifiable due to small numbers, were also reported as normal (not shown). When a partial defect of the ODA was seen in p.His154Pro patients, an assessment of proximity to the epithelial cell surface—as judged by the presence of neighbouring microvilli—showed that ODA loss typically occurred at the distal end of the cilia, towards the tips.

A summary shown in [table 2](#) highlights that the majority of *CCDC103* p.His154Pro cases clearly differed from the non-*CCDC103* comparator group, which have a near complete absence of both dynein arms. This is highly significant in terms of ODA loss (p<0.05), with only 19% ODA absence in the *CCDC103* p.His154Pro cilia compared with 89% in the comparator group which carry mutations in other dynein arm loss-associated PCD genes. IDAs were also clearly more retained in *CCDC103* p.His154Pro cilia, but this difference did not reach statistical significance. The overall relative lack of disturbance to dynein arm structures means that *CCDC103* p.His154Pro TEM overlaps with that of non-PCD controls.

Figure 2 Transmission electron microscopy of *CCDC103* p.His154Pro patients. Representative cilia cross sections show within the same micrograph (A) absent outer and partially absent inner dynein arms and (B) presence of outer dynein arms only and presence of both inner and outer dynein arms. (C) Inset shows presence of outer but not inner dynein arms. Black arrows indicate example of outer dynein arms. White arrows indicate inner dynein arms. Scale bar, 100 nm.

Biochemical analysis of the *CCDC103* p.His154Pro mutation reveals an abrogation of its oligomerisation capacity

We conducted gel filtration of purified *CCDC103* p.His154Pro protein and compared this to the normal protein using previously established methods²¹ to assess the functional viability of the mutant form of the protein. The p.His154Pro variant appears to be a highly disruptive mutation since the ability of *CCDC103* to oligomerise is significantly disrupted by the mutation, as shown in the chromatogram (figure 4). The lack of oligomers that the wild-type protein forms (~250 kDa) is however accompanied by a large increase in the void volume of the mutant protein. We speculate that this material, which is >2 MDa, is aggregated protein, rather than organised oligomers.

DISCUSSION

We report from genetic screening of a PCD cohort ascertained through UK National PCD Services that the *CCDC103* c.461A>C; p.His154Pro mutation accounts for disease in approximately one-fifth of affected individuals in the highly consanguineous South Asian PCD community, associated with loss of both the ODA and IDA of cilia. Therefore, a significant risk of disease arises from the presence of this important frequent mutation which is typically carried in homozygous state, one copy inherited from each parent. Direct screening for

this specific p.His154Pro mutation in South Asian PCD cases could be an economical diagnostic approach.

Despite the relatively high prevalence of *CCDC103* p.His154Pro, this mutation is not the sole cause of the increased incidence of PCD reported in the British South Asian community.³ Our screen also revealed patients in the group of 86 screened that carry causal mutations in a number of other genes including *CCDC40*, *DNAF1*, *DNAF3*, *HEATR2*, *LRRC6*, *ZMYND10* and *RSPH4A*. The *CCDC103* p.His154Pro defect is not exclusive to the UK Asian community and has also been detected in a patient from an Irish travelling family³¹ as well as in two North European origin families with PCD from our studies (unpublished data), all of whom have dynein arm defects.

This study highlights extensive variability in the diagnostic results for patients carrying biallelic *CCDC103* p.His154Pro mutations. Normal standard diagnostic test results have led to *CCDC103* p.His154Pro patients being discharged from the PCD clinic, and in some cases the diagnosis of PCD in a *CCDC103* p.His154Pro patient was only finally confirmed following the genetic test result. Fifteen (94%) patients had situs inversus, suggesting *CCDC103* p.His154Pro patients without situs inversus may be missed either due to lack of referral or due to normal diagnostic tests.

Figure 3 Quantitative transmission electron microscopy survey of inner and outer dynein arm loss in *CCDC103* p.His154Pro patients versus a comparator group. By surveying >100 cross sections in each patient sample we performed quantitative electron microscopy to determine the percentage of arm defects in cilia from individuals homozygous for the hypomorphic p.His154Pro *CCDC103* mutation. Note that quantitative plots have only been included within this graph if more than 100 cilia were counted and that all data have been collected at a single centre to ensure uniform results; data collected for one sample from a separate centre were therefore excluded. Red diamonds and triangles indicate results from 16 *CCDC103* p.His154Pro homozygote PCD patients, where triangles indicate the result of four repeat nasal brushings performed on patients marked by a diamond. The comparator group of 16 individuals with PCD, indicated by other symbols as shown, consists of three cases with *DNAAF1* mutations (open diamonds), two *DNAAF3* (black squares), two *DNAH5* (dark blue diamonds), three *LRRC6* (two as light blue diamonds, one contained within the filled purple circle), two *ZMYND10* (contained within the filled purple circle) and four cases in whom no mutations in *CCDC103* or other known genes could be identified (grey squares). Six *CCDC103* p.His154Pro samples (27%) of p.His154Pro samples showed complete lack of both outer and inner dynein arms comparable to other gene mutations in the graph (purple circle). The grey shaded area represents normal range counts from >200 non-PCD respiratory controls. Four *CCDC103* p.His154Pro patients had counts within this normal range (one is a repeat sample (triangle) which showed similar data). Individuals with *CCDC103* p.His154Pro mutations have a trend towards a distinctive pattern of partial loss of dynein arms that diverges from total dynein arm loss in the comparator group.

The homozygous p.His154Pro-positive patients represent an expanded phenotype for PCD, since without genetic results many in this group may not have been considered to meet the current UK clinical diagnostic criteria for PCD. This would lead to uncertainty for the patient and their parents and may result in unnecessary further investigation into the cause of their symptoms. One South Asian patient in the comparator group had a TEM phenotype similar to that of the *CCDC103* p.His154Pro group. This individual also had an nNO level of >77 nL/min, but a screen for the *CCDC103* mutation was negative (figure 3, lower left grey square). This case, along with other patients in the UK national diagnostic clinics, suggests that there will be other mutations which cause PCD with normal diagnostic results.

It is difficult to explain the variability in the p.His154Pro patients nNO (table 1). This variability is temporal within a patient and seen even between siblings. We speculate that this may be a broader reflection of ciliary function in the sinuses such that NO levels may only be low in *CCDC103* p.His154Pro patients when cilia are static (maybe during an infection or other external insult) and may normalise with improved ciliary function. The possible reasons why levels might be low in PCD were recently summarised.³²

Interestingly, preservation of some cilia structure and motility in *CCDC103* p.His154Pro patients is not apparently accompanied by significantly preserved lung function, since the FEV₁ range in p.His154Pro patients is equivalent to the comparator group and respiratory capacity is overall equally reduced in these individuals. Another missense mutation causing PCD, *ZMYND10*

Figure 4 *CCDC103* p.His154Pro oligomerisation capacity chromatograms of wild-type (solid trace) and His154Pro (dashed trace) *CCDC103* native proteins separated in a calibrated Superose 6 10/300 gel filtration column. The data are plotted as absorbance at 280 nm (in mAU) against elution volume (mL). Both proteins show strong dimer peaks at ~60 kDa. However, only the wild-type form generates a series of higher order oligomers with an approximate mass of ~250 kDa. Aggregated material (>2 MDa) eluted in the void volume.

p.Val16Gly, was similarly found to cause a mixed cilia beat defect with a significant degree of retained cilia motility but this is also not yet associated with milder disease course.²⁸ In future such studies, larger numbers of *CCDC103* p.His154Pro patients and controls should be analysed to confirm this observation and genotype—phenotype relationships could be further investigated with more sensitive tests for staging lung disease such as Lung Clearance Index, CT scan or radiolabelling methods screening mucociliary clearance rates.^{33 34}

Patients with the *CCDC103* p.His154Pro mutation represent an interesting cohort for targeted pharmacogenetic therapies due to their prevalence and since the dynein components required for normal ciliary beating appear to still be present in the patient's cilia, although at variable levels.

The finding of nNO levels within the normal range in *CCDC103* p.His154Pro patients highlights the importance of considering the full clinical history in conjunction with nNO testing in patients with suspected PCD. Clinicians should proceed to further testing in cases with a high index of clinical suspicion, using the NO test as part of a multidisciplinary diagnostic protocol rather than a stand-alone screening test. This is not the first report of normal nNO results in patients with PCD. Some patients with *RSPH1* mutations are also reported to have levels of NO close to normal and we expect further such cases to

be reported as the complex genetic landscape of PCD diagnosis is unravelled further.³⁵

The apparently normal/mixed cilia beat pattern found in half of p.His154Pro cases was notably high, but this finding is supported by previous work indicating that this represents a hypomorphic allele in cilia function tests.¹³ Awareness of these cases is critical when assessing diagnostic samples in the laboratory, if the observer inadvertently tends to select beating strips for analysis over those that are static when scanning the sample at 5–20 times magnification before full analysis with high-speed video, thereby inadvertently missing this defect when a significant portion of the sample has a normal coordinated beat. This places an emphasis on diagnostic centres to increase awareness and expertise of the operators when assessing nasal biopsies by high-speed video microscopy. HSVM does not quantify the power of the ciliary beating, but in cases with a normal stroke and frequency there could be weakness in the strength of the beating that might not be detected by any of the current diagnostic tests.

In this study, TEM diagnosis often revealed a pattern of intermittent IDA and ODA loss in *CCDC103* p.His154Pro patients which was distinct from cases of dynein arm absence due to mutations in other dynein loss-associated PCD genes, for example, *LRRC6*, *DNAAF1* and *ZYMN10*. The distinction was

most significant for ODA retention with *CCDC103* p.His154Pro cases, with comparatively high levels of preserved ODAs seen in patient's cilia while the IDAs were more often missing. Despite analysis of >100 cilia cross sections, usually sufficient for diagnosis, the loss of dynein was not always detected by electron microscopy. However, it has long been accepted that normal ultrastructure by electron microscopy cannot exclude PCD.^{36 37} Our study appears to confirm in *CCDC103* p.His154Pro patients the previous evidence that *CCDC103* mutations confer a loss of distal ODAs containing DNAH9, not ODAs in the proximal half of the cilia closer to the epithelial surface.¹³

This study contributes strong evidence that *CCDC103* p.His154Pro is pathogenic rather than a benign polymorphism. It is present at very low frequency in the non-PCD population and no *CCDC103* p.His154Pro homozygote individuals were detected in large-scale screening of 9798 South Asian and 33,345 North European controls. Our segregation analysis performed where parental samples were available in the affected families studied here showed an inheritance pattern fully consistent with recessive disease (table 1). The *CCDC103* protein remains poorly characterised but previous studies show that it usually forms dimers and higher order oligomers. It is thought to help in generating a high-affinity site on the doublets for outer arm assembly, either through direct interactions or indirectly, by modifying the underlying microtubule lattice.²¹ The oligomerisation capacity of *CCDC103* is a property of the central region of the protein (RSPK and SMK, unpublished observations) which contains a highly conserved RPAP3_C domain that spans residues 96–189 of the protein and is predicted to function in protein–protein interactions.²¹ The His154 amino acid is located within an alpha helix of this RPAP3_C functional domain and substitution of the cyclic side chain of proline at this position would enforce a conformation predicted incompatible with alpha helical secondary structure. Thus, altering His154 is expected to disrupt the protein's secondary structure and consequently its function. Our current results reveal that the His154Pro mutant form retains the ability to dimerise but shows little oligomer formation, suggesting that this property is disrupted by the mutation and that *CCDC103* may have two distinct self-interaction domains. We hypothesise that gene mutations causing instability or depletion of *CCDC103* protein may make the attachment of the dynein arms more susceptible to physical, infective or inflammatory insult. We cannot determine if the presence of normal cilia in our patients is temporal or spatial; however, as results appear to vary from one biopsy to the next, for example, in TEM data for cases 11, 12, 15 and 16 in table 1, therefore cell culture and repeating investigations may be useful in these cases.

It can often be difficult to make or exclude a diagnosis of PCD due to poor sensitivity of electron microscopy and genetic testing and the poor specificity of nNO measurement.⁵ Currently, diagnosis using multiple tests is recommended. These tests are often complex and require specialist equipment and skills to interpret. Consequently, several patients continue to have an indeterminate diagnosis. Awareness of variants such as *CCDC103* p.His154Pro may allow targeted gene screening in patients with an indeterminate diagnosis.

In conclusion, the *CCDC103* p.His154Pro variant is prevalent in the UK South Asian community and likely to be found in South Asian patients worldwide. This patient group should therefore undergo genetic testing for c.461A>C, especially if (partial) dynein arm absence is suspected. These patients frequently present a diagnostic dilemma, due to inconclusive results of multiple clinical diagnostic tests. This study expands

the diagnostic phenotype which we consider to be PCD, since in some cases pathology-based tests can be equivocal as the cilia can beat at least partially in a coordinated manner, at the correct speed and may appear structurally normal. PCD is widely understood to be an underdiagnosed condition and this appears to be the case for *CCDC103* p.His154Pro patients, who demonstrate a high level of situs inversus probably indicative of a lack of recognition making their diagnosis liable to be missed.³⁸ We anticipate that studies such as this, in combination with easier access to high throughput and economically achievable genetic screening, should greatly increase disease recognition and understanding. We have highlighted the importance of multidisciplinary testing, repeat testing and genotyping in patients with a highly suggestive history for PCD.

Author affiliations

¹Department of Paediatric Respiratory Medicine, Royal Brompton and Harefield NHS Trust, National Heart and Lung Institute, London, UK

²Division of Services for Women and Children, Women's and Newborn Unit Bradford Royal Infirmary, University of Bradford, Bradford, UK

³Department of Infection, Centre for PCD Diagnosis and Research, Immunity and Inflammation, University of Leicester, Leicester, UK

⁴Genetics and Genomic Medicine, University College London, UCL Great Ormond Street Institute of Child Health, London, UK

⁵North East Thames Regional Genetics Service, Great Ormond Street Hospital for Children, London, UK

⁶Neuroscience and Mental Health Research Institute, School of Medicine and School of Bioscience, Cardiff University, London, UK

⁷Human Genetics Department, Medical Research Institute, Alexandria University, Alexandria, Egypt

⁸Genome Research Division, Human Genetics Department, Radboud University Medical Center and Radboud Institute for Molecular Life Sciences, Nijmegen, The Netherlands

⁹Pediatric Genetics Division, Center for Pediatrics and Adolescent Medicine, University of Freiburg Medical Center, Freiburg, Germany

¹⁰Department of Molecular Biology and Biophysics, University of Connecticut Health Center, Farmington, Connecticut, USA

¹¹Department of Respiratory Medicine, Royal Brompton and Harefield NHS Trust, London, UK

¹²Primary Ciliary Dyskinesia Centre, University Hospital Southampton NHS Foundation Trust and Clinical and Experimental Sciences Academic Unit, University of Southampton Faculty of Medicine, Southampton, UK

¹³NIHR Southampton Respiratory Biomedical Research Unit, University of Southampton and University Hospital Southampton NHS Foundation Trust, Southampton, UK

¹⁴Department of Paediatrics, Luton and Dunstable Hospital NHS Trust, Luton, UK

¹⁵Department of Paediatric Respiratory Medicine, National Heart and Lung Institute, Imperial College, London, UK

¹⁶Centre for Translational Omics-GOSgene, Genetics and Genomic Medicine, University College London, Institute of Child Health, London, UK

¹⁷Yorkshire Regional Genetics Service and School of Medicine, University of Leeds, St. James's University Hospital, Leeds, UK

¹⁸Population, Policy and Practice Programme, University College London, UCL Great Ormond Street Institute of Child Health, London, UK

¹⁹Department of Respiratory Paediatrics, Birmingham Children's Hospital NHS Foundation Trust, Birmingham, UK

²⁰Infection, Immunity, Inflammation and Physiological Medicine, University College London, Institute of Child Health, London, UK

²¹Institute for Systems Genomics, University of Connecticut Health Center, Farmington, Connecticut, USA

Acknowledgements We are very grateful to the families with PCD who have participated in this study and to the UK PCD Family Support Group for their support. We thank Louise Ocaka, Chela James (UCL GOSgene) and Emily Frost (Royal Brompton Hospital) for experimental support and data analysis.

Contributors AS, HMM, EAR and EM were responsible for study concept and design. MPP, JH, JS, MRF, WL, HW, CB, LJ, CMW, TC and HMM performed the genetic analyses. RSPK and SMK performed the biochemical analysis. AS, MD, AVR, AR and PG performed electron microscopy. AS, RAH, MD, AVR, CLJ and SO performed cilia cell culture and cilia functional testing. EM, EAR, SC, WW, BA, MRL, BR, RW, AB, ES, JSL, PK, COC and CH provided clinical information. AS, SMK, CH and HMM wrote the manuscript and all authors contributed to the final version. HMM is the guarantor responsible for the overall content.

Funding The research is supported by the BEAT-PCD: Better Evidence to Advance Therapeutic options for PCD network (COST Action 1407). AB was supported by the NIHR Respiratory Disease Biomedical Research Unit at the Royal Brompton and Harefield NHS Foundation Trust and Imperial College London. Work by AS is independent research funded by a postdoctoral research fellowship from the National Institute of Health Research and Health Education England. RSPK and SMK are supported by NIH grant GM051293. MS is supported by a Radboudumc Hypatia Tenure Track fellowship, a Radboud University Excellence fellowship, an ERC starting grant (TREATCilia, grant no.716344) and received funding from the German Research Foundation (DFG), collaborative Research Center (CRC) 1140 KIDGEM. This research and the Centre for Translational Omics (GOSgene) is supported by the National Institute for Health Research Biomedical Research Centre at Great Ormond Street Hospital for Children NHS Foundation Trust and University College London. HMM was supported by grants from Action Medical Research (GN2101), Newlife Foundation (10-11/15) and the Great Ormond Street Hospital Children's Charity. Work in Southampton is supported by the NIHR Respiratory Biomedical Research Unit and NIHR Wellcome Trust Clinical Research Facility.

Competing interests None declared.

Ethics approval London Bloomsbury Research Ethics Committee.

Provenance and peer review Not commissioned; externally peer reviewed.

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2018. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- Lucas JS, Burgess A, Mitchison HM, *et al.* Diagnosis and management of primary ciliary dyskinesia. *Arch Dis Child* 2014;99:850–6.
- Shapiro AJ, Davis SD, Ferkol T, *et al.* Laterality defects other than situs inversus totalis in primary ciliary dyskinesia: insights into situs ambiguus and heterotaxy. *Chest* 2014;146:1176–86.
- O'Callaghan C, Chetcuti P, Moya E. High prevalence of primary ciliary dyskinesia in a British Asian population. *Arch Dis Child* 2010;95:51–2.
- Shah A, Shoemark A, MacNeill SJ, *et al.* A longitudinal study characterising a large adult primary ciliary dyskinesia population. *Eur Respir J* 2016;48:441–50.
- Lucas JS, Barbato A, Collins SA, *et al.* European Respiratory Society guidelines for the diagnosis of primary ciliary dyskinesia. *Eur Respir J* 2016 (49:1601090).
- Jackson CL, Behan L, Collins SA, *et al.* Accuracy of diagnostic testing in primary ciliary dyskinesia. *Eur Respir J* 2016;47:837–48.
- El Khouri E, Thomas L, Jeanson L, *et al.* Mutations in DNAJB13, encoding an HSP40 Family Member, Cause Primary Ciliary Dyskinesia and Male Infertility. *Am J Hum Genet* 2016;99:489–500.
- Jeanson L, Copin B, Papon JF, *et al.* RSPH3 mutations cause Primary Ciliary Dyskinesia with Central-Complex Defects and a near absence of Radial spokes. *Am J Hum Genet* 2015;97:153–62.
- Wallmeier J, Al-Mutairi DA, Chen CT, *et al.* Mutations in CCNO result in congenital mucociliary clearance disorder with reduced generation of multiple motile cilia. *Nat Genet* 2014;46:646–51.
- Wallmeier J, Shiratori H, Dougherty GW, *et al.* TTC25 Deficiency results in defects of the Outer Dynein arm docking Machinery and primary ciliary dyskinesia with Left-Right body asymmetry randomization. *Am J Hum Genet* 2016;99:460–9.
- Olbrich H, Cremers C, Loges NT, *et al.* Loss-of-Function GAS8 mutations cause Primary Ciliary Dyskinesia and disrupt the Nexin-Dynein Regulatory complex. *Am J Hum Genet* 2015;97:546–54.
- Watson CM, Crinnion LA, Morgan JE, *et al.* Robust diagnostic genetic testing using solution capture enrichment and a novel variant-filtering interface. *Hum Mutat* 2014;35:434–41.
- Panizzi JR, Becker-Heck A, Castleman VH, *et al.* CCDC103 mutations cause primary ciliary dyskinesia by disrupting assembly of ciliary dynein arms. *Nat Genet* 2012;44:714–9.
- Lucas JS, Chetcuti P, Copeland F, *et al.* Overcoming challenges in the management of primary ciliary dyskinesia: the UK model. *Paediatr Respir Rev* 2014;15:142–5.
- Le Quesne Stabej P, Williams HJ, James C, *et al.* STAG3 truncating variant as the cause of primary ovarian insufficiency. *Eur J Hum Genet* 2016;24:135–8.
- Trump N, McTague A, Brittain H, *et al.* Improving diagnosis and broadening the phenotypes in early-onset seizure and severe developmental delay disorders through gene panel analysis. *J Med Genet* 2016;53:310–7.
- Beydon N, Davis SD, Lombardi E, *et al.* An official American Thoracic Society/European Respiratory Society statement: pulmonary function testing in preschool children. *Am J Respir Crit Care Med* 2007;175:1304–45.
- Quanjer PH, Stanojevic S, Cole TJ, *et al.* Multi-ethnic reference values for spirometry for the 3–95-yr age range: the global lung function 2012 equations. *Eur Respir J* 2012;40:1324–43.
- Chilvers MA, Rutman A, O'Callaghan C. Ciliary beat pattern is associated with specific ultrastructural defects in primary ciliary dyskinesia. *J Allergy Clin Immunol* 2003;112:518–24.
- Shoemark A, Dixon M, Corrin B, *et al.* Twenty-year review of quantitative transmission electron microscopy for the diagnosis of primary ciliary dyskinesia. *J Clin Pathol* 2012;65:267–71.
- King SM, Patel-King RS. The oligomeric outer dynein arm assembly factor CCDC103 is tightly integrated within the ciliary axoneme and exhibits periodic binding to microtubules. *J Biol Chem* 2015;290:7388–401.
- Sheridan E, Wright J, Small N, *et al.* Risk factors for congenital anomaly in a multiethnic birth cohort: an analysis of the born in Bradford study. *Lancet* 2013;382:1350–9.
- Lek M, Karczewski KJ, Minikel EV, *et al.* Analysis of protein-coding genetic variation in 60,706 humans. *Nature* 2016;536:285–91.
- Loges NT, Olbrich H, Becker-Heck A, *et al.* Deletions and point mutations of LRRC50 cause primary ciliary dyskinesia due to dynein arm defects. *Am J Hum Genet* 2009;85:883–9.
- Mitchison HM, Schmidts M, Loges NT, *et al.* Mutations in axonemal dynein assembly factor DNAAF3 cause primary ciliary dyskinesia. *Nat Genet* 2012;44:381–9.
- Olbrich H, Häffner K, Kispert A, *et al.* Mutations in DNAH5 cause primary ciliary dyskinesia and randomization of left-right asymmetry. *Nat Genet* 2002;30:143–4.
- Kott E, Duquesnoy P, Copin B, *et al.* Loss-of-function mutations in LRRC6, a gene essential for proper axonemal assembly of inner and outer dynein arms, cause primary ciliary dyskinesia. *Am J Hum Genet* 2012;91:958–64.
- Moore DJ, Onoufriadis A, Shoemark A, *et al.* Mutations in ZMYND10, a gene essential for proper axonemal assembly of inner and outer dynein arms in humans and flies, cause primary ciliary dyskinesia. *Am J Hum Genet* 2013;93:346–56.
- Leigh MW, Hazucha MJ, Chawla KK, *et al.* Standardizing nasal nitric oxide measurement as a test for primary ciliary dyskinesia. *Ann Am Thorac Soc* 2013;10:574–81.
- Raidt J, Wallmeier J, Hjej R, *et al.* Ciliary beat pattern and frequency in genetic variants of primary ciliary dyskinesia. *Eur Respir J* 2014;44:1579–88.
- Casey JP, Goggin P, McDaid J, *et al.* A case report of primary ciliary dyskinesia, laterality defects and developmental delay caused by the co-existence of a single gene and chromosome disorder. *BMC Med Genet* 2015;16:45.
- Walker WT, Jackson CL, Lackie PM, *et al.* Nitric oxide in primary ciliary dyskinesia. *Eur Respir J* 2012;40:1024–32.
- Walker WT, Young A, Bennett M, *et al.* Pulmonary radioaerosol mucociliary clearance in primary ciliary dyskinesia. *Eur Respir J* 2014;44:533–5.
- Nielsen JG. Lung clearance index: should we really go back to nitrogen washout? *Eur Respir J* 2014;43:655–6.
- Knowles MR, Ostrowski LE, Leigh MW, *et al.* Mutations in RSPH1 cause primary ciliary dyskinesia with a unique clinical and ciliary phenotype. *Am J Respir Crit Care Med* 2014;189:707–17.
- Olbrich H, Schmidts M, Werner C, *et al.* Recessive HYDIN mutations cause primary ciliary dyskinesia without randomization of left-right body asymmetry. *Am J Hum Genet* 2012;91:672–84.
- Schwabe GC, Hoffmann K, Loges NT, *et al.* Primary ciliary dyskinesia associated with normal axoneme ultrastructure is caused by DNAH11 mutations. *Hum Mutat* 2008;29:289–98.
- Kuehni CE, Frischer T, Strippoli MP, *et al.* Factors influencing age at diagnosis of primary ciliary dyskinesia in European children. *Eur Respir J* 2010;36:1248–58.