

SUPPLEMENTARY FILE

eTable 1. Sensitivity analysis

<i>Pharmacological intervention</i>	<i>Network meta-analysis estimates</i>	<i>A) Haemodynamically stable</i>	<i>B) > 65 years</i>	<i>C) Alternative statistical model</i>
ALL-CAUSE MORTALITY				
Compared with anticoagulation				
Full-dose thrombolysis	0.60 (0.36-1.01)	0.59 (0.29-1.22)	0.75 (0.31-1.81)	0.61 (0.36-1.04)
Low-dose thrombolysis	0.47 (0.14-1.59)	1.13 (0.04-31.83)	0.36 (0.01-20.70)	0.58 (0.04-8.70)
Catheter-directed thrombolysis	0.31 (0.01-7.96)	0.31 (0.01-7.96)	-	0.55 (0.13-2.31)
Compared with full-dose thrombolysis				
Low-dose thrombolysis	1.28 (0.40-4.12)	1.90 (0.07-49.60)	0.48 (0.01-25.18)	0.89 (0.23-3.41)
Catheter-directed thrombolysis	1.93 (0.07-51.33)	0.53 (0.02-14.54)	-	0.51 (0.02-13.56)
Compared with low-dose thrombolysis				
Catheter-directed thrombolysis	1.50 (0.05-47.94)	0.28 (0-29.11)	-	0.67 (0.02-21.26)
MAJOR BLEEDING				
Compared with anticoagulation				
Full-dose thrombolysis	2.00 (1.06-3.78)	2.29 (0.78-6.71)	3.39 (1.01-11.42)	1.99 (1.04-3.79)
Low-dose thrombolysis	0.90 (0.25-3.21)	0.43 (0.01-18.62)	4.93 (0.30-80.95)	1.11 (0.02-78.75)
Catheter-directed thrombolysis	0.97 (0.02-56.03)	0.97 (0.01-66.76)	-	0.89 (0.23-3.41)
Compared with full-dose thrombolysis				
Low-dose thrombolysis	2.22 (0.71-6.89)	0.19 (0.01-6.94)	1.45 (0.12-18.09)	0.45 (0.14-1.46)
Catheter-directed thrombolysis	2.07 (0.03-126.08)	0.42 (0.01-33.32)	-	0.49 (0.01-29.81)
Compared with low-dose thrombolysis				
Catheter-directed thrombolysis	0.93 (0.01-65.65)	2.23 (0.01-640.48)	-	1.09 (0.02-78.62)