

# State of the art review: from the seaside to the bedside: insights from comparative diving physiology into respiratory, sleep and critical care

Paul J Ponganis<sup>1</sup>

Scripps Institution of Oceanography, University of California San Diego, La Jolla, CA, USA

## Correspondence to

Dr. Paul J Ponganis, Scripps Institution of Oceanography, University of California San Diego, La Jolla, CA 92093-0204, USA; pponganis@ucsd.edu

Received 22 October 2018  
Revised 20 January 2019  
Accepted 28 January 2019  
Published Online First  
2 March 2019


## ABSTRACT

Anatomical and physiological adaptations of animals to extreme environments provide insight into basic physiological principles and potential therapies for human disease. In that regard, the diving physiology of marine mammals and seabirds is especially relevant to pulmonary and cardiovascular function, and to the pathology and potential treatment of patients with hypoxaemia and/or ischaemia. This review highlights past and recent progress in the field of comparative diving physiology with emphasis on its potential relevance to human medicine.


## INTRODUCTION

Anatomical and physiological adaptations underlie the dive capacities, foraging behaviours and ecology of marine mammals and seabirds. The dive response (the bradycardia and vasoconstriction associated with a breath hold), enhanced oxygen stores, hypoxaemic tolerance and pressure tolerance of these animals allow for remarkable physiological and behavioural feats. Severe bradycardias (figure 1), arterial  $pO_2$  near 10 mm Hg (1.3 kPa) (figure 2) and maximum dive depths of 500–1500 m occur in multiple species at sea.<sup>1</sup> The deepest and longest dives have been reported for Cuvier's beaked whale at 2992 m and 163 min, respectively.<sup>2,3</sup>

These physiological adaptations for diving are also relevant to basic physiological principles and to potential therapies in cardiovascular and respiratory medicine. For instance, the activation of trigeminal



**Figure 1** Heart rate profiles in dives to greater than 350 m depth of emperor penguins (A) and California sea lions (B) demonstrate the bradycardia of diving.<sup>105 106</sup> In such deep dives, heart rates routinely approach 10 beats per minute, and at times exhibit considerable variability (adapted from McDonald and Ponganis<sup>105</sup> and Wright *et al*<sup>106</sup>).



**Figure 2** During these routine 15–20 min dives of northern elephant seals,<sup>60</sup> arterial  $pO_2$  profiles demonstrated an initial compression hyperoxia, and then a gradual decline to about 20 mm Hg (2.67 kPa). Data were collected with an intravascular electrode and backpack recorder. Shaded area indicated dive (adapted from Meir *et al*<sup>60</sup>).

nerve reflexes and the dive response by cold water application to the face is a classic vagal manoeuvre to treat supraventricular tachycardia.<sup>4</sup> In addition, simultaneous activation of both sympathetic and parasympathetic reflexes during cold water immersion of humans has been considered to contribute to autonomic conflict and cardiac arrhythmias.<sup>5</sup>

In this paper, I will review the relevance of comparative physiology and some of the adaptations for diving to the teaching of basic physiological principles and to potential therapies in medicine. These topics include (A) airway anatomy and respiratory flow rates, (B) alveolar collapse at depth, and surfactant function, (C) the roles of the dive response, vena caval sphincter and aortic bulb of seals in regulation of myocardial oxygen supply/demand, (D) haemoglobin (Hb) oxygen affinity in relation to oxygen transport during hypoxaemia, (E) cerebral hypoxaemic tolerance, (F) antioxidants and avoidance of reperfusion injury, and (G) the potential roles of the gasotransmitters—nitric oxide, hydrogen sulfide and carbon monoxide.


## LUNG MECHANICS AND AIR FLOW

The importance of tracheobronchial anatomy and compliance in respiratory mechanics is exemplified in the flow volume curves, maximum expiratory flow rates and tidal volumes of cetaceans (whales and dolphins).<sup>6</sup> In contrast to human subjects, and especially, patients with emphysema (figure 3), flow volume curves of the bottlenose dolphin and of excised harbour porpoise lungs are remarkable for



© Author(s) (or their employer(s)) 2019. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Ponganis PJ. *Thorax* 2019;74:512–518.



**Figure 3** Flow volume curves (expiratory flow rate vs lung volume) demonstrate the magnitude of high flow rates as well as the maintenance of high flow at lower lung volumes in the harbour porpoise.<sup>8</sup> In comparison, flow rates (expressed in % of vital capacity (VC)) of a healthy human and a patient with emphysema are considerably lower and not maintained at lower lung volumes.<sup>107</sup> Maintenance of such high flow rates in the porpoise is attributed to cartilaginous reinforcement throughout the distal airways (adapted from Kooyman and Sinnott<sup>8</sup> and Hyatt *et al*<sup>107</sup>).


the maintenance of high flows at low lung volumes.<sup>7,8</sup> Furthermore, maximum expiratory flow rates in cetaceans and sea lions (expressed in terms of vital capacity (VC)) are in the range of 5–8 VC/s<sup>7–9</sup>, much higher than maximum human expiratory flow rates of about 2 VC/s<sup>10</sup>. Such flows allow for a tidal volume as high as 88% of total lung capacity (TLC) in the pilot whale (*Globicephala melena*),<sup>11</sup> and 95% of TLC in bottlenose dolphins (*Tursiops truncatus*).<sup>12</sup> In comparison, the average human tidal volume is less than 10% of TLC.<sup>10</sup>

The reinforcement of the distal tracheobronchial tree in marine mammals is considered to promote such rapid air exchange in marine mammals. The anatomical basis of such reinforcement varies in different species, and includes the presence of (A) cartilaginous structures in the distal airway walls of cetaceans and sea lions, (B) prominent muscle fibres in the bronchial walls of seals, and (C) large vascular plexuses that have been postulated to become engorged at depth to reinforce the airways in deep-diving whales.<sup>13–16</sup>

Lastly, a recent report of high collateral ventilation flow rates in the unicameral dolphin lung should be of interest in relation to the role of collateral ventilation in emphysema and to its effects on outcomes from endoscopic lung volume reduction procedures.<sup>17</sup> In a novel hypothesis, it has been postulated that high collateral flow and constriction of the dolphin's peribronchiolar sphincter muscles during a dive contribute to optimisation of ventilation-perfusion matching.<sup>18</sup>

### ALVEOLAR COLLAPSE AND SURFACTANT

Scholander's observations of cartilaginous distal airways in whales led him to hypothesise that more rigid airways would also allow (1) movement of air into those airways during compression of the lungs at depth, (2) collapse of alveoli, (3) cessation of gas exchange at depth, and (4) the avoidance of excess nitrogen absorption during dives.<sup>16</sup> This concept has been supported by (A) low relaxation volumes of excised lungs of various marine mammals relative to those of terrestrial counterparts,<sup>8,19,20</sup> (B) pulmonary shunts in pressure chamber studies,<sup>21</sup> (C) blood



**Figure 4** The arterial partial pressure of nitrogen ( $P_a N_2$ ) remained low during a free dive of a Weddell seal (WS) to about 90 m depth, and during a simulated dive of an elephant seal (ES) in a pressure chamber to 136 m depth.<sup>22,23</sup> Alveolar collapse and lack of gas exchange was estimated to occur at 20–50 m depth from these data. Depth and  $P_a N_2$  are both expressed in atmospheres absolute (ATA). In the absence of a pulmonary shunt or alveolar collapse, arterial  $P_a N_2$  should approach alveolar nitrogen levels, approximately 80% the ambient pressure at depth (adapted from Falke *et al*<sup>22</sup> and Kooyman *et al*<sup>23</sup>).

nitrogen levels that were less than expected in the absence of shunting or alveolar collapse at depth<sup>22–24</sup> (figure 4), (D) characteristic arterial  $pO_2$  profiles at threshold depths in sea lions,<sup>25</sup> and (E) modelling studies based on lung volumes and respiratory compliance measurements.<sup>26,27</sup> Although the depth at which gas exchange ceases may vary in marine mammals (above references<sup>22–27</sup>), routine collapse and re-expansion of marine mammal lungs during dives has led to the question as to whether there are adaptations in the pulmonary surfactant of these animals. Elephant seals, for example, spend at least 80% of a 3-month trip to sea underwater at average depths of 400 m with typical dive durations of 20 min and surface intervals of only 2 min.<sup>28</sup> Although alveolar collapse was originally considered to occur at 20–50 m depth in these seals,<sup>22,23</sup> modelling studies now suggest a deeper threshold, near 160 m depth.<sup>29</sup> Regardless of the exact depth of alveolar collapse, most gas exchange is limited to the brief surface intervals during which lung expansion and function are critical.

Analysis of bronchoalveolar lavage fluid from young seals, sea lions and other marine mammals revealed a greater concentration of phospholipid and more fluidic species of phosphatidylcholine than in terrestrial mammals.<sup>30,31</sup> This was considered consistent with more rapid spreading of phospholipids during alveolar re-expansion. Surfactant from elephant seals produced moderately elevated minimum surface tension measurements. Poor surface activity and minimal reduction of surface tension were also later observed in analyses of lavage fluid from excised lungs of other pinniped species.<sup>32</sup> In addition, further analyses of surfactant composition of pinnipeds revealed a relative decrease in anionic phospholipids, an increase in short-chain phospholipids and a decrease in surfactant protein B.<sup>33,34</sup> Based on all these findings, it has been suggested that the primary function of surfactant in deep-diving mammals may be an antiadhesive function rather than only surface tension reduction.<sup>35</sup> This antiadhesive function has been considered to include the reduction of adhesive interactions between respiratory surfaces due to variation in surface tension.<sup>36</sup> In this situation, as two surfaces,

apposed against each other in a collapsed alveolus, begin to pull apart during lung expansion, surfactant lipids are considered to rise to the surface at the air–liquid interface, thereby lowering surface tension and decreasing the work required to expand the lung.

### MYOCARDIAL OXYGEN SUPPLY AND DEMAND


The dive response and cardiovascular adaptations of seals provide an excellent demonstration of the regulation and optimisation of myocardial oxygen supply and demand.<sup>1,37</sup> Heart rate reduction during the breath hold is variable, dependent on the nature of a dive and is most severe during forced submersions.<sup>38,39</sup> The bradycardia of diving optimises myocardial oxygen supply and demand by increasing diastolic perfusion time for the heart while at the same time decreasing myocardial workload. In addition, despite increased sympathetic tone and intense vasoconstriction during the severe bradycardia of forced submersion, left ventricular maximum rate of pressure change ( $dp/dt_{max}$ ) and left ventricular wall segment shortening were decreased in seals.<sup>40,41</sup> Thus, myocardial contractility and its associated oxygen demand are also decreased during the dive response.

Optimisation of cardiac preload is necessary to prevent increased wall stress and overdistention of the heart. In seals, a caval sphincter, composed of striated muscle, and innervated by the right phrenic nerve, surrounds the thoracic portion of posterior vena cava at the level of the diaphragm.<sup>42</sup> The vena caval sphincter, which has been observed to open and contract with each cardiac cycle during forced submersions, represents a unique mechanism to regulate venous return and optimise preload of the heart.<sup>43,44</sup> During the severe bradycardia of forced submersion, decreased left ventricular wall segment dimensions, decreased left ventricular  $dp/dt_{max}$ , unchanged left ventricular diastolic pressures, unchanged central venous pressures and decreased stroke volumes were all consistent with decreased ventricular filling, decreased wall tension and decreased contractility.<sup>40,41,45</sup> All these factors would minimise myocardial oxygen demand. The lack of any increase in left ventricular diastolic pressure will also aid in maintenance of myocardial perfusion because the driving pressure for coronary perfusion during diastole is the difference between the arterial pressure and the intra-ventricular pressure.

In seals, the effects of widespread vasoconstriction and increased systemic vascular resistance on the afterload and oxygen consumption of the left ventricle are minimised by the aortic bulb, an elastic dilatation of the aortic root.<sup>46,47</sup> The distensible walls of the aortic bulb decrease the afterload the ventricle would otherwise face. Thus, the workload of the heart during the dive response is decreased by yet another mechanism. In addition, the elastic wall of the expanded bulb gradually contracts during diastole, maintaining both blood flow and diastolic blood pressure which is necessary for coronary perfusion<sup>48</sup> (figure 5). It is remarkable that the physiological effects of the aortic bulb in seals are exactly analogous to the therapeutic effects (afterload reduction and maintenance of diastolic perfusion pressure) of the intra-aortic balloon pump.<sup>49</sup> Aortic bulbs also occur in other types of marine mammals, and an elastic, highly compliant aortic arch in the large baleen whales is hypothesised to accomplish the same optimisation of myocardial oxygen supply/demand and maintenance of diastolic blood flow.<sup>50,51</sup>

### HB'S OXYGEN AFFINITY

Recently, GBT1118, a novel compound that reversibly binds to Hb and increases oxygen affinity, has been demonstrated




**Figure 5** A comparison of blood flow in the descending aorta during a single cardiac cycle in humans and seals illustrated the maintenance of diastolic flow in the seal.<sup>48,108</sup> This higher diastolic flow was attributed to the slow contraction of the distended aortic bulb in the seal. Data were obtained with MRI techniques (adapted from Thornton *et al*<sup>48</sup> and Hochachka<sup>108</sup>).

in murine models to decrease Hb polymerisation in sickle cell disease, attenuate hypoxaemia and pulmonary fibrosis in bleomycin toxicity and improve survival in acute lung injury.<sup>52–54</sup> The rationale for modification of Hb's oxygen affinity to increase blood oxygen content in the treatment of hypoxaemia in human patients is supported by the left shift (increased affinity) of the oxygen-Hb dissociation curve in penguins and high-altitude birds.<sup>55–58</sup> The  $P_{50}$  values ( $pO_2$  at 50% Hb saturation) of deep-diving emperor penguins and high-flying bar-headed geese are 10–15 mm Hg (1.3–2.0 kPa) less than those of low-altitude birds (figure 6). However, such leftward shifts in the dissociation curve are not apparent in marine mammals; the  $P_{50}$  values of two of the most accomplished pinniped divers, the elephant seal and Weddell seal, were 27–30 mm Hg (3.6–4.0).<sup>59,60</sup> Similarly, although the Hbs of some high-altitude mammals have increased oxygen affinity, the evidence for such an adaptation among all high-altitude mammals was equivocal.<sup>61</sup> Further investigation is necessary.

### CEREBRAL HYPOXAEMIC TOLERANCE


In forced submersion studies of cerebral hypoxaemic tolerance, seals tolerated arterial  $pO_2$  as low as 10 mm Hg (1.3 kPa).<sup>62,63</sup> During the routine 20 min dives of elephant seals (figure 7), arterial Hb saturations were consistently between 80% and 20% for most of the dive duration.<sup>60</sup> The lowest saturation recorded was 8%. Significant desaturation also occurred during the frequent spontaneous sleep apnoeas that occur on land (figure 8).<sup>64</sup> The mechanisms underlying such tolerance appear to be multiple, and may provide insights to the design of therapeutic options for cerebral hypoxaemia and ischaemia. In the seal, high Hb concentrations (25 g/dL), hypercarbia (with both its associated rightward shift (decreased affinity) in the oxygen-Hb dissociation curve and its increase in cerebral blood flow) and increased brain capillary density all enhance oxygen delivery under these circumstances.<sup>63</sup> In addition, selective brain cooling can occur via arteriovenous shunting in the foreflippers with brain temperature declines of 3°C–4°C during 15 min forced submersions.<sup>65,66</sup> Such findings lend support to current advanced cardiac life support guidelines in regard to avoidance of hyperventilation during resuscitation



**Figure 6**  $O_2$ -haemoglobin dissociation curves of the emperor penguin and bar-headed goose demonstrated a left shift relative to that of the domestic duck.<sup>55,56</sup> It was postulated that the increased oxygen affinity of the penguin and high-altitude goose was beneficial in providing higher blood oxygen content at the lower  $pO_2$  experienced by these birds (adapted from Black and Tenney<sup>55</sup> and Meir and Ponganis<sup>56</sup>).

from cardiac arrest and to the potential role of therapeutic hypothermia in neurological recovery from ischaemic insult.

Remarkably, synaptic transmission is maintained in seal brain tissue even under conditions of extreme (<1%  $O_2$ ) hypoxia.<sup>67</sup> In comparison to terrestrial controls, biochemical adaptations and molecular mechanisms that support the seal brain under such conditions include: (A) a threefold elevation in brain glycogen, (B) increased tolerance to both elevated lactate and decreased glucose concentrations, but no increase in pH buffering capacity, (C) normal lactate dehydrogenase (LDH) activity with >70%




**Figure 8** Blood and muscle oxygen store depletion patterns during spontaneous sleep apnoeas of young northern elephant seals revealed near-complete depletion of the blood oxygen store but 80%–90% preservation of the muscle oxygen store.<sup>64,113</sup> Arterial and venous haemoglobin (Hb) data were obtained with intermittent blood sampling during multiple breath holds of different animals; myoglobin (Mb) saturation was determined with  $^1H$  NMR spectroscopy (adapted from Stockard *et al*<sup>64</sup> and Ponganis *et al*<sup>113</sup>).

LDH1 and 2 isoenzymes (lactate oxidation), primarily located in glial cells, (D) increased gene expression of S100B (a stress protein with calcium binding activity), clusterin (an extracellular chaperone molecule) and most glycolytic enzymes, but decreased expression of pyruvate dehydrogenase, and (E) normal neuroglobin and cytochrome oxidase gene expression, but, in contrast to terrestrial mammals, located in glial cells not neurons.<sup>68–74</sup> Thus, in the seal in contrast to terrestrial mammals, most oxidative activity and consumption of lactate occurs in the astrocytes, not the neurons. Neither oxidative nor glycolytic activity appears enhanced. And, again in contrast to terrestrial mammals, there appears to be a reverse astrocyte-neuron lactate shuttle (astrocytes consume lactate produced by neurons).


Another remarkable aspect of neural function in deep-diving marine mammals is the apparent absence of high-pressure nervous syndrome. Recent molecular studies in cetaceans have revealed that substitutions in the N-methyl-D-aspartate receptor change its tertiary structure and may alter activation of the receptor, thus preventing the neuronal hyperexcitability that occurs at high pressure in other animals, including humans.<sup>75</sup>

#### AVOIDANCE OF REPERFUSION INJURY

Absence of reperfusion injury in marine mammals was well documented more than 40 years ago during complete, warm ischaemia of seal kidneys.<sup>76</sup> The mechanisms responsible for such tolerance are still incompletely understood, but again are potentially relevant to ischaemic preconditioning, organ preservation for transplantation and avoidance of reperfusion injury during revascularisation procedures. In the seal heart, although a 10-fold elevation in glycogen content may provide a large glycolytic energy store and prevent intracellular calcium accumulation during ischaemia/hypoxaemia,<sup>77</sup> an even more impressive 25-fold elevation in glutathione content should enhance the potential for scavenging of reactive oxygen species (ROS) during reperfusion.<sup>78</sup> Significant elevations in glutathione content were



**Figure 7** During routine 20 min dives at sea, elephant seals consistently experienced arterial haemoglobin saturations between 80% and 20%.<sup>60</sup> The seal's arterial saturation was at or below the common clinical criterion (90%) for ventilation and inspired oxygen concentration adjustments in ventilated patients, below the supplemental nasal cannula oxygen treatment threshold for patients with severe lung disease, below the human shallow water blackout level and, by the end of the dive, even less than the arterial saturations of humans breathing ambient air on Mt Everest (adapted from Grocott *et al*<sup>109</sup>, Lindholm and Lundgren<sup>110</sup>, Mason *et al*<sup>111</sup> and Nunn<sup>112</sup>).



**Figure 9** Carboxyhaemoglobin levels in non-smokers, heavy smokers (40 cigarettes per day), children with haemolytic diseases, patients with haemoglobin Zurich and northern elephant seals.<sup>100 114–116</sup> Data are mean values from the studies.

also found in seal kidney, lung and muscle. In all tissues, enzymes associated with the recycling of glutathione were elevated. In addition, multiple other ROS-scavenging enzyme activities are elevated in many tissues of both seals and penguins.<sup>79–81</sup> Lastly, in marine mammal muscle, the elevated myoglobin concentrations may also affect both ROS scavenging and production.<sup>82 83</sup> The peroxidase activity of myoglobin can break down hydrogen peroxide. The nitrite reductase activity of myoglobin may increase nitric oxide production, which, in turn, can inhibit cytochrome c oxidase and decrease ROS formation.

It has also been found that the whole blood inflammatory response of seals on exposure to a potent endotoxin (lipopolysaccharide, LPS) is significantly blunted.<sup>84</sup> Interleukin-6 cytokine production in blood was 50–100 times lower in elephant seals and Weddell seals than in humans. In cell cultures of mononuclear cells from Weddell seal blood, addition of seal serum to the culture media decreased the LPS-induced production of interleukin-6 and other cytokines. It thus appears that seal blood contains an anti-inflammatory compound that may also play a role in protection against ischaemic and hypoxaemic insult. The mechanism(s) of this anti-inflammatory response in seals may represent another route to explore to minimise inflammatory responses in various clinical situations, including acute lung injury and organ transplantation (as emphasised by those authors), and also in others (ie, during cardiopulmonary bypass).<sup>85</sup> Changes in granulocyte and monocyte function induced by pressure exposure have also been reported in the beluga, a deep-diving whale.<sup>86</sup> It was postulated that such alterations in immune function might decrease susceptibility to decompression sickness.

### NITRIC OXIDE, HYDROGEN SULFIDE AND CARBON MONOXIDE

The diving physiology of marine mammals provides an opportunity to examine the functional roles and therapeutic potential of the gasotransmitters, nitric oxide, hydrogen sulfide and carbon monoxide.<sup>87–89</sup> These intracellular messengers have multiple physiological effects including vasodilatation. In that regard, it is notable that seals, which have common, spontaneous (non-obstructive) sleep apnoeas, do not develop pulmonary hypertension

despite the routine hypoxia and hypercarbia that occur during these 6–12 min breath holds.<sup>90</sup>

Increased nitric oxide production in the blood in marine mammals has been postulated because of elevated blood nitrite levels in porpoises and the high Hb concentrations found in many marine mammals.<sup>91</sup> Nitrite is a potential substrate for nitric oxide formation by deoxyhaemoglobin due to deoxyhaemoglobin's nitrite reductase activity.<sup>92–94</sup> The presence of exhaled nitric oxide in porpoises supported this hypothesis.<sup>95</sup> However, nitric oxide was absent in exhaled air of Weddell seals.<sup>96</sup> Falke *et al* suggested that the absence of exhaled nitric oxide may be secondary to reduced nitric oxide production in the seal, although another possible cause was the lack of paranasal sinuses in seals as paranasal sinuses are considered to contribute significantly to nitric oxide levels in exhaled air. Further examination of the role of nitric oxide and nitrite in these animals is needed, and is particularly relevant to current investigations of nitrite therapy in hypoxaemic and ischaemic disease.<sup>97</sup>

Hydrogen sulfide may also play a role in the prevention of pulmonary hypertension in marine mammals. During *in vitro* exposure to hypoxia, the pulmonary arteries of sea lions exhibit hypoxic pulmonary vasodilation, not hypoxic pulmonary vasoconstriction.<sup>98</sup> Vasodilatation also occurs in response to addition of hydrogen sulfide to the tissue bath. In contrast, bovine pulmonary arteries constrict on exposure to either hypoxia or hydrogen sulfide. It has been postulated that hydrogen sulfide accumulates within smooth muscle cells on exposure to hypoxia in both species, but that the contractile response to this intracellular messenger has changed in the sea lion to that of vasodilatation rather than vasoconstriction as in terrestrial mammals.

Carboxyhaemoglobin levels as high as 10% in elephant seals and elevated blood carbon monoxide levels in Weddell seals raise the possibility of a role for carbon monoxide in vascular regulation and prevention of reperfusion injury in these deep-diving seals.<sup>99 100</sup> Carboxyhaemoglobin levels were as high as in smokers (figure 9). These high carbon monoxide levels have been attributed to generation of carbon monoxide in the metabolism of haem to bilirubin, and to the high Hb concentrations (25 g/dL), blood volumes (200–220 mL/kg body mass) and myoglobin concentrations (5.4–7.8 g 100/g muscle) in these seals.<sup>101</sup> Although paradoxical in that total blood oxygen stores would be decreased by the binding of carbon monoxide to Hb, the left shift (increased O<sub>2</sub> affinity) of the Hb dissociation curve induced by carbon monoxide may be beneficial and actually increase blood oxygen content at low arterial pO<sub>2</sub>.

In this era of the design of carbon monoxide releasing molecules and the potential utilisation of carbon monoxide as a therapeutic agent in vascular regulation and prevention of apoptosis/inflammatory responses during reperfusion,<sup>88 102–104</sup> these seals are potentially excellent models for investigation of carbon monoxide's function. The high concentrations of carbon monoxide in seals may play a significant role in the diving physiology of the seal, especially in vascular regulation and hypoxaemic/ischaemic protection.

### CONCLUSIONS

In conclusion, the exceptional breath-hold capacities and dive behaviours of many marine mammals and seabirds are dependent on anatomy and physiology. Investigation of these adaptations is relevant to the understanding of their ecology and their responses to environmental challenges. Such research could benefit from collaboration of field biologists, comparative physiologists and medical investigators. Furthermore, the hypoxaemic

and pressure tolerance of these animals and the avoidance of reperfusion injury provide the opportunity to illustrate basic physiological principles and explore potential therapeutic applications.

**Funding** This study was funded by the National Science Foundation Division of Polar Programs (PLR-1643532).

**Competing interests** None declared.

**Patient consent for publication** Not required.

**Provenance and peer review** Commissioned; externally peer reviewed.

## REFERENCES

- Ponganis PJ. *Diving physiology of marine mammals and seabirds*. Cornwall: Cambridge University Press, 2015.
- Falcone EA, Schorr GS, Watwood SL, et al. Diving behaviour of Cuvier's beaked whales exposed to two types of military sonar. *R Soc Open Sci* 2017;4.
- Schorr GS, Falcone EA, Moretti DJ, et al. First long-term behavioral records from Cuvier's beaked whales (*Ziphius cavirostris*) reveal record-breaking dives. *PLoS One* 2014;9.
- Hamilton J, Moodie D, Levy J. The use of the diving reflex to terminate supraventricular tachycardia in a 2-week-old infant. *Am Heart J* 1979;97:371-4.
- Shattock MJ, Tipton MJ. 'Autonomic conflict': a different way to die during cold water immersion? *J Physiol* 2012;590:3219-30.
- Piscitelli MA, Raverty SA, Lillie MA, et al. A review of cetacean lung morphology and mechanics. *J Morphol* 2013;274:1425-40.
- Kooyman GL, Cornell LH. Flow properties of expiration and inspiration in a trained Bottle-Nosed porpoise. *Physiol Zool* 1981;54:55-61.
- Kooyman GL, Sinnett EE. Mechanical properties of the harbor porpoise lung, *Phocoena phocoena*. *Respir Physiol* 1979;36:287-300.
- Kerem DH, Kylstra JA, Saltzman HA. Respiratory flow rates in the sea lion. *Undersea Biomedical Research* 1975;2:20-7.
- Camporesi EM, Ventilation BG. Gas Exchange and Exercise Under Pressure. In: Brubakk AO, Neuman TS, eds. *Physiology and medicine of diving*. Edinburgh: Saunders, 2003: 77-114.
- Olsen CR, Elsner R, Hale FC, et al. "Blow" of the pilot whale. *Science* 1969;163:953-5.
- Fahlman A, Loring SH, Levine G, et al. Lung mechanics and pulmonary function testing in cetaceans. *J Exp Biol* 2015;218:2030-8.
- Cozzi B, Bagnoli P, Accocella F, et al. Structure and biomechanical properties of the trachea of the striped dolphin *Stenella coeruleoalba*: evidence for evolutionary adaptations to diving. *Anat Rec A Discov Mol Cell Evol Biol* 2005;284:500-10.
- Davenport J, Cotter L, Rogan E, et al. Structure, material characteristics and function of the upper respiratory tract of the pygmy sperm whale. *J Exp Biol* 2013;216:4639-46.
- Denison DM, Kooyman GL. The structure and function of the small airways in pinniped and sea otter lungs. *Respir Physiol* 1973;17:1-10.
- Scholander PF. Experimental investigations on the respiratory function in diving mammals and birds. *Hvalradets Skrifter* 1940;22:1-131.
- Gompelmann D, Eberhardt R, Michaud G, et al. Predicting atelectasis by assessment of collateral ventilation prior to endobronchial lung volume reduction: a feasibility study. *Respiration* 2010;80:419-25.
- García Parraga D, Moore M, Fahlman A, et al. Pulmonary ventilation-perfusion mismatch: a novel hypothesis for how diving vertebrates may avoid the bends. *Proc R Soc B* 2018;285.
- Denison DM, Warrell DA, West JB. Airway structure and alveolar emptying in the lungs of sea lions and dogs. *Respir Physiol* 1971;13:253-60.
- Fahlman A, Loring SH, Ferrigno M, et al. Static inflation and deflation pressure-volume curves from excised lungs of marine mammals. *J Exp Biol* 2011;214:3822-8.
- Kooyman GL, Sinnett EE. Pulmonary shunts in harbor seals and sea lions during simulated dives to depth. *Physiological Zoology* 1982;55:105-11.
- Falke K, Hill R, Qvist J, et al. Seal lungs collapse during free diving: evidence from arterial nitrogen tensions. *Science* 1985;229:556-8.
- Kooyman GL, Schroeder JP, Denison DM, et al. Blood nitrogen tensions of seals during simulated deep dives. *Am J Physiol* 1972;223:1016-20.
- Ridgway SH, Howard R. Dolphin lung collapse and intramuscular circulation during free diving: evidence from nitrogen washout. *Science* 1979;206:1182-3.
- McDonald BI, Ponganis PJ. Lung collapse in the diving sea lion: hold the nitrogen and save the oxygen. *Biol Lett* 2012;8:1047-9.
- Fahlman A, Loring SH, Johnson SP, et al. Inflation and deflation pressure-volume loops in anesthetized pinnipeds confirms compliant chest and lungs. *Front Physiol* 2014;5.
- Fahlman A, Moore MJ, Garcia-Parraga D. Respiratory function and mechanics in pinnipeds and cetaceans. *J Exp Biol* 2017;220:1761-73.
- Robinson PW, Costa DP, Crocker DE, et al. Foraging behavior and success of a mesopelagic predator in the Northeast Pacific Ocean: insights from a data-rich species, the northern elephant seal. *PLoS One* 2012;7:e36728.
- Fahlman A, Hooker SK, Olszowka A, et al. Estimating the effect of lung collapse and pulmonary shunt on gas exchange during breath-hold diving: the Scholander and Kooyman legacy. *Respir Physiol Neurobiol* 2009;165:28-39.
- Spragg RG, Ponganis PJ, Marsh JJ, et al. Surfactant from diving aquatic mammals. *J Appl Physiol* 2004;96:1626-32.
- Gutierrez DB, Fahlman A, Gardner M, et al. Phosphatidylcholine composition of pulmonary surfactant from terrestrial and marine diving mammals. *Respir Physiol Neurobiol* 2015;211:29-36.
- Miller NJ, Daniels CB, Schürch S, et al. The surface activity of pulmonary surfactant from diving mammals. *Respir Physiol Neurobiol* 2006;150:220-32.
- Miller NJ, Postle AD, Orgeig S, et al. The composition of pulmonary surfactant from diving mammals. *Respir Physiol Neurobiol* 2006;152:152-68.
- Miller NJ, Postle AD, Schürch S, et al. The development of the pulmonary surfactant system in California sea lions. *Comp Biochem Physiol A Mol Integr Physiol* 2005;141:191-9.
- Foot NJ, Orgeig S, Daniels CB. The evolution of a physiological system: the pulmonary surfactant system in diving mammals. *Respir Physiol Neurobiol* 2006;154:118-38.
- Daniels CB, Orgeig S. Pulmonary surfactant: the key to the evolution of air breathing. *Physiology* 2003;18:151-7.
- Kjekshus JK, Blix AS. How seals avoid myocardial infarction during diving: do they show us a new approach to treatment of patients with ischaemic heart injury? *Scand J Clin Lab Invest* 1977;37:95-8.
- Ponganis PJ, Meir JU, Williams CL. In pursuit of Irving and Scholander: a review of oxygen store management in seals and penguins. *J Exp Biol* 2011;214:3325-39.
- Blix AS. Adaptations to deep and prolonged diving in phocid seals. *J Exp Biol* 2018;221. [Epub ahead of print: 22 Jun 2018].
- Elsner R, Millard RW, Kjekshus JK, et al. Coronary blood flow and myocardial segment dimensions during simulated dives in seals. *Am J Physiol* 1985;249:H1119-H1126.
- Kjekshus KJ, Blix AS, Elsner R, et al. Myocardial blood flow in the diving seal. *Amer J Physiol* 1982;242:R97-R104.
- Harrison RJ, Tomlinson JDW. Observations on the venous system in certain Pinnipedia and Cetacea. *Proceedings of the Zoological Society of London* 1956;126:205-34.
- Elsner R, Hanafee WN, Hammond DD. Angiography of the inferior vena cava of the harbor seal during simulated diving. *Amer J Physiol* 1971;220:1155-7.
- Ronald K, McCarter R, Selley LJ. Venous circulation of the Harp seal (*Pagophilus groenlandicus*). In: Harrison RJ, ed. *Functional anatomy of marine mammals*. New York: Academic Press, 1977: 235-70.
- Hol R, Blix AS, Myhre HO. Selective redistribution of the blood volume in the diving seal (*Pagophilus groenlandicus*). *Rapp. P.-v. Re'un. Cons. int. Explor. Mer* 1975;169:423-32.
- Drabek CM. Some anatomical aspects of the cardiovascular system of Antarctic seals and their possible functional significance in diving. *J Morphol* 1975;145:85-105.
- Rhode EA, Elsner R, Peterson TM, et al. Pressure-volume characteristics of aortas of harbor and Weddell seals. *Am J Physiol Regul Integr Comp Physiol* 1986;251:R174-R180.
- Thornton SJ, Pelc NJ, Spielman JR, et al. Vascular flow dynamics in a diving elephant seal (*Mirovanga angustirostris*). *Int Soc Magn Reson Med* 1997;2.
- Papaioannou TG, Stefanadis C. Basic principles of the intraaortic balloon pump and mechanisms affecting its performance. *Asaio J* 2005;51:296-300.
- Goldbogen JA, Shadwick RE, Lillie MA, et al. Using morphology to infer physiology: case studies on orca whales (Balaenopteridae). *Can J Zool* 2015;93:687-700.
- Shadwick RE, Gosline JM. Arterial mechanics in the fin whale suggest a unique hemodynamic design. *Am J Physiol Regul Integr Comp Physiol* 1994;267:R805-R818.
- Geng X, Dufu K, Hutchaleelaha A, et al. Increased hemoglobin-oxygen affinity ameliorates bleomycin-induced hypoxemia and pulmonary fibrosis. *Physiol Rep* 2016;4:e12965.
- Oksenberg D, Dufu K, Patel MP, et al. GBT440 increases haemoglobin oxygen affinity, reduces sickling and prolongs RBC half-life in a murine model of sickle cell disease. *Br J Haematol* 2016;175:141-53.
- Putz ND, Shaver CM, Dufu K, et al. GBT1118, a compound that increases the oxygen affinity of hemoglobin, improves survival in murine hypoxic acute lung injury. *J Appl Physiol* 2018;124:899-905.
- Black CP, Tenney SM. Oxygen transport during progressive hypoxia in high-altitude and sea-level waterfowl. *Respir Physiol* 1980;39:217-39.
- Meir JU, Ponganis PJ. High-affinity hemoglobin and blood oxygen saturation in diving emperor penguins. *J Exp Biol* 2009;212:3330-8.
- Milsum WK, Johansen K, Millard RW. Blood respiratory properties in some Antarctic birds. *Condor* 1973;75:472-4.
- Petschow D, Würdinger I, Baumann R, et al. Causes of high blood O<sub>2</sub> affinity of animals living at high altitude. *J Appl Physiol Respir Environ Exerc Physiol* 1977;42:139-43.
- Qvist J, Weber RE, Zapol WM. Oxygen equilibrium properties of blood and hemoglobin of fetal and adult Weddell seals. *J Appl Physiol* 1981;50:999-1005.

- 60 Meir JU, Champagne CD, Costa DP, *et al.* Extreme hypoxemic tolerance and blood oxygen depletion in diving elephant seals. *Am J Physiol Regul Integr Comp Physiol* 2009;297:R927–R939.
- 61 Storz JF. Hemoglobin-oxygen affinity in high-altitude vertebrates: is there evidence for an adaptive trend? *J Exp Biol* 2016;219:3190–203.
- 62 Elsner R, Shurley JT, Hammond DD, *et al.* Cerebral tolerance to hypoxemia in asphyxiated Weddell seals. *Respir Physiol* 1970;9:287–97.
- 63 Kerem D, Elsner R. Cerebral tolerance to asphyxial hypoxia in the harbor seal. *Respir Physiol* 1973;19:188–200.
- 64 Stockard TK, Levenson DH, Berg L, *et al.* Blood oxygen depletion during rest-associated apneas of northern elephant seals (*Mirounga angustirostris*). *J Exp Biol* 2007;210:2607–17.
- 65 Blix AS, Walløe L, Messelt EB, *et al.* Selective brain cooling and its vascular basis in diving seals. *J Exp Biol* 2010;213:2610–6.
- 66 Odden A, Folkow LP, Caputa M, *et al.* Brain cooling in diving seals. *Acta Physiol Scand* 1999;166:77–8.
- 67 Geiseler SJ, Larson J, Folkow LP. Synaptic transmission despite severe hypoxia in hippocampal slices of the deep-diving hooded seal. *Neuroscience* 2016;334:39–46.
- 68 Czech-Damal NU, Geiseler SJ, Hoff ML, *et al.* The role of glycogen, glucose and lactate in neuronal activity during hypoxia in the hooded seal (*Cystophora cristata*) brain. *Neuroscience* 2014;275:374–83.
- 69 Fabrizio A, Hoff MLM, Engler G, *et al.* When the brain goes diving: transcriptome analysis reveals a reduced aerobic energy metabolism and increased stress proteins in the seal brain. *BMC Genomics* 2016;17:1–11.
- 70 Folkow LP, Ramirez JM, Ludvigsen S, *et al.* Remarkable neuronal hypoxia tolerance in the deep-diving adult hooded seal (*Cystophora cristata*). *Neurosci Lett* 2008;446:147–50.
- 71 Hoff MLM, Fabrizio A, Czech-Damal NU, *et al.* Transcriptome analysis identifies key metabolic changes in the hooded seal (*Cystophora cristata*) brain in response to hypoxia and reoxygenation. *PLoS One* 2017;12:e0169366.
- 72 Hoff ML, Fabrizio A, Folkow LP, *et al.* An atypical distribution of lactate dehydrogenase isoenzymes in the hooded seal (*Cystophora cristata*) brain may reflect a biochemical adaptation to diving. *J Comp Physiol B* 2016;186:373–86.
- 73 Mitz SA, Reuss S, Folkow LP, *et al.* When the brain goes diving: glial oxidative metabolism may confer hypoxia tolerance to the seal brain. *Neuroscience* 2009;163:552–60.
- 74 Schreiner M, Flachsbarth S, Czech-Damal NU, *et al.* Neuroglobin of seals and whales: evidence for a divergent role in the diving brain. *Neuroscience* 2012;223:35–44.
- 75 Bliznyuk A, Golan H, Grossman Y. Marine mammals' NMDA receptor structure: possible adaptation to high pressure environment. *Front Physiol* 2018;9.
- 76 Halasz NA, Elsner R, Garvie RS, *et al.* Renal recovery from ischemia: a comparative study of harbor seal and dog kidneys. *Am J Physiol* 1974;227:1331–5.
- 77 Henden T, Aasum E, Folkow L, *et al.* Endogenous glycogen prevents Ca<sup>2+</sup> overload and hypercontracture in HARP seal myocardial cells during simulated ischemia. *J Mol Cell Cardiol* 2004;37:43–50.
- 78 Vázquez-Medina JP, Zenteno-Savín T, Elsner R. Glutathione protection against dive-associated ischemia/reperfusion in Ringed seal tissues. *J Exp Mar Bio Ecol* 2007;345:110–8.
- 79 Vázquez-Medina JP, Zenteno-Savín T, Elsner R. Antioxidant enzymes in Ringed seal tissues: potential protection against dive-associated ischemia/reperfusion. *Comp Biochem Physiol A Mol Integr Physiol Part C* 2006;142:198–204.
- 80 Zenteno-Savín T, St. Leger J, Ponganis PJ. Hypoxemic and ischemic tolerance in emperor penguins. *Comp Biochem Physiol A Mol Integr Physiol Part C* 2010;152:18–23.
- 81 Zenteno-Savín T, Vázquez-Medina JP, Cantu-Medellín N, *et al.* Ischemia/reperfusion in diving birds and mammals: how they avoid oxidative damage. In: Zenteno-Savín T, Vázquez-Medina JP, Abele D, *et al.*, eds. *Oxidative stress in aquatic ecosystems*. Oxford: Wiley-Blackwell, 2011: 178–89.
- 82 Helbo S, Fago A. Functional properties of myoglobins from five whale species with different diving capacities. *J Exp Biol* 2012;215:3403–10.
- 83 Helbo S, Weber RE, Fago A. Expression patterns and adaptive functional diversity of vertebrate myoglobins. *Biochimica et Biophysica Acta (BBA) - Proteins and Proteomics* 1834;2013:1832–9.
- 84 Bagchi A, Batten AJ, Levin M, *et al.* Intrinsic anti-inflammatory properties in the serum of two species of deep-diving seal. *J Exp Biol* 2018;221. doi:10.1242/jeb.178491. [Epub ahead of print: 09 Jul 2018].
- 85 Warren OJ, Smith AJ, Alexiou C, *et al.* The inflammatory response to cardiopulmonary bypass: part 1—mechanisms of pathogenesis. *J Cardiothorac Vasc Anesth* 2009;23:223–31.
- 86 Thompson LA, Romano TA. Beluga (*Delphinapterus leucas*) granulocytes and monocytes display variable responses to in vitro pressure exposures. *Front Physiol* 2015;6.
- 87 Lundberg JO, Weitzberg E, Gladwin MT. The nitrate-nitrite-nitric oxide pathway in physiology and therapeutics. *Nat Rev Drug Discov* 2008;7:156–67.
- 88 Motterlini R, Otterbein LE. The therapeutic potential of carbon monoxide. *Nat Rev Drug Discov* 2010;9:728–43.
- 89 Szabó C. Hydrogen sulphide and its therapeutic potential. *Nat Rev Drug Discov* 2007;6:917–35.
- 90 Ponganis PJ, Stockard TK, Levenson DH, *et al.* Intravascular pressure profiles in elephant seals: hypotheses on the caval sphincter, extradural vein and venous return to the heart. *Comp Biochem Physiol A Mol Integr Physiol* 2006;145:123–30.
- 91 Soegaard LB, Hansen MN, van Elk C, *et al.* Respiratory properties of blood in the harbor porpoise, *Phocoena phocoena*. *J Exp Biol* 2012;215:1938–43.
- 92 Flögel U, Fago A, Rassaf T. Keeping the heart in balance: the functional interactions of myoglobin with nitrogen oxides. *J Exp Biol* 2010;213:2726–33.
- 93 Lei J, Vodovotz Y, Tzeng E, *et al.* Nitric oxide, a protective molecule in the cardiovascular system. *Nitric Oxide* 2013;35:175–85.
- 94 Fago A, Parraga DG, Petersen EE, *et al.* A comparison of blood nitric oxide metabolites and hemoglobin functional properties among diving mammals. *Comp Biochem Physiol A Mol Integr Physiol* 2017;205:35–40.
- 95 Yeates LC, Carlin KP, Baird M, *et al.* Nitric oxide in the breath of bottlenose dolphins: effects of breath hold duration, feeding, and lung disease. *Marine Mammal Science* 2014;30:272–81.
- 96 Falke KJ, Busch T, Hoffmann O, *et al.* Breathing pattern, CO<sub>2</sub> elimination and the absence of exhaled NO in freely diving Weddell seals. *Respir Physiol Neurobiol* 2008;162:85–92.
- 97 Calvert JW, Lefer DJ. Clinical translation of nitrite therapy for cardiovascular diseases. *Nitric Oxide* 2010;22:91–7.
- 98 Olson KR, Whitfield NL, Bearden SE, *et al.* Hypoxic pulmonary vasodilation: a paradigm shift with a hydrogen sulfide mechanism. *Am J Physiol Regul Integr Comp Physiol* 2010;298:R51–R60.
- 99 Pugh LG. Carbon monoxide content of the blood and other observations on Weddell seals. *Nature* 1959;183:74–6.
- 100 Tift MS, Ponganis PJ, Crocker DE. Elevated carboxyhemoglobin in a marine mammal, the northern elephant seal. *J Exp Biol* 2014;217:1752–7.
- 101 Hassrick JL, Crocker DE, Teutschel NM, *et al.* Condition and mass impact oxygen stores and dive duration in adult female northern elephant seals. *J Exp Biol* 2010;213:585–92.
- 102 Kevin LG, Laffey JG. Carbon monoxide: from poison to therapy for cardiopulmonary bypass-induced lung injury? *Anesthesiology* 2008;108:977–8.
- 103 Nakao A, Neto JS, Kanno S, *et al.* Protection against ischemia/reperfusion injury in cardiac and renal transplantation with carbon monoxide, biliverdin and both. *Am J Transplant* 2005;5:282–91.
- 104 Romão CC, Blättler WA, Seixas JD, *et al.* Developing drug molecules for therapy with carbon monoxide. *Chem Soc Rev* 2012;41:3571–83.
- 105 McDonald BI, Ponganis PJ. Deep-diving sea lions exhibit extreme bradycardia in long-duration dives. *J Exp Biol* 2014;217:1525–34.
- 106 Wright AK, Ponganis KV, McDonald BI, *et al.* Heart rates of emperor penguins diving at sea: implications for oxygen store management. *Marine Ecology Progress Series* 2014;496:85–98.
- 107 Hyatt RE, Schilder DP, Fry DL. Relationship between maximum expiratory flow and degree of lung inflation. *J Appl Physiol* 1958;13:331–6.
- 108 Hochachka PW. Pinniped diving response mechanism and evolution: a window on the paradigm of comparative biochemistry and physiology. *Comp Biochem Physiol A Mol Integr Physiol* 2000;126:435–58.
- 109 Grocott MP, Martin DS, Levett DZ, *et al.* Arterial blood gases and oxygen content in climbers on Mount Everest. *N Engl J Med* 2009;360:140–9.
- 110 Lindholm P, Lundgren CE. The physiology and pathophysiology of human breath-hold diving. *J Appl Physiol* 2009;106:284–92.
- 111 Mason RJ, Murray JF, Broaddus VC, *et al.* *Murray and Nadel's Textbook of Respiratory Medicine*. Philadelphia: Elsevier Saunders, 2005.
- 112 Nunn JF. *Applied respiratory physiology*. 2nd edn. London: Butterworths, 1977.
- 113 Ponganis PJ, Kreutzer U, Stockard TK, *et al.* Blood flow and metabolic regulation in seal muscle during apnea. *J Exp Biol* 2008;211:3323–32.
- 114 Engel RR, Rodkey FL, Krill CE. Carboxyhemoglobin levels as an index of hemolysis. *Pediatrics* 1971;47:723–30.
- 115 Law MR, Morris JK, Watt HC, *et al.* The dose-response relationship between cigarette consumption, biochemical markers and risk of lung cancer. *Br J Cancer* 1997;75:1690–3.
- 116 Zinkham WH, Houtchens RA, Caughey WS. Carboxyhemoglobin levels in an unstable hemoglobin disorder (Hb Zürich): effect on phenotypic expression. *Science* 1980;209:406–8.