



OPEN ACCESS

Impact of banning smoking in cars with children on exposure to second-hand smoke: a natural experiment in England and Scotland

Anthony A Laverty ¹, Thomas Hone,¹ Eszter P Vamos,¹ Philip E Anyanwu,² David Taylor-Robinson,³ Frank de Vocht,⁴ Christopher Millett,¹ Nicholas S Hopkinson ⁵

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/thoraxjnl-2019-213998>).

¹Public Health Policy Evaluation Unit, School of Public Health, Imperial College London, London, UK

²Global Digital Health Unit, Department of Primary Care and Public Health, Imperial College London, London, UK

³Department of Public Health and Policy, University of Liverpool, Liverpool, UK

⁴Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK

⁵National Heart and Lung Institute, Imperial College London, London, UK

Correspondence to

Dr Anthony A Laverty, Dept Primary Care and Public Health, Imperial College, London W6 8RP, UK; a.laverty@imperial.ac.uk

Received 21 August 2019
Revised 4 November 2019
Accepted 5 November 2019
Published Online First
27 January 2020



Listen to Podcast
thorax.bmj.com



► <http://dx.doi.org/10.1136/thoraxjnl-2020-214660>



© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY. Published by BMJ.

To cite: Laverty AA, Hone T, Vamos EP, et al. *Thorax* 2020;**75**:345–347.

ABSTRACT

England banned smoking in cars carrying children in 2015 and Scotland in 2016. We used survey data from 3 years for both countries ($N_{\text{England}}=3483-6920$, $N_{\text{Scotland}}=232-319$) to assess effects of the English ban using logistic regression within a difference-in-differences framework. Among children aged 13–15 years, self-reported levels of regular exposure to smoke in cars for Scotland were 3.4% in 2012, 2.2% in 2014 and 1.3% in 2016 and for England 6.3%, 5.9% and 1.6%. The ban in England was associated with a -4.1% (95% CI -4.9% to -3.3%) absolute reduction (72% relative reduction) in exposure to tobacco smoke among children.

INTRODUCTION

Exposure to secondhand tobacco smoke is a significant cause of illness in children and particularly affects more disadvantaged groups.¹ Exposure of children to smoking inside cars is especially concerning owing to the very high concentrations of smoke in enclosed spaces and associations with child smoking uptake.^{2,3} Both smoking uptake and secondhand exposure to smoke are mechanisms which sustain health inequality.^{4–6}

One policy response has been bans on smoking in private vehicles with children present. Evaluations of such bans are scarce and present a mixed picture.^{7–9} A ban on smoking in private vehicles with anyone aged ≤ 18 years present came into effect on 1 October 2015 in England, and on 5 December 2016 in Scotland, providing a natural experiment to evaluate policy effects.

METHODS

Data for England came from the Smoking, Drinking and Drug Use (SDDU) surveys and for Scotland from the Scottish health surveys in 2012, 2014 and 2016 (online supplementary appendix 1). We restricted the sample to children aged 13–15, as exposure for 11 and 12 year olds was reported by caregivers, which is likely to lead to under-recording.

Our primary exposure was child-reported regular exposure to smoking inside cars, assessed, in England, with the question “In the past year, how often were you in a car with somebody smoking?” (table 1). We categorised responses of *every day or most days* as ‘regular exposure’. In Scotland, children were asked “Are you regularly exposed to other

people’s tobacco smoke in any of these places?” (Responses: yes/no for cars/vehicles). We harmonised a measure of deprivation using free school meals and the Family Affluence Scale and include this as well as age and sex in analyses (table 1).

We used survey-weighted logistic regression to assess changes in exposure over time using a differences-in-differences analysis, which controls for all time-invariant differences between the intervention and comparison populations.¹⁰ We considered interaction between time and country to assess changes in each country separately and with policy implementation in England in 2016. We present relative percentage changes (based on odds ratios) and absolute percentage changes (average marginal effects compared with baseline).

We conducted various additional analyses: an unadjusted model (online supplementary appendix table 2), analyses with linear trends for each country (online supplementary appendix table 3) and analyses for England only (ages 11–15), using more granular exposure data to analyse ever, monthly and regular exposure (online supplementary appendix table 4).

RESULTS

A total of 15 318 responses were received in England and 822 in Scotland (online supplementary appendix 1). Self-reported regular exposure to smoke in cars was 6.3% in 2012, 5.9% in 2014 and 1.6% in 2016 for England and, respectively, 3.4%, 2.2% and 1.3% in Scotland (figure 1).

Implementation of the smoke-free policy in England was associated with an absolute reduction of -4.1% (95% CI -4.9% to -3.3%) (72% relative reduction) in the percentage of children self-reporting exposure to smoke in cars (table 2). Girls and those in the deprived group were more likely to report exposure.

Analyses unadjusted for covariates and using linear time trends were similar (online supplementary appendix tables 2 and 3). Analyses within England only, using the wider age range of 11 to 15 years, identified substantial changes in exposure in all levels of exposure assessed after implementation of the ban (online supplementary appendix table 4).

DISCUSSION

Child-reported frequency of exposure to tobacco smoke in cars fell after the 2015 introduction of the ban in England, a finding made more robust by

Table 1 Key characteristics of data sources

Country	England	Scotland
Data name	Smoking Drinking Drug Use (SDDU) survey	Scottish health survey
Years and interview dates	2012: September 2012 – December 2012 2014: September 2014 – December 2014 2016: September 2016 – January 2017	2012: January 2012 – December 2012 2014: January 2014 – February 2015 2016: January 2016 – January 2017
Individuals included	2012: 4915 2014: 3483 2016: 6920	2012: 319 2014: 271 2016: 232
Age range	11–15 years	13–17 years
Exposure to tobacco smoke in cars question	2012: Two separate questions: "In the past year, how often were you in your family's car (OR someone else's car with somebody smoking?" (Responses: every day or most days; once or twice a week; once or twice a month; less often than once a month; never in the past year; don't know) 2014 and 2016: In the past year, how often were you in a car with somebody smoking? This could be your family's car or someone else's car. (Responses: every day or most days; once or twice a week; once or twice a month; less often than once a month; never in the past year; don't know) In this study 'every day or most days' classed as 'regular exposure'	2012–2016: "Are you regularly exposed to other people's tobacco smoke in any of these places?" (responses: yes/no for in cars/ vehicles, etc)
Deprivation marker	2012 and 2014: Child reported receipt of free school meals (FSM) 2016: Family Affluence Scale (low, middle, high). Harmonised so low groupe equivalent to receiving FSM	2012– 2016: Scottish index of multiple deprivation in five groups Harmonised so most deprived group comparable to receiving FSM or being in lowest affluence band

comparison with Scotland where the policy was not introduced until the following year. This exposure remains higher in children from more deprived groups, which serves as a reminder of the socially patterned risks of smoking. These findings provide support for introducing this policy in other jurisdictions, although further evaluations are warranted. Previous research from Canadian provinces enacting such bans found more marked effects on exposure in provinces with comprehensive strategies including discouraging smoking uptake, while recent evidence has additionally pointed to exposure to smoking in cars resulting in an increase in the incidence of asthma.^{8 9}

The ban is an example of the 'Protect' element of the MPOWER policy approach to delivering the WHO Framework Convention on Tobacco Control programme to reduce the harms caused by smoking. Importantly, the purpose of the ban is to reduce child exposure to tobacco smoke, for which this study provides evidence, and not to drive prosecutions (eg, coverage presenting the legislation as a failure due to the low number of arrests <https://www.mirror.co.uk/news/uk-news/car-smoking-ban-massive-flop-10858407>).




Figure 1 Percentages of child-reported regular exposure to smoke in cars in England and Scotland 2012–2016

Strengths and limitations

The strength of this experiment using high-quality data from Scotland and England is that the design permits observed changes to be plausibly ascribed to the policy intervention. However, differences in the data used between countries deserve consideration. Different measurements of exposure in England and Scotland mean that we cannot directly compare exposure levels between countries, and the measure of deprivation we use has been harmonised from measures which have not previously been combined. However, analyses of data for England alone, using different levels of exposure, and not including the deprivation measure, all produced concordant results. Combining data sources also meant that we could employ a difference-in-difference design, which controls for time-invariant differences between countries and provides stronger evidence than relying

Table 2 Results from logistic regression difference in difference analyses of impact of policy implementation on self-reported exposure to smoking in vehicles

	AOR (95% CI)	Absolute % difference (95% CI)
Scotland 2012–2014	0.34 (0.15 to 0.80)	-0.45 (-2.70 to 1.80)
Scotland 2014–2016	0.25 (0.07 to 0.79)	0.00 (-1.05 to 1.06)
England 2012–2014	1.00 (0.82 to 1.22)	-1.04 (-3.34 to 1.27)
England 2014–2016 (policy implementation)	0.28 (0.21 to 0.37)	-4.11 (-4.91 to -3.31)
Age 13 years	ref	ref
Age 14 years	1.01 (0.80 to 1.28)	0.04 (-0.77 to 0.84)
Age 15 years	1.22 (0.99 to 1.51)	0.74 (-0.05 to 1.53)
Boys	ref	ref
Girls	1.61 (1.34 to 1.93)	1.75 (1.07 to 2.43)
Not deprived group	ref	ref
Deprived group	1.98 (0.82 to 1.21)	2.53 (1.76 to 3.29)

Absolute % differences derived from marginal effect compared with baseline. AOR, adjusted odds ratio.

on England alone. Exposure was based on self-report only, and reporting bias might have changed over time, although this would probably have been similar in both countries. Interview dates were not available for the Scottish data, and sampling in 2016 included almost 2 months after the introduction of their ban. Misclassification from sampling dates would probably bias associations towards the null point, but the direction of any biases due to harmonising data sources is difficult to predict. Reported frequencies of exposure frequency in Scotland were low, meaning that there might have been some floor effects and, as these analyses use only three data points from each country, future analyses with more data are recommended and may provide discrepant results.

CONCLUSIONS

Our results suggest that banning smoking in private vehicles carrying children has been successful in its main aim of reducing their exposure to tobacco smoke. Given children's known vulnerability to secondhand smoke, reductions in exposure will probably result in improved health.

Twitter Nicholas S Hopkinson @COPDdoc

Contributors AAL, CM and NSH conceived the study idea. Analyses were by AAL with TH. All authors provided intellectual input to the interpretation of results and approved the final manuscript.

Funding This study is funded by the National Institute for Health Research (NIHR) School for Public Health Research (grant reference number PD-SPH-2015). The views expressed are those of the author(s) and not necessarily those of the NIHR or the Department of Health and Social Care. The funder had no input in the writing of the manuscript or decision to submit for publication. The NIHR School for Public Health Research is a partnership between the Universities of Sheffield; Bristol; Cambridge; Imperial; and University College London; The London School for Hygiene and Tropical Medicine (LSHTM); LiLaC – a collaboration between the Universities of Liverpool and Lancaster; and Fuse - The Centre for Translational Research in Public Health a collaboration between Newcastle, Durham, Northumbria, Sunderland and Teesside Universities. DT-R is funded by the MRC on a clinician scientist fellowship (MR/P008577/1).

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

ORCID iDs

Anthony A Laverty <http://orcid.org/0000-0003-1318-8439>

Nicholas S Hopkinson <http://orcid.org/0000-0003-3235-0454>

REFERENCES

- 1 Royal College of Physicians. Passive smoking and children, 2010. Available: <https://cdn.shopify.com/s/files/1/0924/4392/files/passive-smoking-and-children.pdf?15599436013786148553> [Accessed 27 Jul 2019].
- 2 Sendzik T, Fong GT, Travers MJ, et al. An experimental investigation of tobacco smoke pollution in cars. *Nicotine Tob Res* 2009;11:627–34.
- 3 Okoli CTC, Kodet J. A systematic review of secondhand tobacco smoke exposure and smoking behaviors: smoking status, susceptibility, initiation, dependence, and cessation. *Addict Behav* 2015;47:22–32.
- 4 Laverty AA, Filippidis FT, Taylor-Robinson D, et al. Smoking uptake in UK children: analysis of the UK millennium cohort study. *Thorax* 2019;74:607–10.
- 5 Hopkinson NS, Lester-George A, Ormiston-Smith N, et al. Child uptake of smoking by area across the UK. *Thorax* 2014;69:873–5.
- 6 Jones LL, Moodie C, MacKintosh AM, et al. Young people's exposure to and perceptions of smoking in cars and associated harms in the United Kingdom. *Drugs Educ Prev Policy* 2014;21:225–33.
- 7 Nguyen HV. Do smoke-free car laws work? Evidence from a quasi-experiment. *J Health Econ* 2013;32:138–48.
- 8 Elton-Marshall T, Leatherdale ST, Driezen P, et al. Do provincial policies banning smoking in cars when children are present impact youth exposure to secondhand smoke in cars? *Prev Med* 2015;78:59–64.
- 9 Patel M, Thai CL, Meng Y-Y, et al. Smoke-free car legislation and student exposure to smoking. *Pediatrics* 2018;141:S40–50.
- 10 Craig P, Katikireddi SV, Leyland A, et al. Natural experiments: an overview of methods, approaches, and contributions to public health intervention research. *Annu Rev Public Health* 2017;38:39–56.

Impact of banning smoking in cars with children on exposure to second-hand smoke: a natural experiment in England and Scotland - APPENDIX

Anthony A Laverty^{1*}, Thomas Hone¹, Philip E. Anyanwu², David Taylor Robinson³, Frank de Vocht⁴, Christopher Millett¹, Nicholas S Hopkinson⁵

¹ Public Health Policy Evaluation Unit, School of Public Health, Imperial College London, UK

² Global Digital Health Unit, School of Public Health, Imperial College London, London, UK

³ Centre for Public Health, Bristol Medical School, University of Bristol, Bristol, UK

⁴ Department of Public Health and Policy, University of Liverpool, Liverpool, England

⁵ National Heart and Lung Institute, Imperial College London, London, UK

* Room 322, Reynolds Building, Imperial College London, St Dunstan's Road, W6 8RP, London, England.

Appendix contents

1. Further information on data sources
2. Appendix table 1 of study sample
3. Appendix table 2 of analyses using linear trends for time
4. Appendix table 3 unadjusted analyses
5. Appendix table 4 of analyses in England only including children aged 11 – 15 years

Further information on data sources

The Smoking Drinking Drug Use (SDDU) is a survey of children in school years 7 – 11 (aged 11-15 years) and is used to monitor the performance of the Government tobacco strategy [1][2][3][4][5]. Data came from questionnaires administered to children at school in exam conditions and is designed to be representative of the gender, age, region and type of school in England. Data for Scotland come from the years 2012, 2014 and 2016, where children aged 13 years and older were asked to report their exposure to smoking in cars [6][7][8]. For children below this age, caregivers were asked to report exposure, but we have excluded this data due to concerns over the accuracy of caregiver reporting of exposure.

Both surveys use a using a multi-stage stratified design and we have applied survey weights as recommended in the reports from both the SDDU and the Scottish health survey. Both surveys recommend the use of one weighting variable for analyses. The SDDU recommends the use of one weighting variable (*pupilwt*) which accounts for unequal selection probabilities between regions of England, sex and school year [9]. The Scottish Health Survey also recommends the use of one survey weight in analyses of children (*cmint[year]wt*). This weight is designed to adjust for selection of only two children per household as well as differences between responding and non-responding households. These survey weights were used for all regression analyses, and analyses were conducted in Stata using the survey routine *svy: logistic*. There were differences in the collection of data on markers of deprivation over time in England. In 2012 and 2014 children were asked if they received Free School Meals (FSM), but this measure was not used in 2016. The 2016 data used the Family Affluence Scale which asks children how many computers, vehicles and bathrooms their family has and assigned them a band from low to high [10]. We have harmonised these two measures by considered those receiving FSM or in the low FAS band as deprived. Scottish data used the Scottish Index of Multiple Deprivation as a marker of deprivation, and we harmonised the data by using the most deprived group as equivalent to receiving FSM or being in the lowest FAS band.

Appendix table 1: Study sample

	England		Scotland	
	Overall N	% exposed	Overall N	% exposed
2012	4915	6.3	319	3.4
2014	3483	5.9	271	2.2
2016	6920	1.6	232	1.3
Age 13 years	5056	4.0	272	0.7
Age 14 years	4702	3.8	266	3.4
Age 15 years	5560	4.5	284	3.2
Boys	7617	3.3	429	1.4
Girls	7701	4.8	393	3.6
Not deprived group	13572	3.5	552	2.2
Deprived group	1746	8.8	270	3.0
Total	15318	4.1	822	2.4

Appendix Table 2: Results from analyses using linear trends for time

	AOR	Lower CI	Upper CI
Policy Implementation	0.27	0.19	0.41
Scotland trend	0.77	0.42	1.42
England Trend	1.01	0.82	1.21
Scotland baseline	ref	ref	ref
England baseline	2.35	1.28	4.30
Age 13 years	ref	ref	ref
Age 14 years	1.01	0.80	1.28
Age 15 years	1.22	0.99	1.51
Boys	ref	ref	ref
Girls	1.61	1.34	1.93
Not deprived group	ref	ref	ref
Deprived group	1.98	1.61	2.43

AOR = Adjusted Odds Ratio, CI = Confidence Interval

In these analyses we modelled a linear trend for time individually for both Scotland and England individually and a binary variable for 2016 in England as the one post-intervention data point, interacted with time.

Results as odds ratios from logistic regression

Appendix table 3 : Results from unadjusted analyses

	Absolute % difference	Lower CI	Upper CI
Scotland 2012 - 2014	-0.66	-3.35	2.03
Scotland 2014-2016	0.02	-1.06	1.09
England 2012 - 2014	-1.55	-4.14	1.03
England 2014 - 2016 (policy implementation)	-4.14	-4.94	-3.33

Appendix table 4: Difference in difference analyses in England only including children aged 11 – 15 years

	Absolute % difference	Lower CI	Upper CI
Regular exposure			
Change 2012 - 2014	-2.96	-4.87	-1.05
Change 2014 - 2016	-10.83	-12.99	-8.67
Monthly exposure			
Change 2012 - 2014	-3.93	-5.59	-2.26
Change 2014 - 2016	-11.33	-13.09	-9.56
Ever exposure			
Change 2012 - 2014	0.17	-0.66	1.01
Change 2014 - 2016	-3.86	-4.48	-3.24

These analyses use data from England only and are controlled for age (11 – 15 years, categorically), sex and a marker of deprivation and survey weights for non-response.

References

- 1 NHS Digital. Smoking, Drinking and Drug Use Among Young People in England. 2016.
- 2 Department of Health (2017) Towards a Smokefree Generation: A Tobacco Control Plan for England.
https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/630217/Towards_a_Smoke_free_Generation_-_A_Tobacco_Control_Plan_for_England_2017-2022__2_.pdf
- 3 National Centre for Social Research, National Foundation for Educational Research. (2013). Smoking, Drinking and Drug Use among Young People, 2012. [data collection]. UK Data Service. SN: 7393, <http://doi.org/10.5255/UKDA-SN-7393-1>.
- 4 NatCen Social Research. (2015). Smoking, Drinking and Drug Use among Young People, 2014. [data collection]. UK Data Service. SN: 7811, <http://doi.org/10.5255/UKDA-SN-7811-1>.
- 5 NHS Digital. (2018). Smoking, Drinking and Drug Use among Young People, 2016. [data collection]. UK Data Service. SN: 8320, <http://doi.org/10.5255/UKDA-SN-8320-1>.
- 6 ScotCen Social Research, University College London, Department of Epidemiology and Public Health, University of Glasgow, MRC/CSO Social and Public Health Sciences Unit. (2015). Scottish Health Survey, 2012. [data collection]. 3rd Edition. UK Data Service.
- 7 ScotCen Social Research. (2019). Scottish Health Survey, 2014. [data collection]. 3rd Edition. UK Data Service. SN: 7851, <http://doi.org/10.5255/UKDA-SN-7851-3>.
- 8 ScotCen Social Research. (2017). Scottish Health Survey, 2016. [data collection]. UK Data Service. SN: 8290, <http://doi.org/10.5255/UKDA-SN-8290-1>.
- 9 NHS England (2017). Smoking, Drinking and Drug Use among Young People in England 2016 Appendices <https://files.digital.nhs.uk/publication/j/b/sdd-2016-app1.pdf>.
- 10 Currie C, Molcho M, Boyce W, *et al*. Researching health inequalities in adolescents: The development of the Health Behaviour in School-Aged Children (HBSC) Family Affluence Scale. *Soc Sci Med* Published Online First: 2008. doi:10.1016/j.socscimed.2007.11.024

Impact of banning smoking in cars with children on exposure to second-hand smoke: a natural experiment in England and Scotland - APPENDIX

Anthony A Laverty^{1*}, Thomas Hone¹, Philip E. Anyanwu², David Taylor Robinson³, Frank de Vocht⁴,
Christopher Millett¹, Nicholas S Hopkinson⁵

¹ Public Health Policy Evaluation Unit, School of Public Health, Imperial College London, UK

² Global Digital Health Unit, School of Public Health, Imperial College London, London, UK

³ Centre for Public Health, Bristol Medical School, University of Bristol, Bristol, UK

⁴ Department of Public Health and Policy, University of Liverpool, Liverpool, England

⁵ National Heart and Lung Institute, Imperial College London, London, UK

* Room 322, Reynolds Building, Imperial College London, St Dunstan's Road, W6 8RP, London, England.

Appendix contents

1. Further information on data sources
2. Appendix table 1 of study sample
3. Appendix table 2 of analyses using linear trends for time
4. Appendix table 3 unadjusted analyses
5. Appendix table 4 of analyses in England only including children aged 11 – 15 years

Further information on data sources

The Smoking Drinking Drug Use (SDDU) is a survey of children in school years 7 – 11 (aged 11-15 years) and is used to monitor the performance of the Government tobacco strategy [1][2][3][4][5]. Data came from questionnaires administered to children at school in exam conditions and is designed to be representative of the gender, age, region and type of school in England. Data for Scotland come from the years 2012, 2014 and 2016, where children aged 13 years and older were asked to report their exposure to smoking in cars [6][7][8]. For children below this age, caregivers were asked to report exposure, but we have excluded this data due to concerns over the accuracy of caregiver reporting of exposure.

Both surveys use a using a multi-stage stratified design and we have applied survey weights as recommended in the reports from both the SDDU and the Scottish health survey. Both surveys recommend the use of one weighting variable for analyses. The SDDU recommends the use of one weighting variable (*pupilwt*) which accounts for unequal selection probabilities between regions of England, sex and school year [9]. The Scottish Health Survey also recommends the use of one survey weight in analyses of children (*cmint[year]wt*). This weight is designed to adjust for selection of only two children per household as well as differences between responding and non-responding households. These survey weights were used for all regression analyses, and analyses were conducted in Stata using the survey routine *svy: logistic*. There were differences in the collection of data on markers of deprivation over time in England. In 2012 and 2014 children were asked if they received Free School Meals (FSM), but this measure was not used in 2016. The 2016 data used the Family Affluence Scale which asks children how many computers, vehicles and bathrooms their family has and assigned them a band from low to high [10]. We have harmonised these two measures by considered those receiving FSM or in the low FAS band as deprived. Scottish data used the Scottish Index of Multiple Deprivation as a marker of deprivation, and we harmonised the data by using the most deprived group as equivalent to receiving FSM or being in the lowest FAS band.

Appendix table 1: Study sample

	England		Scotland	
	Overall N	% exposed	Overall N	% exposed
2012	4915	6.3	319	3.4
2014	3483	5.9	271	2.2
2016	6920	1.6	232	1.3
Age 13 years	5056	4.0	272	0.7
Age 14 years	4702	3.8	266	3.4
Age 15 years	5560	4.5	284	3.2
Boys	7617	3.3	429	1.4
Girls	7701	4.8	393	3.6
Not deprived group	13572	3.5	552	2.2
Deprived group	1746	8.8	270	3.0
Total	15318	4.1	822	2.4

Appendix Table 2: Results from analyses using linear trends for time

	AOR	Lower CI	Upper CI
Policy Implementation	0.27	0.19	0.41
Scotland trend	0.77	0.42	1.42
England Trend	1.01	0.82	1.21
Scotland baseline	ref	ref	ref
England baseline	2.35	1.28	4.30
Age 13 years	ref	ref	ref
Age 14 years	1.01	0.80	1.28
Age 15 years	1.22	0.99	1.51
Boys	ref	ref	ref
Girls	1.61	1.34	1.93
Not deprived group	ref	ref	ref
Deprived group	1.98	1.61	2.43

AOR = Adjusted Odds Ratio, CI = Confidence Interval

In these analyses we modelled a linear trend for time individually for both Scotland and England individually and a binary variable for 2016 in England as the one post-intervention data point, interacted with time.

Results as odds ratios from logistic regression

Appendix table 3 : Results from unadjusted analyses

	Absolute % difference	Lower CI	Upper CI
Scotland 2012 - 2014	-0.66	-3.35	2.03
Scotland 2014-2016	0.02	-1.06	1.09
England 2012 - 2014	-1.55	-4.14	1.03
England 2014 - 2016 (policy implementation)	-4.14	-4.94	-3.33

Appendix table 4: Difference in difference analyses in England only including children aged 11 – 15 years

	Absolute % difference	Lower CI	Upper CI
Regular exposure			
Change 2012 - 2014	-2.96	-4.87	-1.05
Change 2014 - 2016	-10.83	-12.99	-8.67
Monthly exposure			
Change 2012 - 2014	-3.93	-5.59	-2.26
Change 2014 - 2016	-11.33	-13.09	-9.56
Ever exposure			
Change 2012 - 2014	0.17	-0.66	1.01
Change 2014 - 2016	-3.86	-4.48	-3.24

These analyses use data from England only and are controlled for age (11 – 15 years, categorically), sex and a marker of deprivation and survey weights for non-response.

References

- 1 NHS Digital. Smoking, Drinking and Drug Use Among Young People in England. 2016.
- 2 Department of Health (2017) Towards a Smokefree Generation: A Tobacco Control Plan for England.
https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/630217/Towards_a_Smoke_free_Generation_-_A_Tobacco_Control_Plan_for_England_2017-2022__2_.pdf
- 3 National Centre for Social Research, National Foundation for Educational Research. (2013). Smoking, Drinking and Drug Use among Young People, 2012. [data collection]. UK Data Service. SN: 7393, <http://doi.org/10.5255/UKDA-SN-7393-1>.
- 4 NatCen Social Research. (2015). Smoking, Drinking and Drug Use among Young People, 2014. [data collection]. UK Data Service. SN: 7811, <http://doi.org/10.5255/UKDA-SN-7811-1>.
- 5 NHS Digital. (2018). Smoking, Drinking and Drug Use among Young People, 2016. [data collection]. UK Data Service. SN: 8320, <http://doi.org/10.5255/UKDA-SN-8320-1>.
- 6 ScotCen Social Research, University College London, Department of Epidemiology and Public Health, University of Glasgow, MRC/CSO Social and Public Health Sciences Unit. (2015). Scottish Health Survey, 2012. [data collection]. 3rd Edition. UK Data Service.
- 7 ScotCen Social Research. (2019). Scottish Health Survey, 2014. [data collection]. 3rd Edition. UK Data Service. SN: 7851, <http://doi.org/10.5255/UKDA-SN-7851-3>.
- 8 ScotCen Social Research. (2017). Scottish Health Survey, 2016. [data collection]. UK Data Service. SN: 8290, <http://doi.org/10.5255/UKDA-SN-8290-1>.
- 9 NHS England (2017). Smoking, Drinking and Drug Use among Young People in England 2016 Appendices <https://files.digital.nhs.uk/publication/j/b/sdd-2016-app1.pdf>.
- 10 Currie C, Molcho M, Boyce W, *et al*. Researching health inequalities in adolescents: The development of the Health Behaviour in School-Aged Children (HBSC) Family Affluence Scale. *Soc Sci Med* Published Online First: 2008. doi:10.1016/j.socscimed.2007.11.024