OPEN ACCESS

ORIGINAL RESEARCH

Performance at stair-climbing test is associated with postoperative complications after lung resection: a systematic review and meta-analysis

Fairuz Boujibar,^{1,2} André Gillibert,³ Francis Edouard Gravier,^{4,5} Timothée Gillot,^{6,7} Tristan Bonnevie ,^{4,5} Antoine Cuvelier,^{5,8} Jean-marc Baste^{1,2}

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/thoraxjnl-2019-214019>).

For numbered affiliations see end of article.

Correspondence to

Fairuz Boujibar, Department of General and Thoracic Surgery, CHU Rouen, Rouen, France; fairuz.boujibar@chu-rouen.fr

Received 29 August 2019

Revised 15 May 2020

Accepted 5 June 2020

Published Online First

10 July 2020

ABSTRACT

Background Thoracic surgery is the optimal treatment for early-stage lung cancer, but there is a high risk of postoperative morbidity. Therefore, it is necessary to evaluate patients' preoperative general condition and cardiorespiratory capacity to determine the risk of postoperative complications. The objective of this study was to assess whether the stair-climbing test could be used in the preoperative evaluation of lung resection patients to predict postoperative morbidity following thoracic surgery.

Methods We performed a systematic review and a meta-analysis on the association between stair-climbing test result and morbidity/mortality after thoracic surgery. We analysed all articles published until May 2020 in the following databases: Pubmed/Medline, Pedro, The Cochrane library, Embase and CINAHL. The risk of bias was assessed using the Quality in Prognosis Studies tool. This meta-analysis is registered as PROSPERO CRD42019121348.

Results 13 articles were included in the systematic review for a total of 2038 patients and 6 in the meta-analysis. There were multiple test evaluation criteria: rise time, height, desaturation and heart rate change. For the meta-analysis, we were able to pool data on the height of rise at a variable threshold: risk ratio 2.34 (95% CI 1.59 to 3.43) with $I^2=53\%$ ($p=0.06$). The threshold for occurrence of complications was estimated at a 10 m climb.

Conclusions Our results indicate that the stair-climbing test could be used as a first-line functional screening test to predict postoperative morbidity following thoracic surgery and that patients with a poor test result (<10 m) should be referred to formal cardiopulmonary exercise testing.

INTRODUCTION

Surgery is considered the optimal treatment for patients with early stage lung cancer.¹ Nevertheless, lung resection remains a major intervention with significant morbidity.² Candidate selection influences morbidity and mortality; therefore, an appropriate assessment is primordial. According to the guidelines of the American College of Chest Physicians,³ the European Respiratory Society (ERS) and the European Society of Thoracic Surgeons (ESTS),⁴ all patients with a predicted postoperative value of diffusing capacity of the lung for carbon monoxide (DLCO) or forced expiratory volume in

Key messages**What is the key question?**

- Can stair-climbing test be used to predict postoperative complications in lung resection candidates?

What is the bottom line?

- This meta-analysis provides quality of evidence that a threshold of 10 m climbed during the stair-climbing test can predict postoperative morbidity.

Why read on?

- In this original systematic review and meta-analysis, we summarise current evidence of the role of the stair-climbing test in the preoperative assessment of patients referred for pulmonary resection.

1 s (FEV1) under 60% would benefit from further exercise testing as cardiopulmonary exercise testing (CPET). CPET on ergocycle with incremental exercise protocol is currently recognised as the gold standard stress test, allowing cardiopulmonary evaluation.⁵ It is essential to determine whether a patient is at risk after surgery or even operable and whether there is a need for prehabilitation to improve cardiorespiratory function before surgery.⁶

However, CPET is largely underused, with surgeons declaring that only between 10% and 30% of their patients receive this evaluation even though 75% of them prescribe CPET.⁷ As an alternative to CPET, some low-technology tests as the incremental shuttle walking test, the 6 min walk test (6MWT) and the stair-climbing test (SCT) have been described.⁸

For the SCT, patients are required to climb stairs to evaluate their cardiorespiratory function. In the context of thoracic surgery, this assessment is intended to identify patients at risk for postoperative complications. Indeed, there is a high risk of complications after pulmonary resection, especially if the patient is operated by thoracotomy.⁹ This test was first described in 1968¹⁰ and has been used for years. It is safe, easy to understand, fast and cheap. SCT can be performed as a submaximal test (a number of flights to climb) or as a maximal test (symptom limited test).⁸

► <http://dx.doi.org/10.1136/thoraxjnl-2020-214966>

© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Boujibar F, Gillibert A, Gravier FE, et al. *Thorax* 2020;**75**:791–797.

The objective of this study was to assess whether the SCT could be used in the preoperative evaluation of lung resection patients to predict postoperative complications.

METHODS

Protocol

A systematic review and meta-analysis were performed and are reported in accordance with the 'Preferred Reported Items for Systematic reviews and Meta-Analyses' (PRISMA) statement¹¹ and Cochrane Handbook for systematic reviews of diagnostic test accuracy.¹² A review protocol was written before starting the literature search.

The research protocol was prospectively published online at the PROSPERO International prospective register of systematic reviews (<http://www.crd.york.ac.uk/PROSPERO/>) under registration number: CRD42019121348.

Literature search

We queried Pubmed/Medline, Pedro, The Cochrane library, Embase and CINAHL in October 2018 and updated the search on May 2020. The search strategy is described in online supplementary material S1.

Eligibility criteria

Inclusion criteria were studies with a cohort design and investigating patients aged over 18 years who were diagnosed with lung cancer and who had SCT before surgery to predict postoperative complications. Studies with patients who had SCT for other reasons were excluded. There were no exclusion criteria based on language or publication date. For the analysis, we retained only studies in which SCT was performed in a maximal way, that is, the patient had to provide an uninterrupted maximal effort until exhaustion.

Outcome reporting

Since all studies that defined postoperative complications, defined it as any adverse event causing a cardiorespiratory disorder or death, this definition was kept for the meta-analysis. Since follow-up of complications was limited to hospital stay,¹³ up to 1 month,^{14 15} up to 3 months postoperatively¹⁶ or do not define the time limit,¹⁷ the maximum follow-up period of each article was used to define complications. Table 1 shows a high level of heterogeneity for categories of complications, with 98% of respiratory complications in Arruda¹⁸ vs 36% of respiratory and 59% of cardiac complications in Dong.¹⁹

Study selection and methodological quality assessment

Two independent reviewers screened and selected studies based on title, abstract and full text. The articles were translated if they were not reported in English or French.

Assessment of risk of bias was done by means of the Quality in Prognostic Studies (QUIPS) tool²⁰ used by Cochrane. A low risk of bias was given if all six domains were scored as low or if no more than two moderate or unknown risks of bias were identified. A moderate risk of bias was given if three or less than three domains were scored as moderate or unknown, in combination with no high risk of bias. A moderate risk of bias was also given if one domain was scored as high in combination with one or less than one moderate or unknown risk of bias. A high risk of bias was given when two or more domains were scored as high or four or more domains were scored as moderate or unknown risk of bias.^{4 5} The QUIPS assessment for each study

was independently completed by two authors. Differences were resolved by referral to a third author.

Data extraction and analysis

The following data were extracted from the studies: name of first author, publication year, country, study period, size of cohort, frequency of postoperative complications and crude number of complications in groups below and above a study-specific threshold separating low-risk and high-risk patients for each exposure. When there was no specified threshold, but detailed data were available, the median height was chosen as the threshold. When no threshold was defined and there were insufficient data to calculate the median or number of subjects above, the study was excluded from meta-analysis pooling.

Statistical analysis

Pooling effects of desaturation, heart rate change and rise time at the SCT was planned but cancelled because of heterogeneity and too few studies for each exposure.

Unadjusted relative risks (RRs) were pooled in a random effect (primary analysis) model by the DerSimonian-Laird method and a fixed effects model based on the Mantel-Haenszel estimator (sensitivity analysis). Heterogeneity was tested by Cochran's Q test²¹ and described by tau² and I². A fixed effects model based on the Mantel-Haenszel estimator for RRs was used in a sensitivity analysis.

Due to the threshold variance between studies, sensitivity and specificity are variable and negatively correlated, calling for joint interpretation. We modelled the bivariate distribution of the logit of sensitivity and logit of specificity in the five parameter model of Reitsma *et al.*²² mean sensitivity, mean specificity, between-study variance of sensitivity, between-study variance of specificity, between-study covariance of sensitivity and specificity. It is equivalent to the hierarchical summary receiver operating characteristic model.²³ It was estimated by restricted maximum likelihood by the 'mada' package of the R statistical software. Bivariate confidence regions and univariate CIs of sensitivity and specificity were generated from this model. Positive predictive value (PPV) and negative predictive values (NPV) were modelled in another Reitsma model, inverting roles between the test and the outcome as suggested by Leeftang *et al.*²⁴

All analyses were performed using R statistical software (V3.5.0, The R Foundation for Statistical Computing, Vienna, Austria).

RESULTS

Database searches and reference lists yielded a total of 168 articles (figure 1). After different removals, we identified 13 studies that were included in the systematic review.

Of the 13 cohort studies included, 8 studies described the climbed height as evaluation criteria, of which 2 gave means and SD in the group with and without complications but did not provide enough data to assess the RR of complication at a given threshold. Therefore, 6 of the 13 studies were included in the meta-analysis assessing the predictive value of the climbed height.

Of the studies included in the systematic review, four used desaturation as an evaluation criterion,^{19 25-27} but only two of them had data on RRs, and they used different definitions of desaturation. Consequently, results were not pooled for this evaluation criterion. Three articles used time to climb as evaluation criteria and two analysed heart rate but none of them had data on RR.

Table 1 Characteristics of included studies on systematic review and meta-analysis

Author, year, country	Participants	Age Mean (SD)	Interventions	FEV1 Mean (SD)	Maximal or submaximal test/variables reported	Type of complications	Surgical approach
Prospective cohort studies							
Ambrozio <i>et al.</i> , ²⁸ 2013, Brazil	98	52.7 (17.2)	Lobectomy/ pneumonectomy (repartition not specified)	2.3±0.8 L	Maximal/time	Respiratory: 75% Cardiac: 19% Death: 4% Other: 2%	Thoracotomy
Arruda ¹⁸ 2013, Brazil	48	54.0 (16.7)	All major lung resection without details	2.3±0.7 L	Maximal/time	Respiratory: 98% Death: 2%	Not specified
Brunelli, 2008, Italy ¹⁴	640	66.7 (9.3)	533 lobectomies/bilobectomies 107 pneumonectomies	85.4%±18.8%	Maximal/climbed height	Not reported	Thoracotomy
Brunelli, 2008, ²⁷ Italy	536	67.0 (9.0)	440 lobectomies 96 pneumonectomies	85.6±18.9%	Maximal/oxygen desaturation Climbed height	Cardiac: 50% Respiratory: 47% Other: 3%	Thoracotomy
Nikolić <i>et al.</i> , ²⁵ 2008, Croatia	101	61.1 (8.4)	55 lobectomies 14 bilobectomies 17 pneumonectomies 15 others	Complications (n=124) 1.63±0.44 L No complications (n=170) 1.75±0.49 L	Maximal/time Oxygen desaturation	Cardiac: 41% Respiratory: 32% Other: 27%	Thoracotomy
Toker <i>et al.</i> , ²⁶ 2008, Turkey	150	60.0 (10.6)	101 lobectomies 27 pneumonectomies 22 wedge resections	Complications (n=31) 2.29±0.69 No complications (n=119) 2.43±0.73	Submaximal/time Oxygen desaturation Heart rate change	Respiratory: 62% Cardiac: 38%	Thoracotomy
Salahuddin <i>et al.</i> , ²⁰ 2005, Pakistan	78	52.9 (16.2)	All major lung resection without details	<1 floor (n=19) 1.3±0.17 L ≥1 floor (n=59) 4.5±2.37 L	Maximal/climbed height	Respiratory: 67% Cardiac: 33%	Thoracotomy/VATS
Zurauskas <i>et al.</i> , ¹³ 2002, Lithuania	52	62.2 (4.6)	35 lobectomies 11 pneumonectomies 6 others	No data	Maximal/climbed height	Cardiac: 36% Respiratory: 59% Death: 5%	Thoracotomy
Girish <i>et al.</i> , ¹⁵ 2001, USA	83	66.3 (1.3)	31 lobectomies 6 wedge resections 3 pneumonectomies 3 thymectomies 39 others	Complications (n=21) 1.8±0.77 L No complications (n=62) 2.3±0.82 L	Maximal/climbed height	Respiratory: 67% Cardiac: 25% Death: 8%	Thoracotomy
Pate <i>et al.</i> , ²⁹ 1996, USA	11	63.6 (4.9)	5 lobectomies 2 segmentectomies 2 wedge resections 1 pneumonectomy 1 other	1.38±0.3 L	Maximal/climbed height	Respiratory: 71% Cardiac: 14.5% Death: 14.5%	Thoracotomy
Holden <i>et al.</i> , ¹⁶ 1992, USA	16	68.5 (8.7)	6 lobectomies 4 pneumonectomies 3 wedge resections 3 segmentectomies	Complications (n=5) 1.38±0.08 L No complications (n=62) 1.44±0.13 L	Maximal/climbed height	Respiratory: 53% Cardiac: 40% Death: 6%	Thoracotomy
Retrospective cohort studies							
Dong <i>et al.</i> , ¹⁹ 2017, China	171	65.0 (9.0)	27 wedge resection 122 lobectomies 22 pneumonectomies	Complications (n=124) 1.63±0.44 L No complications (n=170) 1.75±0.49 L	Submaximal/ Oxygen desaturation Heart rate change Climbed height	Cardiac: 59% Respiratory: 36% Death: 5%	Thoracotomy/VATS
Olsen <i>et al.</i> , ¹⁷ 1991, USA	54	60.6 (9.9)	25 lobectomies 21 wedge resections 4 pneumonectomies 4 others	2.41±0.62 L	Maximal/climbed height	Respiratory: 55% Cardiac: 34% Other: 9% Death: 2%	Thoracotomy

FEV1, forced expiratory volume in 1 s; VATS, video-assisted thoracic surgery.

Study characteristics

The study characteristics of the 13 included articles are summarised in [table 1](#). There were 11 prospective cohort studies and 2 retrospective cohort studies. Studies were published between 1991 and 2017 and included data from 2038 patients, with a mean age ranging from 52.7 years to 68.5 years. All were conducted in a hospital setting.

Two studies published by the same team included patients during overlapping periods.^{14 27} However, the first study¹⁴ reported climbed height as assessment criteria while the second²⁷ reported desaturation and height during the test. Only the first was included in the meta-analysis.

Exposure reporting

Desaturation

Four cohort studies investigated the association between oxygen desaturation and postoperative complications. Distance to climb was different between studies. In the first study, patients had to climb six floors and the saturation change was noted at this term with a mean desaturation of 6.15% (SD 5.86%, n=46) in patients with complications vs 4.06% (SD 3.94%, n=124) in patients without complications¹⁹ (p=0.008). In the second study, in which patients had to climb three floors,²⁶ the mean desaturation was 2.50% (SD 2.58%) in patients with complications vs 1.18% (SD 0.98%) in patients without complications

Figure 1 PRISMA diagram showing selection of studies for systematic review and meta-analysis. SCT, stair-climbing test.

($p=0.017$). In the third study, in which patients had to provide a maximal effort climb, a saturation $<90\%$ at the end of the test was observed in 26 (51.0%) patients with major complications or death and in 28 (56.0%) patients without complications or with minor complications²⁵ ($p=0.61$). In the fourth study, after a maximal effort, 27 (21%) patients with complications had a SpO₂ drop $\geq 4\%$ vs 48 (12%) patients without complications ($p=0.008$). In the same study, 6 (4.7%) patients with complications had a SpO₂ under 90% at the end of the test vs 21 (5.1%) patients without complications ($p=0.80$).²⁷ Definitions of desaturations and evaluation methods differed between studies; hence, we were not able to pool data because of this heterogeneity.

Time

Three cohort studies reported time as assessment criteria.^{25 26 28} For Nikolić *et al*,²⁵ the duration of maximal effort SCT was only predictive of postoperative complications in the lobectomy subgroup, with a mean test duration of 96.1 s (SD 14.5) in patients without complications or with minor complications and of 80.7 s (SD 14.2) in patients with complications. Ambrozin *et al*²⁸ reported, for six floors climbed, 46.6 s (SD 17.4) for patients with complications vs 35.8 s (SD 12.8) for patients without complications ($p=0.005$), while Toker *et al*²⁶ reported 21.67 s (SD 6.55) vs 19.65 s (SD 6.32) for 3 m ($p=0.16$).

Heart rate change

Only two studies reported this exposure. Dong *et al*¹⁹ reported, for five floors climbed, a mean heart rate change of 48.8 (SD 13.5) in patients with complications vs 54.1 (SD 14.5) in patients without complications ($p=0.03$), while Toker *et al*²⁶ reported,

for 3 m climbed, respectively, 14.85 (SD 3.77) and 14.93 (SD 3.55) ($p=0.90$).

Climbed height

Eight studies investigated the association between the height climbed and postoperative morbidity. The overall homogeneity allowed us to perform a meta-analysis. One study²⁹ was excluded from quantitative analysis ($n=11$) because it did not define a climb threshold, and RRs at the median could not be calculated. All patients included in the meta-analysis were instructed to climb the maximum number of floors until exhaustion. Studies in which patients did not perform the test under maximum conditions were excluded from the analysis.¹⁹ For the study of Olsen *et al*,¹⁷ two thresholds were presented (19 m and 10 m). The one closest to the mean of the other studies was selected (10 m).

Results of meta-analysis: relative risk

Figure 2 shows RR estimations of the random effects model at the threshold defined by the study. The overall pooled RR of complications was 2.34 (95% CI 1.59 to 3.43) with heterogeneity coefficient $I^2=53\%$ ($p=0.06$). The pooled RR of complications in the fixed-effects model (sensitivity analysis) was 2.31 (95% CI 1.79 to 2.96). A sensitivity analysis was performed by using fixed-effect models by excluding the two studies with the lowest quality score showing an RR estimated at 2.00 (95% CI 1.53 to 2.61), with a non-significant heterogeneity ($I^2=34\%$, $p=0.21$) (online supplementary figure S2).

Each study compared the risk of complications of patients exceeding a threshold to that of those not exceeding it. This threshold was different in each study, varying between 2.43 m³⁰

Figure 2 Forest plot of the relative risks of postoperative cardiopulmonary complications for a height of climbing below (vs above) the study-specific threshold.

and 12 m.¹⁴ We therefore analysed for each study the association between complications and height climbed at its own threshold (figure 2). A pooled threshold of height climbed, indicating a high risk of complications, was obtained by the unweighted mean of thresholds of all the studies included in the meta-analysis. In a sensitivity analysis, the two studies with poor methodological quality were excluded. The mean threshold including the studies with poor methodological quality^{13 30} was 8.11 m whereas the threshold excluding these studies was 9.91 m.

Predictive performances

The sensitivity, specificity, PPV and NPV of the SCT were estimated respectively at 43% (95% CI 21% to 68%), 88% (95% CI 82% to 92%), 62% (95% CI 42% to 78%) and 75% (95% CI 65% to 83%). Confidence regions of predictive values are shown in figure 3.

Risk of bias

The overall 'Risk of Bias' of each included study is presented in figure 4. Overall crude agreement on methodological quality scores between the reviewers was 84.6%. Five studies had a low risk of bias, four had a moderate risk of bias and four had a high risk of bias, mainly due to the variety of confounding factors.

We assessed the overall level of evidence of the univariate prognosis value of the SCT for postoperative complications with the adapted GRADE framework defined by Huguet *et al* for prognostic meta-analyses.³¹ The investigational phase was at 2 for all studies. There was no problem on: inconsistency, indirectness, publication bias. There was a downgrade for serious study limitations (high risk of bias in several studies) and an upgrade for moderate/large effect size. Overall, the level of evidence was high.

Figure 3 Sensitivity, specificity (panel A), PPV and NPV (panel B) of each study (grey crosses and 95% thin ovoid confidence regions) and of the meta-analysis (round and bold 95% ovoid confidence region) in Retisma's random effect model taking in account the correlation between sensitivity and specificity and between PPV and NPV. NPV, negative predictive value; PPV, positive predictive value.

Assessment of risk of bias

Selection bias: two studies do not include only thoracic surgery candidates, but also a minor proportion of abdominal surgery.^{15 30}

Information bias on exposition: the heterogeneity of SCT is not perfectly standardised. The types of stairs are different, and this can have an impact on the result. Furthermore, the reproducibility of the test is unknown.

Information bias on outcome: we observe a difference in follow-up complications with a lack of standardised definition of postoperative complications.

DISCUSSION

This is the first meta-analysis of observational studies on the association between the height climbed during SCT and postoperative morbidity. We found evidence of a 2.34-fold (95%

	Study participation	Study attrition	Prognostic factor measurement	Outcome measurement	Study confounding	Statistical analysis and reporting	Overall risk of bias
Dong 2017	Low	Low	Low	Moderate	Moderate	Low	Low
Ambrozio 2013	Moderate	Low	Low	Moderate	High	Low	Moderate
Amuda 2013	Low	Low	Low	Moderate	Moderate	Low	Low
Brunelli 2008	Low	Low	Low	Moderate	Moderate	Low	Low
Brunelli 2008	Low	Low	Low	Moderate	Moderate	Low	Low
Nikolic 2008	Moderate	Low	Moderate	Moderate	High	Low	High
Toker 2008	Low	Low	High	Moderate	Moderate	Low	Moderate
Salahuddin 2005	High	High	High	Moderate	High	High	High
Zurauskas 2002	Moderate	Moderate	Moderate	High	High	High	High
Girish 2001	Moderate	Low	Low	Moderate	Moderate	Moderate	Moderate
Pate 1996	High	Moderate	Low	Moderate	High	High	High
Holden 1992	Low	Low	Moderate	Moderate	High	Low	Moderate
Olsen 1991	Moderate	Low	Low	Moderate	Moderate	Low	Low

Figure 4 Risk of bias according to the QUIPS tool. Red circle: high risk of bias, orange circle: moderate risk of bias, green circle: low risk of bias. QUIPS, Quality in Prognosis Studies.

CI 1.59 to 3.43) increased risk of postoperative complications beyond the climb threshold of SCT.

This meta-analysis included two low-quality studies. However, their exclusion had little impact on estimations of effects. The high methodological heterogeneity of studies included in this systematic review made it impossible to pool any effect, but the climbed height. Even that analysis was limited by variable climbing thresholds. The climbing threshold being variable from study to study, the predictive value at the mean threshold (9.91 m) could not be computed on aggregated data. Individual data meta-analysis was not possible since we failed to acquire study databases despite our requests to the authors.

Exercise tests simulate surgical stress and detect potential defects in the lung, heart and/or muscular chain of the oxygen transport system.⁵ This encourages the widespread use of preoperative exercise in thoracic surgery units. The SCT involves a large muscle mass and it is a functional test, informing the clinicians about the daily capacities of patients. The number and height of steps in stair flights varied between centres. We tried to standardise the results by reporting the performance in metres. This study shows that the number of floors was not a reliable value, 12 m corresponding in some studies to six floors,¹⁴ while 10 m corresponded to 2.5 floors in others.¹⁷ Metric altitude standardisation allows therapists to adapt the risk threshold according to patients' consultation environment.

ERS/ESTS guidelines recommend that SCT should be used as a first-line functional screening test to select patients for safe surgery (height of ascent >22 m) (grade of recommendation: B).⁴ The 10 m threshold computed in our work is based on pooling studies that directly estimated the risk of complications, while the 22 m threshold is based on a single study³² that did not measure postoperative complications, but estimated it indirectly via VO₂max. A total of 98% of patients who were above the 22 m threshold had high VO₂max (>15 mL/kg/min) but only 23% of patients who were below the threshold had low VO₂max (<15 mL/kg/min). From the same study, a threshold of 14 m would provide a PPV of 56%, a sensitivity of 64% and a specificity of 87%.

The climbed height or climbing speed is correlated ($r=0.63-0.72$) with the VO₂peak reached during SCT.^{19 28} VO₂peak reflects a patient's aerobic activity and ability to respond to intense physical stress, as during major surgery. SCT yields greater values of VO₂peak than ergocycle.³³ This can be explained by the visual feedback and the motivation of the patient to reach the next floor before stopping. Only 10% of patients performing SCT stopped the test before reaching the next floor.³²

After reanalysis of raw data published by Bolliger *et al*, VO₂peak PPV, at threshold 15 mL/min/kg was 47% (23%–72%) and NPV was 87% (77%–94%). In our meta-analysis, the PPV of SCT was higher (62%), but NPV was lower (75%).³⁴

The SCT as the 6MWT can be used to predict postoperative outcomes.³⁵ The 6MWT is very easy to perform but requires a 30 m long corridor which is not always available. The SCT requires stairs that are high enough. To improve risk control, some teams prefer the use of ergocycle in a secured environment.³⁶

As SCT is a highly demanding cardiorespiratory test, there are associated risks.³⁷ It can identify patients presenting risks, but it is difficult to determine if the risk is cardiac, respiratory and/or neuromuscular. Regarding these results, we suggest using SCT more as a screening tool to identify patients who would benefit from CPET, rather than as a substitute for this major examination.

Currently, the use of CPET is based on European and American recommendations.^{4 5} These recommendations are relative to

resting respiratory parameters (FEV1/DLCO) at a time when the patient's evaluation must be dynamic. In addition, these recommendations, like the majority of the studies included in this meta-analysis, are relative to thoracotomy-operated patients, whereas at present, surgery by video or robotic-assisted thoracic surgery (VATS or RATS) is largely preferred. There are too few prospective studies including these techniques.

The strength of this meta-analysis is an extensive literature search, reporting based on PRISMA statement and standardised assessment of bias, whereas most articles report small case series limited to a single institution. As part of this analysis, patients treated with VATS were combined with those treated with thoracotomy.^{19 30} Compared with thoracotomy, VATS enables less postoperative pain, fewer complications and shorter hospitalisation.³⁸ Most of the reviewed studies did not include VATS resection. Modern thoracic surgery may allow more complex cases to be performed. All patients in the meta-analysed studies had major pulmonary resection except in the Salahuddin *et al*³⁰ and Girish *et al*¹⁵ studies, which included 65% and 50% of patients with thoracic surgery, respectively, and the remainder had upper abdominal surgery.

As with all meta-analyses, the quality of pooled results is dependent on the quality of the included studies. The included studies were small, with variable methodological quality. The high variability of the average height climbed suggests that the populations and/or test conditions were variable. The surgery extent (from wedge resection to pneumonectomy) was variable and subgroup analyses were impossible with the published data. Postoperative complications were defined in all studies as cardiorespiratory impairment resulting in morbidity or mortality. However, complications were not detailed in all studies. It would be helpful to know which complications were most frequently detected by the SCT. Some may be relatively easily managed such as atrial arrhythmia whereas others such as ventilation insufficiency may not. None used the Clavien-Dindo classification to rank complications.³⁹

Moreover, postoperative follow-up was not of the same duration (hospital stay of up to 90 days), even if according to Agostini's work, postoperative complications occur mostly during hospital stay.⁴⁰

Our meta-analysis completes the systematic review of Moran *et al* investigating the predictive capacity of field tests in abdominal surgery and calls for further evaluation of SCT in a context of minimally invasive surgery or combined with other low-technology tests.⁴¹

CONCLUSION

This meta-analysis and the systematic review highlight an association between height climbed at stair-climbing test and postoperative complications. SCT could be used as a screening tool in patients with lung cancer in order to select candidates for thoracic surgery who could benefit from preoperative CPET and prehabilitation. However, better designed prospective studies are necessary to assess the predictive value of low-technology tests and risk classification algorithms in a context of minimally invasive thoracic surgery.

Author affiliations

¹Department of General and Thoracic Surgery, CHU Rouen, Rouen, France

²Normandie University UNIROUEN, INSERM U1096, Rouen, France

³Biostatistics Unit, CHU Rouen, Rouen, Normandie, France

⁴ADIR Association, Bois Guillaume, France

⁵Normandie University, UNIROUEN, UPRES EA 3830, Rouen University Hospital, Haute Normandie Research and Biomedical Innovation, Rouen, France

⁶CETAPS EA 3832, Mont Saint Aignan, France

⁷ERFPS, CHU Rouen, Rouen, France

⁸Pulmonary & Respiratory Intensive Care Department, CHU Rouen, Rouen, Normandie, France

Twitter Fairuz Boujibar @boujibarf

Acknowledgements The authors are grateful to Nikki Sabourin-Gibbs, Rouen University Hospital, for her help in editing the manuscript.

Contributors Contributed to conception and design, acquisition, analysis and interpretation of data: FB, AG, JMB, TG, FEG, AC, TB. Contributed to drafting the work and critical revisions: FB, AG, JMB, FEG, TB. Contributed to final approval of the version submitted: FB, AG, JMB, TG, FEG, AC, TB

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available on reasonable request.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Tristan Bonnevie <http://orcid.org/0000-0003-0550-813X>

REFERENCES

- Howington JA, Blum MG, Chang AC, *et al.* Treatment of stage I and II non-small cell lung cancer: diagnosis and management of lung cancer, 3rd ED: American College of chest physicians evidence-based clinical practice guidelines. *Chest* 2013;143:e27–85–313.
- de Leyn P, Decker G. [Surgical treatment of non-small cell lung cancer]. *Rev Mal Respir* nov 2004;21:971–82.
- Brunelli A, Kim AW, Berger KI, *et al.* Physiologic evaluation of the patient with lung cancer being considered for resectional surgery: diagnosis and management of lung cancer, 3rd ED: American College of chest physicians evidence-based clinical practice guidelines. *Chest* 2013;143:e1665–90.
- Brunelli A, Charloux A, Bolliger CT, *et al.* The European respiratory Society and European Society of thoracic surgeons clinical guidelines for evaluating fitness for radical treatment (surgery and chemoradiotherapy) in patients with lung cancer. *Eur J Cardiothorac Surg* 2009;36:181–4.
- American Thoracic Society, American College of Chest Physicians. ATS/ACCP statement on cardiopulmonary exercise testing. *Am J Respir Crit Care Med* 2003;167:211–77.
- Cavalheri V, Granger C. Preoperative exercise training for patients with non-small cell lung cancer. *Cochrane Database Syst Rev* 2017;2017:CD012020.
- Charloux A, Brunelli A, Bolliger CT, *et al.* Lung function evaluation before surgery in lung cancer patients: how are recent advances put into practice? A survey among members of the European Society of thoracic surgeons (ESTs) and of the thoracic oncology section of the European respiratory Society (ERS). *Interact Cardiovasc Thorac Surg* 2009;9:925–31.
- Brunelli A, Pompili C, Salati M. Low-Technology exercise test in the preoperative evaluation of lung resection candidates. *Monaldi Arch Chest Dis* 2010;73:72–8.
- Agostini P, Cieslik H, Rathinam S, *et al.* Postoperative pulmonary complications following thoracic surgery: are there any modifiable risk factors? *Thorax* 2010;65:815–8.
- Van Nostrand D, Kjelsberg MO, Humphrey EW. Preresectional evaluation of risk from pneumonectomy. *Surg Gynecol Obstet* 1968;127:306–12.
- Moher D, Liberati A, Tetzlaff J, *et al.* Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Ann Intern Med* 2009;151:W64:264–9.
- Macaskill P, Gatsonis C, Deeks J, *et al.* *Cochrane Handbook for systematic reviews of diagnostic test accuracy*, 2010: 61.
- Zurauskas A, Tikuisis R, Miliauskas P. [Stair climbing test in prediction of postoperative complications after lung cancer surgery]. *Medicina* 2002;38 Suppl 2:37–9.
- Brunelli A, Refai M, Xiumé F, *et al.* Performance at symptom-limited stair-climbing test is associated with increased cardiopulmonary complications, mortality, and costs after major lung resection. *Ann Thorac Surg* 2008;86:240–8.
- Girish M, Trayner E, Dammann O, *et al.* Symptom-limited stair climbing as a predictor of postoperative cardiopulmonary complications after high-risk surgery. *Chest* 2001;120:1147–51.
- Holden DA, Rice TW, Stelmach K, *et al.* Exercise testing, 6-min walk, and stair climb in the evaluation of patients at high risk for pulmonary resection. *Chest* 1992;102:1774–9.
- Olsen GN, Bolton JW, Weiman DS, *et al.* Stair climbing as an exercise test to predict the postoperative complications of lung resection. two years' experience. *Chest* 1991;99:587–90.
- Arruda KA, Cataneo DC, Cataneo AJM. Surgical risk tests related to cardiopulmonary postoperative complications: comparison between upper abdominal and thoracic surgery. *Acta Cir Bras* 2013;28:458–66.
- Dong J, Mao Y, Li J, *et al.* Stair-Climbing test predicts postoperative cardiopulmonary complications and hospital stay in patients with non-small cell lung cancer. *Med Sci Monit* 2017;23:1436–41.
- Hayden JA, van der Windt DA, Cartwright JL, *et al.* Assessing bias in studies of prognostic factors. *Ann Intern Med* 2013;158:280–6.
- Cochran WG. The combination of estimates from different experiments. *Biometrics* 1954;10:101–29.
- Reitsma JB, Glas AS, Rutjes AWS, *et al.* Bivariate analysis of sensitivity and specificity produces informative summary measures in diagnostic reviews. *J Clin Epidemiol* 2005;58:982–90.
- Harbord RM, Deeks JJ, Egger M, *et al.* A unification of models for meta-analysis of diagnostic accuracy studies. *Biostatistics* 2007;8:239–51.
- Leeflang MMG, Deeks JJ, Rutjes AWS, *et al.* Bivariate meta-analysis of predictive values of diagnostic tests can be an alternative to bivariate meta-analysis of sensitivity and specificity. *J Clin Epidemiol* 2012;65:1088–97.
- Nikolić I, Majerić-Kogler V, Plavec D, *et al.* Stairs climbing test with pulse oximetry as predictor of early postoperative complications in functionally impaired patients with lung cancer and elective lung surgery: prospective trial of consecutive series of patients. *Croat Med J* 2008;49:50–7.
- Toker A, Ziyade S, Bayrak Y, *et al.* Prediction of cardiopulmonary morbidity after resection for lung cancer: stair climbing test complications after lung cancer surgery. *Thorac Cardiovasc Surg* 2007;55:253–6.
- Brunelli A, Refai M, Xiumé F, *et al.* Oxygen desaturation during maximal stair-climbing test and postoperative complications after major lung resections. *Eur J Cardiothorac Surg* 2008;33:77–82.
- Ambrozio ARP, Cataneo DC, Arruda KA, *et al.* Time in the stair-climbing test as a predictor of thoracotomy postoperative complications. *J Thorac Cardiovasc Surg* 2013;145:1093–7.
- Pate P, Tenholder MF, Griffin JP, *et al.* Preoperative assessment of the high-risk patient for lung resection. *Ann Thorac Surg* 1996;61:1494–500.
- Salahuddin N, Fatimi S, Salahuddin N, *et al.* Predicting postoperative cardio-pulmonary complications by a test of stair climbing. *J Coll Physicians Surg Pak* 2005;15:761–4.
- Huguet A, Hayden JA, Stinson J, *et al.* Judging the quality of evidence in reviews of prognostic factor research: adapting the grade framework. *Syst Rev* 2013;2:71.
- Brunelli A, Xiumé F, Refai M, *et al.* Peak oxygen consumption measured during the stair-climbing test in lung resection candidates. *Respiration* 2010;80:207–11.
- Pollock M, Roa J, Benditt J, *et al.* Estimation of ventilatory reserve by stair climbing. A study in patients with chronic airflow obstruction. *Chest* 1993;104:1378–83.
- Bolliger CT, Jordan P, Soler M, *et al.* Exercise capacity as a predictor of postoperative complications in lung resection candidates. *Am J Respir Crit Care Med* 1995;151:1472–80.
- Kubori Y, Matsuki R, Hotta A, *et al.* Comparison between stair-climbing test and six-minute walk test after lung resection using video-assisted thoracoscopic surgery lobectomy. *J Phys Ther Sci* 2017;29:902–4.
- Novoa NM, Esteban P, Rodriguez M, *et al.* Functional evaluation before lung resection: searching for a low technology test in a safer environment for the patient: a pilot study. *Eur J Cardiothorac Surg* 2017;51:856–60.
- Noonan V, Dean E. Submaximal exercise testing: clinical application and interpretation. *Phys Ther* 2000;80:782–807.
- Al-Ameri M, Bergman P, Franco-Cereceda A, *et al.* Video-Assisted thoracoscopic versus open thoracotomy lobectomy: a Swedish nationwide cohort study. *J Thorac Dis* 2018;10:3499–506.
- Dindo D, Demartines N, Clavien P-A. Classification of surgical complications: a new proposal with evaluation in a cohort of 6336 patients and results of a survey. *Ann Surg* 2004;240:205–13.
- Agostini P, Naidu B, Cieslik H, *et al.* Comparison of recognition tools for postoperative pulmonary complications following thoracotomy. *Physiotherapy* 2011;97:278–83.
- Moran J, Wilson F, Guinan E, *et al.* The preoperative use of field tests of exercise tolerance to predict postoperative outcome in intra-abdominal surgery: a systematic review. *J Clin Anesth*. 2016;35:446–55. 1 déc.