Air pollution and chronic bronchitis: the evidence firms up

Frank Kelly

Chronic bronchitis is defined epidemiologically as cough and sputum production for ≥ 3 months in each of at least two consecutive years.¹ It affects about a third of patients with chronic obstructive pulmonary disease but also occurs in individuals with normal lung function, with prevalence estimates varying widely.^{2–7} While it is not, by itself, associated with a substantially higher risk of death,⁸ it is associated with exacerbations of disease, incapacity and poor quality of life. The identification of chronic bronchitis was standardised in the first instance by the British Medical Research Council questionnaire,⁹ and subsequently, the questions were introduced into other respiratory symptom questionnaires.

Ambient air pollution is increasingly being linked to a range of human conditions, and in the case of chronic bronchitis, which affects over two million individuals in the UK,¹⁰ there is some evidence of an association between the incidence or prevalence of chronic bronchitis and long-term exposure to air pollution.^{11–13} Early reports, however, relate to a time when the bulk of outdoor air pollution came from coal burning when pollutant levels were considerably higher than they are today. Of interest, a real-world study by Holland and Reid showed higher rates of cough and phlegm and lower lung function in London postal workers where pollution levels were much higher than in the three county towns where comparative postal workers were recruited.¹⁴

In 2016, the Committee on the Medical Effects of air Pollutants identified 36 studies that examined a possible link between air pollution (primarily particulate matter (PM) pollution) and chronic bronchitis.¹⁵ Some of the longitudinal studies in Europe and USA that were included in this report demonstrated changes in symptoms following a reduction in the concentration of air

pollution.^{16–18} However, the overall body of evidence of associations between chronic bronchitis and long-term exposure to air pollution was inconsistent, and more detailed and well-controlled epidemiological studies were recommended.¹⁵ Rising to this challenge, Doiron and colleagues in this issue of *Thorax* report findings from the largest analysis to date to examine cross-sectional and longitudinal associations between ambient air pollution and chronic bronchitis.¹⁹

In their carefully controlled study, using the Dutch Lifetimes cohort data on 132 595 (baseline) and 65 009 (second assessment) participants, they assessed possible associations of air pollution with prevalence and incidence of chronic bronchitis (winter cough and sputum almost daily for ≥ 3 months/year), chronic cough (winter cough almost daily for ≥ 3 months/year) and prevalence of cough and sputum symptoms. Logistic regression models were deployed to adjust for sex, age, educational attainment, body mass index, smoking status, pack-years smoking and environmental tobacco smoke at home. Whereas the majority of previous studies focused on exposure to particulate pollution alone, Doiron *et al* used land use regression model exposure estimates from the ELAPSE study for PM with a diameter of less than $2.5 \mu\text{m}$, and nitrogen dioxide (NO_2)²⁰ and black carbon (BC) from the ESCAPE study.²¹

In the north of the Netherlands, NO_2 and BC concentrations were found to be associated with increased odds of prevalent and incident chronic bronchitis. Of note, higher effect sizes were seen among women, never smokers and younger individuals supporting earlier findings for women in the European Community Respiratory Health Survey, where NO_2 was associated with the prevalence and new onset of chronic phlegm and chronic productive cough.²² This finding also aligns with Gundersen *et al*, who found a large excess of cough and phlegm associated with moderate and high levels of exposure to traffic among women, which in this study was particularly marked among smokers and not found among men.²³

Given this is the largest study to date examining the relationship between relatively low ambient air pollution exposures and chronic bronchitis, it contributes to an urgent need for research into the link between air pollution and this chronic condition. Furthermore, in relation to the impact on health, the 2005 report of the cost-benefit analyses for the Clean Air for Europe programme estimated (in monetary valuation) that chronic bronchitis was, after adult mortality, the next most important health outcome in relation to long-term exposure to air pollution.²⁴ The findings reported by Doiron and colleagues set the scene for further work to estimate economic costs of this disease attributable to ambient air pollution, which, given its prevalence, are likely to be sizeable.

Contributors FK, the sole author of the study, drafted the work or revised it critically for important intellectual content; provided the final approval of the version to be published; and agreed to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Funding This study was funded by National Institute for Health Research (grant number NIHR200880).

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Commissioned; externally peer reviewed.

OPEN ACCESS

Open access This is an open access article distributed in accordance with the Creative Commons Attribution 4.0 Unported (CC BY 4.0) license, which permits others to copy, redistribute, remix, transform and build upon this work for any purpose, provided the original work is properly cited, a link to the licence is given, and indication of whether changes were made. See: <https://creativecommons.org/licenses/by/4.0/>.

© Author(s) (or their employer(s)) 2021. Re-use permitted under CC BY. Published by BMJ.

To cite Kelly F. *Thorax* 2021;**76**:744–745.

Accepted 1 March 2021

Published Online First 23 March 2021

► <http://dx.doi.org/10.1136/thoraxjnl-2020-216142>

Thorax 2021;**76**:744–745.

doi:10.1136/thoraxjnl-2021-216883

ORCID iD

Frank Kelly <http://orcid.org/0000-0003-2558-8392>

Environmental Research Group, Imperial College London Faculty of Medicine, London, UK

Correspondence to Dr Frank Kelly, Environmental Research Group, Imperial College London Faculty of Medicine, London W12 0BZ, UK; frank.kelly@imperial.ac.uk

REFERENCES

- 1 Definition and classification of chronic bronchitis for clinical and epidemiological purposes. A report to the medical Research Council by their Committee on the aetiology of chronic bronchitis. *Lancet* 1965;1:775–9.
- 2 Cerveri I, Accordini S, Verlato G, *et al.* Variations in the prevalence across countries of chronic bronchitis and smoking habits in young adults. *Eur Respir J* 2001;18:85–92.
- 3 Lu M, Yao W, Zhong N, *et al.* Chronic obstructive pulmonary disease in the absence of chronic bronchitis in China. *Respirology* 2010;15:1072–8.
- 4 de Oca MM, Halbert RJ, Lopez MV, *et al.* The chronic bronchitis phenotype in subjects with and without COPD: the PLATINO study. *Eur Respir J* 2012;40:28–36.
- 5 Jindal SK, Aggarwal AN, Gupta D, *et al.* Indian study on epidemiology of asthma, respiratory symptoms and chronic bronchitis in adults (INSEARCH). *Int J Tuberc Lung Dis* 2012;16:1270–7.
- 6 Ferré A, Fuhrman C, Zureik M, *et al.* Chronic bronchitis in the general population: influence of age, gender and socio-economic conditions. *Respir Med* 2012;106:467–71.
- 7 Huchon GJ, Vergnenègre A, Neukirch F, *et al.* Chronic bronchitis among French adults: high prevalence and underdiagnosis. *Eur Respir J* 2002;20:806–12.
- 8 Peto R, Speizer FE, Cochrane AL, *et al.* The relevance in adults of air-flow obstruction, but not of mucus hypersecretion, to mortality from chronic lung disease. results from 20 years of prospective observation. *Am Rev Respir Dis* 1983;128:491–500.
- 9 MRC (Medical Research Council on the Aetiology of Chronic Bronchitis). Standardized Questionnaires on respiratory symptoms. *BMJ* 1960;2:1665.
- 10 NHS inform. Available: www.nhsinform.scot/illnesses-and-conditions/lungs-and-airways/bronchitis
- 11 Hodgkin JE, Abbey DE, Euler GL, *et al.* Copd prevalence in nonsmokers in high and low photochemical air pollution areas. *Chest* 1984;86:830–8.
- 12 Euler GL, Abbey DE, Magie AR, *et al.* Chronic obstructive pulmonary disease symptom effects of long-term cumulative exposure to ambient levels of total suspended particulates and sulfur dioxide in California Seventh-Day Adventist residents. *Arch Environ Health* 1987;42:213–22.
- 13 Euler GL, Abbey DE, Hodgkin JE, *et al.* Chronic obstructive pulmonary disease symptom effects of long-term cumulative exposure to ambient levels of total oxidants and nitrogen dioxide in California Seventh-Day Adventist residents. *Arch Environ Health* 1988;43:279–85.
- 14 Holland WW, Reid DD. The urban factor in chronic bronchitis. *Lancet* 1965;1:445–8.
- 15 COMEAP. Long-Term exposure to air pollution and chronic bronchitis, 2016. Available: <https://www.gov.uk/government/publications/comeap-long-term-exposure-to-air-pollution-and-chronic-bronchitis#:~:text=The%20Committee%20found%20some%20evidence,to%20imply%20a%20causal%20relationship> [Accessed 15 Mar 2021].
- 16 Heinrich J, Topp R, Gehring U, *et al.* Traffic at residential address, respiratory health, and atopy in adults: the National German health survey 1998. *Environ Res* 2005;98:240–9.
- 17 Schindler C, Keidel D, Gerbase MW, *et al.* Improvements in PM10 exposure and reduced rates of respiratory symptoms in a cohort of Swiss adults (SAPALDIA). *Am J Respir Crit Care Med* 2009;179:579–87.
- 18 Schikowski T, Ranft U, Sugiri D, *et al.* Decline in air pollution and change in prevalence in respiratory symptoms and chronic obstructive pulmonary disease in elderly women. *Respir Res* 2010;11:113.
- 19 Doiron D, Bourbeau J, de Hoogh K, *et al.* Ambient air pollution exposure and chronic bronchitis in the lifelines cohort. *Thorax* 2021;76:772–9.
- 20 de Hoogh K, Chen J, Gulliver J, *et al.* Spatial PM2.5, NO2, O3 and bc models for Western Europe – evaluation of spatiotemporal stability. *Environ Int* 2018;120:81–92.
- 21 de Hoogh K, Gulliver J, Donkelaar Avan, *et al.* Development of West-European PM_{2.5} and NO₂ land use regression models incorporating satellite-derived and chemical transport modelling data. *Environ Res* 2016;151:1–10.
- 22 Sunyer J, Jarvis D, Gotschi T, *et al.* Chronic bronchitis and urban air pollution in an international study. *Occup Environ Med* 2006;63:836–43.
- 23 Gundersen H, Magerøy N, Moen BE, *et al.* Low traffic and respiratory symptoms among smoking females: the Hordaland health study. *Arch Environ Occup Health* 2012;67:189–98.
- 24 Hurley F, Hunt A, Cowie H. Methodology for the cost-benefit analysis for CAFE: volume 2: health impact assessment. *AEA Technology Environment* 2005.