

Supplementary Material

Association of the Dysfunctional Placentation Endotype of Prematurity with Bronchopulmonary Dysplasia: A Systematic Review, Meta-analysis and Meta-regression

1. Methods

1.1. Search strategy

Pubmed

1	<p>(((((outcome AND preterm)) OR (risk factor AND preterm))) AND (((((((case control) OR Case-Control Study) OR Case-Base Studies) OR cohort study) OR Observational study) OR RCT) OR Randomized controlled trial))) AND (((((((((((preterm infant [mesh]) OR premature infant [tiab]) OR premature infant) OR Neonatal Prematurity) OR Infants) OR Premature) OR Prematurity) OR Neonatal) OR Neonatal) OR before 37 completed weeks of gestation))) AND (((pre-eclampsia AND preterm))) OR (((EPH AND preterm AND preterm)) OR (hellp syndrome AND preterm)) OR (gestational hypertensive disorder AND preterm)) OR (pre-existing hypertension AND preterm)) OR (eclampsia AND preterm)) OR (toxemia AND preterm)) OR (pregnancy toxemia AND preterm)) OR (edema proteinuria hypertension gestosis AND preterm)) OR (hypertensive disorders of pregnancy AND preterm)) OR (maternal hypertension AND preterm)))</p>
2	<p>(bronchopulmonary dysplasia [MESH] OR bronchopulmonary dysplasia [tiab] OR BPD [tiab] OR chronic lung disease [tiab] OR CLD [tiab] OR pulmonary hypertension [MESH] OR pulmonary hypertension [tiab]) AND (preterm infant [tiab] OR Premature Infant [tiab] OR Premature Infants [tiab] OR preterm infants [tiab] OR neonatal prematurity [tiab] OR very low birth weight infant [tiab] OR Very-Low-Birth-Weight Infant [tiab] OR Very-Low-Birth-Weight Infants [tiab] OR very low birth weight infants [tiab] OR Extremely Low Birth Weight Infant [tiab] OR Extremely Low Birth Weight Infants [tiab] OR preterm infant [MESH] OR Premature Infant [MESH] OR Premature Infants [MESH] OR preterm infants [MESH] OR neonatal prematurity [MESH] OR very low birth weight infant [MESH] OR Very-Low-Birth-Weight Infant [MESH] OR Very-Low-Birth-Weight Infants [MESH] OR very low birth weight infants [MESH] OR Extremely Low Birth Weight Infant [MESH] OR Extremely Low Birth Weight Infants) AND (cohort [MESH] OR Incidence Studies [MESH] OR Incidence Study [MESH] OR concurrent studies [MESH] OR concurrent studies [MESH] OR cohort analysis [MESH] OR observational studies [MESH] OR observational study [MESH] OR case control study [MESH] OR case control studies [MESH] OR case control [MESH] OR cohort [tiab] OR Incidence Studies [tiab] OR Incidence Study [tiab] OR concurrent studies [tiab] OR concurrent studies [tiab] OR cohort analysis [tiab] OR observational studies [tiab] OR observational study [tiab] OR case control study [tiab] OR case control studies [tiab] OR case control [tiab])</p>

3	<p>(fetal growth restriction [tiab] OR intrauterine growth [tiab] OR absent end diastolic [tiab] OR reversed end diastolic [tiab] OR small for gestational age [tiab] OR SGA [tiab] OR small for date [tiab] OR IUGR [tiab] OR fetal growth restriction [MESH] OR intrauterine growth [MESH] OR absent end diastolic [MESH] OR reversed end diastolic [MESH] OR small for gestational age [MESH] OR SGA [MESH] OR small for date [MESH] OR IUGR [MESH])</p> <p>AND</p> <p>(preterm infant [tiab] OR Premature Infant [tiab] OR Premature Infants [tiab] OR preterm infants [tiab] OR neonatal prematurity [tiab] OR very low birth weight infant [tiab] OR Very-Low-Birth-Weight Infant [tiab] OR Very-Low-Birth-Weight Infants [tiab] OR very low birth weight infants [tiab] OR Extremely Low Birth Weight Infant [tiab] OR Extremely Low Birth Weight Infants [tiab] OR preterm infant [MESH] OR Premature Infant [MESH] OR Premature Infants [MESH] OR preterm infants [MESH] OR neonatal prematurity [MESH] OR very low birth weight infant [MESH] OR Very-Low-Birth-Weight Infant [MESH] OR Very-Low-Birth-Weight Infants [MESH] OR very low birth weight infants [MESH] OR Extremely Low Birth Weight Infant [MESH] OR Extremely Low Birth Weight Infants [MESH])</p> <p>AND</p> <p>(cohort [MESH] OR Incidence Studies [MESH] OR Incidence Study [MESH] OR concurrent studies [MESH] OR concurrent studies [MESH] OR cohort analysis [MESH] OR observational studies [MESH] OR observational study [MESH] OR case control study [MESH] OR case control studies [MESH] OR case control [MESH] OR cohort [tiab] OR Incidence Studies [tiab] OR Incidence Study [tiab] OR concurrent studies [tiab] OR concurrent studies [tiab] OR cohort analysis [tiab] OR observational studies [tiab] OR observational study [tiab] OR case control study [tiab] OR case control studies [tiab] OR case control [tiab])</p>
----------	--

EMBASE

1	(pre-eclampsia or preeclampsia or EPH or hellp syndrome or gestational hypertensive disorder or pre-existing hypertension or eclampsia or toxemia or pregnancy toxemia or edema proteinuria hypertension gestosis or maternal hypertension or pregnancy-induced hypertension).af.
2	(premature infant or Neonatal Prematurity or Infants, Premature or Prematurity or Neonatal or Preterm Infants).af.
3	(case control or Case-Control Study or Case-Base Studies or cohort study or RCT).af.
4	(risk factors or outcome or risk factor).af.
5	1 and 2 and 3 and 4
6	('chronic lung disease'/exp OR 'chronic lung disease') AND ('intrauterine growth retardation'/exp OR 'intrauterine growth retardation')
7	('chronic lung disease'/exp OR 'chronic lung disease' OR 'pulmonary hypertension') AND ('prematurity'/exp OR 'prematurity') AND ('cohort analysis'/exp OR 'cohort analysis' OR 'case control study'/exp OR 'case control study')
8	('intrauterine growth retardation'/exp OR 'intrauterine growth retardation') AND ('prematurity'/exp OR 'prematurity') AND ('cohort analysis'/exp OR 'cohort analysis' OR 'case control study'/exp OR 'case control study')

Web of Science

1	TOPIC: (pre-eclampsia or preeclampsia or EPH or hellp syndrome or gestational hypertensive disorder or hypertensive disorders of pregnancy or pre-existing hypertension or eclampsia or toxemia or pregnancy toxemia or edema proteinuria hypertension gestosis or maternal hypertension or pregnancy-induced hypertension) AND TOPIC: (premature infant or Neonatal Prematurity or Infants, Premature or Prematurity or Neonatal or Preterm Infants) AND TOPIC: (case control or Case-Control Study or Case-Base Studies or cohort study or observational) AND TOPIC: (risk factors or outcome or risk factor)
2	((bronchopulmonary dysplasia OR BPD OR chronic lung disease) AND (“preterm infant” OR “Premature Infant” OR “Premature Infants” OR “preterm infants” OR “neonatal prematurity” OR “very low birth weight” OR “Extremely Low Birth Weight” OR Extremely Low Birth Weight Infants) AND (cohort OR “Incidence Studies” OR “Incidence Study” OR “concurrent studies” OR “concurrent studies” OR “cohort analysis” OR “observational studies” OR “observational study” OR case control))

3	((“fetal growth restriction” OR “intrauterine growth retardation” or “intrauterine growth restriction” OR “small for gestational age”) AND (“preterm infant” OR “Premature Infant” OR “Premature Infants” OR “preterm infants” OR “neonatal prematurity” OR “very low birth weight” OR “Extremely Low Birth Weight” OR Extremely Low Birth Weight Infants) AND (cohort OR Incidence Studies OR Incidence Study OR concurrent studies OR concurrent studies OR cohort analysis OR observational studies OR observational study OR case control study OR case control studies OR case control))
----------	---

No language limits were set. Narrative reviews, systematic reviews, case reports, letters, editorials, and commentaries were excluded, but read to identify potential additional studies. Additional strategies to identify studies included manual review of reference lists from key articles that fulfilled our eligibility criteria, use of “related articles” feature in PubMed, and use of the “cited by” tool in Web of Science and Google scholar. Two reviewers independently screened the results of the searches, and included studies according to the inclusion criteria using EndNote (RRID:SCR_014001), using the methodology described by Bramer et al.¹

1.2. Supplementary information on methods

Study selection

Studies were included if they examined preterm (gestational age, GA <37 weeks) or very low birth weight (<1500g) infants and reported primary data that could be used to measure the association between exposure to hypertensive disorders of pregnancy (HDP) or small for GA (SGA)/intrauterine growth restriction (IUGR) and the development of BPD. Therefore, we selected cohort or case-control studies in which the exposure (HDP or SGA/IUGR) was the independent variable and the outcome (BPD) the dependent variable as well as studies in which the outcome was the independent variable and the exposure the dependent variable. Studies that exclusively included late preterm infants (GA \geq 34 weeks) or that combined preterm and term infants were excluded. The absence of a clear definition of BPD was also an exclusion criterion. Due to the high number of included studies, no additional efforts were made to clarify the definitions or other data with the authors. Abstracts and unpublished studies were also excluded. To identify relevant studies, two reviewers (M.P., M. A-F) independently screened the results of the searches and applied inclusion criteria using a structured form. Discrepancies were resolved through discussion or consultation with two other reviewers (E. V-M, E.V.).

Data extraction

Data extracted from each study included citation information, language of publication, location where research was conducted, time period of the study, study objectives, study design, inclusion/exclusion criteria, definition criteria for HDP, IUGR, SGA, BPD, and BPD-PH, patient characteristics, and results (including raw numbers or summary statistics when raw numbers were not available). Outcomes considered in meta-analysis were: 1) BPD28, defined as oxygen requirement on postnatal day 28; 2) BPD36, defined as oxygen requirement at the postmenstrual age (PMA) of 36 weeks; 3) BPD36 or death; 4) Severe BPD, defined as need for \geq 30% oxygen and/or positive pressure at 36 weeks PMA; 5) BPD-associated PH, defined by any echocardiographic criteria as long as the evaluation was performed at a postnatal age >4 weeks. Any definition of HDP or SGA/IUGR was accepted but we performed sub-group analysis based on the different definitions. When a study used more than one definition criteria for growth restriction, definitions based on assessment of fetal growth prevailed over definitions based on BW. When a study used different BW threshold percentiles to define SGA, data from the lowest percentile were included. When a study did not specify the threshold percentile used, it was grouped together with the studies that used the 10th percentile.

Statistical analysis


Studies were combined and analyzed using COMPREHENSIVE META-ANALYSIS V3.0 software (Biostat Inc., Englewood, NJ, USA). For dichotomous outcomes, the odds ratio (OR) with 95% confidence interval (CI) was calculated from the data provided in the studies. Reported OR were included when studies reported them and did not include the numerical data for its calculation. For continuous outcomes (example: gestational age), the mean difference (MD) with 95% CI was calculated. When studies reported continuous variables as median and range or interquartile range, we estimated the mean and standard deviation using the method of Wan et al. and the calculator they provided.²

Due to anticipated heterogeneity, summary statistics were calculated with a random-effects model. This model accounts for variability between studies as well as within studies. Subgroup analyses were conducted according to the mixed-effects model.³ In this model, a random-effects model is used to combine studies within each subgroup, and a fixed-effect model is used to combine subgroups and yield the overall effect. The study-to-study variance (tau-squared) is not assumed to be the same for all subgroups. This value is computed within subgroups and not pooled across subgroups. Statistical heterogeneity was assessed by Cochran's Q statistic and by the I^2 statistic, which is derived from Q and describes the proportion of total variation that is due to heterogeneity beyond chance.⁴ The I^2 statistic was interpreted as follows: low heterogeneity ($25\% \leq I^2 < 50\%$), moderate heterogeneity ($50\% \leq I^2 < 75\%$), and high heterogeneity ($I^2 \geq 75\%$). We used the Egger's regression test and funnel plots to assess publication bias. A probability value of less than 0.05 (0.10 for heterogeneity) was considered statistically significant.


Potential sources of heterogeneity were assessed through subgroup analysis and/or random effects (method of moments) univariate meta-regression analysis.⁵ For continuous covariates (examples: mean gestational age of the cohort, difference in mean gestational age between infants exposed and unexposed to HDP) we used meta-regression analyses to test whether there was a significant relationship between the covariate and effect size, as indicated by a Z-value and an associated p-value. Meta-regression coefficient indicates the change in the log of the OR of the association between BPD and the corresponding exposure for a unit change in the predictor covariate. Subgroups were compared using meta-regression for categorical covariates. For both categorical and continuous covariates, the R^2 analog, defined as the total between-study variance explained by the moderator, was calculated based on the meta-regression matrix.⁵ Covariates defined a priori were: 1) Continuous: mean or median GA of the entire cohort, study time (median year of cohort inclusion), differences between exposed and unexposed infants on GA, sex, rate of use of antenatal corticosteroids, and rate of respiratory distress syndrome (RDS); 2) Categorical: HDP type, SGA/IUGR type, study included only infants with GA<28 weeks (yes/no), DM of GA between exposed and unexposed infants statistically significant ($P < 0.05$, yes/no), and geographical location (continent) of the study. The clinical covariates were selected based on their relevance on the pathogenesis of BPD. On the basis of the recommendation of an anonymous reviewer, we added three categorical covariates related to the design of the studies: Prospective vs. Retrospective; Cohort vs. Case-control; and Exposure vs. Outcome as independent variable.

2. Results

2.1. Supplementary Figures




Supplementary Figure 1. Flow diagram of the systematic search.




Supplementary Figure 2. Meta-analysis of hypertensive disorders of pregnancy (HDP) and bronchopulmonary dysplasia, defined as oxygen requirement on postnatal day 28 (BPD28).

CI: confidence interval; PE: preeclampsia.




Supplementary Figure 3. Meta-analysis of small for gestational age/intrauterine growth restriction and bronchopulmonary dysplasia, defined as oxygen requirement on postnatal day 28 (BPD28).

BW: birth weight; CI: confidence interval; IUGR: intrauterine growth restriction (defined on the basis of fetal growth assessment); P3: 3rd percentile; P10: 10th percentile.




Supplementary Figure 4. Meta-analysis of hypertensive disorders of pregnancy (HDP) and bronchopulmonary dysplasia, defined as oxygen requirement at the postmenstrual age of 36 weeks (BPD36).

CI: confidence interval; PE: preeclampsia.




Supplementary Figure 5. Meta-analysis of the association between birth weight (BW) below the 10th percentile and bronchopulmonary dysplasia, defined as oxygen requirement at the postmenstrual age of 36 weeks (BPD36).




Supplementary Figure 6. Meta-analysis of the association between small for gestational age/intrauterine growth restriction and bronchopulmonary dysplasia, defined as oxygen requirement at the postmenstrual age of 36 weeks (BPD36).

BW: birth weight; CI: confidence interval; IUGR: intrauterine growth restriction (defined on the basis of fetal growth assessment); P5: 5th percentile; P10: 10th percentile.




Supplementary Figure 7. Meta-analysis of hypertensive disorders of pregnancy and severe bronchopulmonary dysplasia.

CI: confidence interval.




Supplementary Figure 8. Meta-analysis of small for gestational age/intrauterine growth restriction and severe bronchopulmonary dysplasia.

BW: birth weight; CI: confidence interval; IUGR: intrauterine growth restriction (defined on the basis of fetal growth assessment); P3: 3rd percentile; P10: 10th percentile.




Supplementary Figure 9. Meta-analysis of hypertensive disorders of pregnancy and the combined outcome bronchopulmonary dysplasia or death.




Supplementary Figure 10. Meta-analysis of small for gestational age/intrauterine growth restriction and the combined outcome bronchopulmonary dysplasia or death.

BW: birth weight; CI: confidence interval; IUGR: intrauterine growth restriction (defined on the basis of fetal growth assessment); P3: 3rd percentile; P5: 5th percentile; P10: 10th percentile.




Supplementary Figure 11. Meta-analysis of hypertensive disorders of pregnancy (HDP) and bronchopulmonary dysplasia-associated pulmonary hypertension.

CI: confidence interval.



Supplementary Figure 12. Meta-analysis of small for gestational age/intrauterine growth restriction and bronchopulmonary dysplasia-associated pulmonary hypertension.

BW: birth weight; CI: confidence interval; IUGR: intrauterine growth restriction (defined on the basis of fetal growth assessment); P3: 3rd percentile; P10: 10th percentile.



Supplementary Figure 13. Funnel plot for publication bias analysis for the studies included in the different meta-analyses.

A: Meta-analysis of hypertensive disorders of pregnancy (HDP) and BPD28. B: Meta-analysis of small for gestational age (SGA)/intrauterine growth restriction (IUGR) and BPD28. C: Meta-analysis of HDP and BPD36. D: Meta-analysis of SGA/IUGR and BPD36. E: Meta-analysis of SGA/IUGR and severe BPD. F: Meta-analysis of SGA/IUGR and BPD or death. G: Meta-analysis of HDP and BPD-associated pulmonary hypertension. H: Meta-analysis of SGA/IUGR and BPD-associated pulmonary hypertension.

Supplementary Table 1. Characteristics of the included studies

First author, year	Country	Design	Prospective?	Total infants	Centers	GA of cohort (weeks)	Independent variable	Exposures	Outcomes (% of incidence in total group)	NOS Selection	NOS Comparability	NOS Outcome/Exposure	NOS Total
Abramovici, 1999 ⁶	USA	cohort	no	2524	1	<36	Exposure	PE-HELLP	BPD28 (64)	3	1	3	7
Altman, 2011 ⁷	Sweden	Cohort	yes	6674	Network	33.1	Exposure	BW<P3	BPD36 (1)	4	1	3	8
Ameenudeen, 2007 ⁸	Malaysia	Cohort	yes	244	1	29.7	Outcome	Any HDP BW<P10	BPD36 (15)	4	2	3	9
Amin, 1997 ⁹	Canada	Ca-Co	no	186	1	28.3	Exposure	BW<P3	BPD28 (54)	4	1	3	8
An, 2010 ¹⁰	Korea	Cohort	no	116	1	26.3	Outcome	BW<P10	BPD-PH (25)	4	1	3	8
Aswani, 2016 ¹¹	USA	Cohort	No	230	1	25.9	Outcome	BW<P10	BPD-PH (8)	4	1	3	8
Aucott, 2004 ¹²	USA	Ca-Co	no	95	1	28.4	Exposure	BW<P5	BPD36 (21)	4	1	3	8
Baer, 2016 ¹³	USA	Cohort	no	78647	Network	33.0	Exposure	BW<P10	BPD36 (2)	4	1	3	8
Baker, 2012 ¹⁴	USA	cohort	yes	62	1	31.6	Outcome	PE BW<P10	BPD36 (21)	4	1	3	8
Balena, 2017 ¹⁵	USA	cohort	yes	113	1	27.9	Outcome	Any HDP	BPD/Death (36)	4	2	3	9
Bardin, 1997 ¹⁶	Canada	Cohort	yes	115	1	25.3	Exposure	BW<P3	BPD28 (86) BPD36 (37)	3	1	3	7
Bhat, 2012 ¹⁷	USA	Cohort	yes	145	1	26	Outcome	BW<P10	BPD-PH (18)	4	2	3	9
Blanca, 2018 ¹⁸	The Netherlands	Cohort	yes	69	1	25.6	Outcome	PE	BPD-PH (12)	3	1	3	7
Bose, 2009 ¹⁹	USA	Cohort	yes	1241	14	<28	Outcome	PE BW<P3	BPD36 (52)	4	2	3	9
Bose, 2011 ²⁰	USA	Cohort	yes	932	14	<28	Outcome	BW<P3	BPD28 (52) sBPD (11)	4	2	3	9
Bossung, 2020 ²¹	Germany	Cohort	yes	16035	62	28.2	Exposure	PE-HELLP	BPD36 (15)	4	2	3	9
Botet, 2012 ²²	Spain	Cohort	no	415	1	27.0	Outcome	BW<P10	BPD/Death (28)	4	2	3	9
Bozzetti, 2013 ²³	Italy	Cohort	yes	310	1	29.7	Exposure	BW<P10	BPD28 (9)	4	2	3	9

First author, year	Country	Design	Prospective?	Total infants	Centers	GA of cohort (weeks)	Independent variable	Exposures	Outcomes (% of incidence in total group)	NOS Selection	NOS Comparability	NOS Outcome/Exposure	NOS Total
Brener, 2017 ²⁴	Argentina	Cohort	no	203	1	29.4	Outcome	BW<P3	BPD36 (22)	4	2	3	9
Brumbaugh, 2018 ²⁵	USA	Cohort	no	151	1	26.6	Outcome	BW<P10	BPD36 (83) sBPD (55)	4	2	3	9
Brumbaugh, 2020 ²⁶	USA	Cohort	yes	2310	Network	24.9	Outcome	BW<P10	BPD36 (48) sBPD (5)	4	2	3	9
Bruno, 2015 ²⁷	USA	Cohort	no	303	1	26.6	Outcome	PE BW<P10	BPD-PH (12)	4	2	3	9
Burns, 1997 ²⁸	Australia	Ca-Co	no	117	5	28.2	Outcome	BW<P10	BPD28 (55)	4	2	3	9
Bursal, 2017 ²⁹	Turkey	cohort	no	284	1	31.2	Exposure	PE	BPD28 (10)	2	1	3	7
Candel, 2016 ³⁰	Spain	Ca-Co	yes	129	1	33.8	Exposure	IUGR	BPD36 (13)	4	1	3	8
Cartón, 2016 ³¹	Spain	Cohort	yes	84	1	27.0	Outcome	Any HDP BW<P10	BPD-PH (26)	4	2	3	9
Cederqvist, 2003 ³²	Finland	cohort	yes	32	1	27.8	Outcome	PE	BPD36 (56)	4	1	3	8
Çetinkaya, 2010 ³³	Turkey	Ca-Co	no	84	1	31.3	Exposure	PE	BPD36 (23)	2	1	3	6
Charafeddine, 1999 ³⁴	USA	Cohort	yes	123	1	27.8	Outcome	BW<P10	BPD28 (72)	4	2	3	9
Check, 2013 ³⁵	USA	Cohort	no	138	1	26.1	Outcome	PE BW<P10	BPD-PH (28)	3	1	3	7
Chen, 2020 ³⁶	USA	Cohort	yes	188	1	26.7	Outcome	BW<P10	BPD-PH (32)	4	2	3	9
Chen, 2016 ³⁷	Switzerland	Cohort	yes	8899	Network	29.2	Outcome	BW<P10	BPD36 (10)	4	2	3	9
Cheng, 2004 ³⁸	Taiwan	Ca-Co	no	89	1	28.3	Exposure	PE	BPD28 (36)	4	1	2	7
Choi, 2015 ³⁹	Korea	Cohort	no	194	1	26.5	Outcome	BW<P10	BPD-PH (26)	4	1	3	8
Choi, 2019 ⁴⁰	Korea	Cohort	no	81	1	25.7	Outcome	PE	BPD-PH (25)	4	1	3	8
Chye, 1995 ⁴¹	Australia	Ca-Co	yes	156	1	28.4	Outcome	BW<P10	BPD28 (50)	4	1	3	8
Claas, 2011 ⁴²	The Netherlands	Cohort	no	101	1	28.0	Exposure	BW<P3	BPD36 (56)	4	1	3	8

First author, year	Country	Design	Prospective?	Total infants	Centers	GA of cohort (weeks)	Independent variable	Exposures	Outcomes (% of incidence in total group)	NOS Selection	NOS Comparability	NOS Outcome/Exposure	NOS Total
Cunha, 2005 ⁴³	Brazil	cohort	yes	86	1	28.9	Outcome	Any HDP	BPD28 (52)	4	1	3	8
Da Silva, 2018 ⁴⁴	Brazil	Cohort	yes	67	3	29.1	Outcome	Any HDP	BPD36 (33)	4	2	3	9
Dasgupta, 2018 ⁴⁵	USA	Cohort	yes	52	1	26.6	Outcome	Any HDP	BPD36 (69) BPD-PH (15)	4	2	3	9
De Jesus, 2013 ⁴⁶	USA	Cohort	no	2971	Network	25.0	Exposure	BW<P10	BPD36 (36)	4	2	3	9
De Souza, 2015 ⁴⁷	Brazil	Ca-Co	yes	60	1	30.0	Exposure	PE	BPD36 (10)	4	2	3	9
Demirel, 2009 ⁴⁸	Turkey	Ca-Co	yes	106	1	30.4	Outcome	PE-HELLP	BPD28 (53)	4	2	3	9
DeVries, 2017 ⁴⁹	USA	Cohort	no	577	1	26.6	Outcome	BW<P10	BPD-PH (3)	4	2	2	8
Dogan, 2020 ⁵⁰	Turkey	Cohort	no	78	1	28.6	Outcome	PE	BPD36 (41)	4	1	3	8
Duan, 2016 ⁵¹	China	Cohort	yes	243	1	30.0	Outcome	Any HDP	BPD28 (29)	4	2	3	9
Duan, 2016b ⁵²	China	Cohort	yes	147	1	29.5	Outcome	Any HDP BW<P10	BPD36 (41)	4	2	3	9
Durrmeyer, 2012 ⁵³	France	Cohort	yes	265	1	26.0	Outcome	BW<P3	BPD36 (15) BPD/Death (25)	4	1	3	8
El Ayoub, 2016 ⁵⁴	Europe	Cohort	yes	4585	Network	28.0	Exposure	IUGR	BPD36 (14)	4	2	3	9
Enginner, 2010 ⁵⁵	UK	Ca-Co	no	121	1	<36	Exposure	IUGR	BPD28 (29)	4	1	3	8
Eriksson, 2015 ⁵⁶	Sweden	Ca-Co	no	2255	Network	30.0	Outcome	Any HDP BW<P10	BPD36 (24)	4	2	3	9
Estevez, 2017 ⁵⁷	Spain	Cohort	no	110	1	29.2	Exposure	BW<P10	BPD36 (5)	4	1	3	8
EXPRESS group, 2010 ⁵⁸	Sweden	Cohort	yes	497	Network	25.0	Outcome	PE BW<P3	sBPD (25)	4	2	3	9
Falciglia, 2003 ⁵⁹	USA	Cohort	yes	46	1	29.5	Exposure	PE	BPD36 (50)	3	2	3	8
Fernandez-Rodriguez, 2021 ⁶⁰	Spain	Cohort	no	95	1	31.4	Exposure	IUGR	BPD36 (9)	3	2	3	8
Fujioka, 2014 ⁶¹	Japan	Cohort	no	97	1	28.0	Outcome	Any HDP	BPD36 (57)	4	1	3	8

First author, year	Country	Design	Prospective?	Total infants	Centers	GA of cohort (weeks)	Independent variable	Exposures	Outcomes (% of incidence in total group)	NOS Selection	NOS Comparability	NOS Outcome/Exposure	NOS Total
								BW<P10					
Gagliardi, 2009 ⁶²	Italy	Cohort	yes	1209	14	28.7	Outcome	BW<P10	BPD36 (16)	4	2	3	9
Gagliardi, 2013 ⁶³	Italy	Cohort	yes	2085	Network	28.7	Exposure	Any HDP	BPD36 (13)	4	2	3	9
Gagliardi, 2014 ⁶⁴	Italy	Cohort	no	3606	82	27.2	Exposure	Any HDP	BPD36 (25)	4	2	3	9
Garite, 2004 ⁶⁵	USA	Cohort	no	24249	124	29.9	Exposure	IUGR	BPD28 (18)	3	2	3	8
Garofoli, 2014 ⁶⁶	Italy	Cohort	no	76	1	31.7	Exposure	BW<P10	BPD36 (9)	4	1	3	8
Gemmell, 2016 ⁶⁷	International	Cohort	no	27846	Network	26.5	Exposure	PE	BPD36 (35)	4	2	3	9
Giapros, 2012 ⁶⁸	Greece	Cohort	no	168	1	28.5	Exposure	BW<P10	BPD36 (35)	4	2	3	9
Gortner, 1999 ⁶⁹	Germany	Cohort	yes	317	6	29.6	Exposure	BW<P10	BPD36 (6)	4	1	3	8
Gortner, 2003 ⁷⁰	Germany	Ca-Co	yes	148	1	34.0	Exposure	BW<P10	BPD28 (6)	4	0	3	7
Gray, 1997 ⁷¹	Australia	Cohort	yes	189	1	27.3	Exposure	PE	BPD28 (49)	4	2	3	9
Greenough, 2005 ⁷²	UK	Ca-Co	yes	34	1	28.5	Outcome	BW<P10	BPD28 (50)	4	2	3	9
Grisaru-G, 2012 ⁷³	Israel	Cohort	yes	9756	Network	28.5	Exposure	BW<P3	BPD36 (15)	4	2	3	9
Guimaraes, 2010 ⁷⁷	Portugal	Cohort	yes	256	5	28.0	Outcome	BW<P10	BPD36 (18)	4	1	3	8
Guo, 2015 ⁷⁵	Taiwan	Cohort	no	75	1	27.6	Outcome	Any HDP BW<P10	sBPD (63)	4	1	3	8
Gursoy, 2015 ⁷⁶	Turkey	Cohort	no	652	1	29.4	Outcome	PE BW<P10	BPD28 (23)	4	2	3	9
Hadchouel, 2020 ⁷⁷	France	Ca-Co	no	19	3	26.7	Outcome	BW<P10	sBPD (58)	3	1	3	7
Hansen, 2010 ⁷⁸	UK	Cohort	yes	107	1	29.0	Outcome	PE	BPD36 (25)	4	2	3	9
Hartung, 2005 ⁷⁹	Germany	Ca-Co	no	88	1	31.0	Exposure	IUGR	BPD36 (9)	4	1	3	8
Hentges, 2015 ⁸⁰	Brazil	Cohort	yes	88	1	30.1	Exposure	PE	BPD28 (26)	4	2	3	9
Hernandez, 2004 ⁸¹	Mexico	Ca-Co	yes	44	1	31.1	Outcome	Any HDP	BPD28 (50)	4	1	3	8

First author, year	Country	Design	Prospective?	Total infants	Centers	GA of cohort (weeks)	Independent variable	Exposures	Outcomes (% of incidence in total group)	NOS Selection	NOS Comparability	NOS Outcome/Exposure	NOS Total
Hiett, 2001 ⁸²	USA	Ca-Co	no	116	2	27.1	Exposure	PE	BPD28 (60)	4	2	3	9
Hikino, 2012 ⁸³	Japan	Cohort	yes	26	1	29.0	Outcome	BW<P10	BPD36 (50)	4	1	3	8
Huusko, 2015 ⁸⁴	Finland, Canada en Hungary	Ca-Co	yes	772	8	27.8	Outcome	BW<P3	BPD36 (28)	4	2	3	9
Imamura, 2017 ⁸⁵	Japan	Cohort	no	169	1	26.0	Outcome	BW<P10	sBPD (40)	4	2	3	9
Iwatani, 2013 ⁸⁶	Japan	Cohort	no	51	1	25.7	Outcome	Any HDP	BPD36 (51)	4	2	3	9
Jakuskiene, 2011 ⁸⁷	Lithuania	Cohort	yes	238	1	27.4	Outcome	BW<P10	BPD36 (6)	4	1	3	8
Jensen, 2019 ⁸⁸	USA	Cohort	no	6708	Network	28.6	Exposure	BW<P10	BPD36 (29) BPD/Death (35)	4	1	3	8
Jeon, 2020 ⁸⁹	Korea	Cohort	yes	521	1	27.4	Outcome	Any HDP	BPD36 (48)	4	1	3	8
Jo, 2015 ⁹⁰	Korea	Cohort	no	2386	55	29.0	Outcome	Any HDP	BPD36 (32)	4	2	3	9
Kalra, 2014 ⁹¹	USA	Ca-Co	yes	60	2	27.0	Outcome	BW<P10	BPD36 (55)	4	1	3	8
Kanaan, 2018 ⁹²	USA	Cohort	no	1340	1	27.8	Outcome	Any HDP BW<P10	BPD-PH (12)	4	2	3	9
Kandasamy, 2015 ⁹³	USA	Cohort	yes	152	1	25.2	Outcome	BW<P10	BPD/Death (23)	4	1	3	8
Karagianni, 2011 ⁹⁴	Greece	Cohort	yes	219	1	29.1	Outcome	BW<P10	BPD36 (28)	4	1	3	8
Karagianni, 2013 ⁹⁵	Greece	Ca-Co	yes	61	1	32.0	Outcome	BW<P10	BPD36 (46)	4	1	3	8
Khan, 2006 ⁹⁶	USA	Cohort	no	306	2	26.7	Outcome	PE	BPD36 (45)	4	2	3	9
Khan, 2020 ⁹⁷	USA	Cohort	yes	68	1	27.0	Outcome	PE BW<P10	BPD36 (65)	4	2	3	9
Khemani, 2007 ⁹⁸	USA	Cohort	no	42	3	26.0	Outcome	BW<P3	BPD-PH (43)	3	1	3	7
Kim, 2014 ⁹⁹	Korea	Ca-Co	no	56	1	26.1	Outcome	PE BW<P3	sBPD (66) BPD-PH (27)	4	1	3	8

First author, year	Country	Design	Prospective?	Total infants	Centers	GA of cohort (weeks)	Independent variable	Exposures	Outcomes (% of incidence in total group)	NOS Selection	NOS Comparability	NOS Outcome/Exposure	NOS Total
Kim, 1996 ¹⁰⁰	USA	Ca-Co	no	117	1	28	Exposure	Any HDP	BPD28 (48) BPD36 (15)	3	1	3	7
Kim, 2018 ¹⁰¹	Korea	cohort	no	199	1	28.9	Exposure	PE	BPD36 (31)	4	2	3	9
Kim, 2020 ¹⁰²	Korea	Cohort	no	117	1	28.8	Outcome	BW<P10	BPD28 (91) sBPD (18)	3	1	3	7
Klinger, 2013 ¹⁰³	Israel	Cohort	yes	12139	28	29.0	Outcome	Any HDP BW<P10	BPD36 (14)	4	2	3	9
Köksal, 2012 ¹⁰⁴	Turkey	Cohort	yes	102	1	28.6	Outcome	PE	BPD28 (30)	4	1	3	8
Korhonen, 2004 ¹⁰⁵	Finland	Cohort	yes	68	1	27.5	Outcome	BW<P3	BPD36 (50)	4	1	2	7
Koroglu, 2013 ¹⁰⁶	Turkey	Ca-Co	yes	41	1	28.9	Outcome	PE	BPD28 (43)	3	2	3	8
Kunjunju, 2017 ¹⁰⁷	Australia	Cohort	no	56	1	26.0	Outcome	PE	BPD-PH (39)	3	1	3	7
Lagatta, 2018 ¹⁰⁸	USA	Cohort	No	1677	23	25.0	Outcome	Any HDP BW<P10	BPD-PH (22)	3	2	3	8
Lal, 2003 ¹⁰⁹	UK	Cohort	yes	2838	Network	29.9	Outcome	BW<P10	BPD28 (23) BPD36 (14) BPD/Death (23)	4	1	3	8
Lapcharoensap, 2015 ¹¹⁰	USA	Cohort	no	15052	132	27.0	Outcome	Any HDP IUGR	BPD/Death (47)	4	2	3	9
Lardon-Fdz, 2017 ¹¹¹	Spain	Cohort	no	129	1	28.7	Outcome	PE BW<P3	BPD28 (47) BPD36 (19) sBPD (7)	3	1	3	7
Lehtinen, 2017 ¹¹²	Finland	Cohort	yes	53	1	29.0	Outcome	BW<P3	BPD36 (40)	4	2	3	9
Leroy, 2018 ¹¹³	Canada	Cohort	yes	62	1	27.0	Outcome	BW<P10	BPD36 (52)	4	1	3	8
Li, 2013 ¹¹⁴	China	Cohort	yes	160	1	30.9	Outcome	PE BW<P10	BPD/Death (36)	4	2	3	9
Lin, 2005 ¹¹⁵	Taiwan	Ca-Co	yes	224	1	26.7	Outcome	BW<P3	BPD36 (50)	4	1	3	8

First author, year	Country	Design	Prospective?	Total infants	Centers	GA of cohort (weeks)	Independent variable	Exposures	Outcomes (% of incidence in total group)	NOS Selection	NOS Comparability	NOS Outcome/Exposure	NOS Total
Lodha, 2014 ¹¹⁶	Canada	Cohort	yes	586	1	28.5	Outcome	BW<P10	BPD36 (25)	4	1	2	7
Löfqvist, 2012 ¹¹⁷	Sweden	Cohort	yes	108	2	27.2	Outcome	PE BW<P3	BPD36 (54)	4	2	3	9
Lohmann, 2014 ¹¹⁸	USA	Cohort	yes	22	1	27.2	Outcome	Any HDP	BPD36 (45)	4	1	3	8
MacKenzie, 2020 ¹¹⁹	Canada	Cohort	No	87	1	25.9	Outcome	Any HDP	BPD-PH (28)	3	1	3	7
Madoglio, 2016 ¹²⁰	Brazil	Cohort	yes	73	1	28.7	Exposure	PE	BPD28 (36) BPD36 (18) BPD/Death (46)	3	1	3	7
Mahlman, 2017 ¹²¹	Finland	Ca-Co	yes	174	5	27.4	Outcome	PE BW<P3	BPD 36 (34)	4	1	3	8
Malavolti, 2018 ¹²²	Switzerland	Cohort	no	610	1	27.9	Outcome	BW<P10	BPD28 (59)	4	2	3	9
Mao, 2018 ¹²⁴	China	Ca-Co	yes	39	1	29.6	Outcome	PE BW<P10	BPD36 (49)	4	1	3	8
Marshall, 1999 ¹²⁴	USA	Cohort	yes	865	13	28.9	Outcome	PE	BPD36 (26)	3	2	3	8
Matic, 2017 ¹²⁵	Australia and UK	Cohort	no	2549	10	26.7	Exposure	Any HDP	BPD36 (30)	4	2	3	9
May, 2009 ¹²⁶	UK	Cohort	yes	80	1	28.0	Outcome	BW<P10	BPD28 (58) BPD36 (39) sBPD (16)	4	1	3	8
McGowan, 2009 ¹²⁷	USA	Ca-Co	no	98	1	26.3	Outcome	PE BW<P10	BPD36 (50)	4	2	3	9
Melamed, 2016 ¹²⁸	Canada	Cohort	no	6567	Network	29.8	Exposure	BW<P10	BPD36 (12)	4	2	3	9
Mello, 2017 ¹²⁹	Brasil	Cohort	yes	112	1	29.5	Outcome	BW<P10	BPD28 (44)	4	2	2	8
Méndez-Abad, 2020 ¹³⁰	Spain	Cohort	yes	101	1	29.0	Outcome	BW<P10	BPD36 (15)	4	1	3	8
Mohamed, 2015 ¹³¹	Canada	Cohort	yes	99	1	26.4	Outcome	PE BW<P10	BPD36 (66)	4	1	3	8

First author, year	Country	Design	Prospective?	Total infants	Centers	GA of cohort (weeks)	Independent variable	Exposures	Outcomes (% of incidence in total group)	NOS Selection	NOS Comparability	NOS Outcome/Exposure	NOS Total
Monier, 2017 ¹³²	France	Cohort	yes	5919	Network	29.0	Exposure	IUGR	BPD36 (48) sBPD (7)	4	2	3	9
Morrow, 2017 ¹³³	USA	Cohort	yes	587	5	27.0	Outcome	PE BW<P10	BPD36 (41)	4	2	3	9
Morsing, 2012 ¹³⁴	Sweden	Ca-Co	yes	62	1	27.0	Exposure	IUGR	sBPD (50)	3	1	3	7
Mourani, 2015 ¹³⁵	USA	Cohort	yes	277	2	27.0	Outcome	PE BW<P10	BPD-PH (14)	4	1	3	8
Naeem, 2020 ¹³⁶	USA	Cohort	yes	26	N	26.1	Outcome	Any HDP	BPD36 (72) BPD-PH (17)	4	1	3	8
Nascimento, 2020 ¹³⁷	UK	Cohort	yes	40	1	28.0	Outcome	BW<P10	BPD36 (53)	4	1	3	8
Natarajan, 2012 ¹³⁸	USA	Cohort	yes	1159	Network	25.7	Outcome	BW<P10	BPD36 (13)	4	2	3	9
Naveda, 2016 ¹³⁹	Venezuela	Ca-Co	yes	144	1	33.0	Outcome	Any HDP	BPD28 (25)	3	2	3	8
Newman, 2011 ¹⁴⁰	USA	Cohort	no	156	1	28.6	Outcome	Any HDP BW<P10	BPD28 (51) sBPD (31)	4	2	3	9
Nobile, 2017 ¹⁴¹	Italy	Cohort	no	515	1	28.0	Exposure	IUGR BW<P10	BPD36 (22) BPD/Death (38)	4	2	3	9
Novitsky, 2015 ¹⁴²	USA	Cohort	no	906	1	28.2	Outcome	PE	BPD36 (20)	3	2	3	8
O'Shea, 2012 ¹⁴³	Australia	Cohort	no	751	Network	26.6	Outcome	PE BW<P3	BPD36 (44)	4	2	3	9
Oh, 2005 ¹⁴⁴	USA	Cohort	no	1382	Network	26.0	Outcome	Any HDP	BPD/Death (58)	4	2	3	9
Ozkan, 2012 ¹⁴⁵	Turkey	Cohort	yes	332	1	29.2	Exposure	PE	BPD28 (26) BPD36 (14) sBPD (8)	4	2	3	9
Palta, 1996 ¹⁴⁶	USA	Cohort	no	632	7	28.7	Exposure	PE-HELLP	BPD28 (24)	3	1	3	7
Park, 2012 ¹⁴⁷	Korea	Cohort	no	191	1	29.4	Exposure	Any HDP	BPD28 (36) BPD36 (29)	4	1	3	8

First author, year	Country	Design	Prospective?	Total infants	Centers	GA of cohort (weeks)	Independent variable	Exposures	Outcomes (% of incidence in total group)	NOS Selection	NOS Comparability	NOS Outcome/Exposure	NOS Total
Patel, 2017 ¹⁴⁸	USA	Cohort	yes	254	1	27.9	Outcome	BW<P10	BPD36 (30)	4	2	3	9
Qi, 2013 ¹⁴⁹	China	Cohort	yes	60	1	29.5	Outcome	Any HDP BW<P10	BPD36 (42)	4	1	3	8
Qiu, 2012 ¹⁵⁰	Canada	Cohort	no	11909	Network	28.8	Exposure	BW<P10	BPD36 (21) BPD/Death (30)	4	2	3	9
Ra, 2013 ¹⁵¹	Korea	Cohort	no	85	1	28.0	Outcome	Any HDP	BPD-PH (21)	3	1	3	7
Rakza, 2007 ¹⁵²	France	Cohort	yes	48	1	29.1	Exposure	IUGR	BPD36 (21) BPD/Death (29)	4	1	3	8
Ranke, 2007 ¹⁵³	Germany	Cohort	yes	97	1	29.0	Exposure	BW<P10	BPD36 (39)	4	1	2	7
Razak 2020 ¹⁵⁴	Canada	Ca-Co	no	237	1	29.2	Exposure	Any HDP	BPD36 (20)	4	2	3	9
Redline, 2002 ¹⁵⁵	USA	cohort	no	371	1	27.6	Outcome	PE	BPD36 (30)	4	1	3	8
Regev, 2015 ¹⁵⁶	Israel	cohort	yes	2139	28	29.7	Exposure	Any HDP	BPD36 (12)	4	1	3	8
Regev, 2003 ¹⁵⁷	Israel	Cohort	yes	2764	Network	27.9	Exposure	BW<P3	BPD36 (22) BPD/Death (43)	4	2	3	9
Reiss, 2003 ¹⁵⁸	Germany	Cohort	yes	1365	1	28.8	Exposure	BW<P10	BPD36 (14)	4	2	3	9
Rijken, 2007 ¹⁵⁹	The Netherlands	Cohort	yes	158	1	28.8	Exposure	BW<P10	BPD36 (22)	4	2	3	9
Rocha, 2018 ¹⁶⁰	Portugal	Cohort	yes	494	11	27.3	Exposure	Any HDP	BPD36 (24)	4	2	3	9
Rocha, 2010 ¹⁶¹	Portugal	Cohort	yes	205	1	29.0	Outcome	BW<P10	BPD36 (22)	4	1	3	8
Rojas, 2012 ¹⁶²	Colombia	Ca-Co	no	212	8	29.2	Outcome	BW<P10	BPD36 (30)	4	2	3	9
Ruiz-Pelaez, 2014 ¹⁶³	Colombia	Cohort	yes	416	12	31.9	Outcome	BW<P10	BPD28 (54) BPD36 (17)	3	1	3	7
Sasi, 2015 ¹⁶⁴	Australia	Cohort	no	459	1	28.7	Exposure	IUGR	BPD36 (27) BPD/Death (30)	4	1	3	8
Schena, 2015 ¹⁶⁵	Italy	Cohort	no	242	1	26.3	Outcome	BW<P10	BPD/Death (43)	4	1	3	8

First author, year	Country	Design	Prospective?	Total infants	Centers	GA of cohort (weeks)	Independent variable	Exposures	Outcomes (% of incidence in total group)	NOS Selection	NOS Comparability	NOS Outcome/Exposure	NOS Total
Schiff, 1993 ¹⁶⁶	USA	Ca-Co	yes	138	1	33.8	Exposure	PE-HELLP	BPD28 (1)	4	2	3	9
Schlapbach, 2010 ¹⁶⁷	Switzerland	Ca-Co	yes	99	1	29.3	Exposure	PE	BPD36 (14)	4	1	3	8
Seo, 2020 ¹⁶⁸	Korea	Cohort	no	81	1	26.4	Outcome	Any HDP BW<P10	BPD-PH (30)	4	1	2	7
Serenius, 2004 ¹⁶⁹	Sweden	Cohort	no	140	2	24.2	Outcome	BW<P3	BPD36 (36)	4	2	3	9
Sharma, 2004 ¹⁷⁰	USA	Cohort	no	2364	1	31.2	Exposure	BW<P10	BPD28 (?) BPD36 (?)	4	1	3	8
Sheth, 2020 ¹⁷¹	USA	Ca-Co	no	220	1	25.9	Outcome	PE BW<P10	BPD-PH (27)	4	2	3	9
Shima, 2013 ¹⁷²	Japan	Cohort	no	306	1	29.0	Outcome	Any HDP BW<P10	BPD28 (42) BPD36 (17)	3	1	3	7
Shin, 2020 ¹⁷³	Korea	Cohort	yes	1827	Network	27.4	Exposure	Any HDP	BPD36 (42)	4	2	3	9
Silveira, 2007 ¹⁷⁴	Brazil	Cohort	yes	86	1	30.9	Exposure	PE	BPD36 (14)	3	1	3	7
Soliman, 2017 ¹⁷⁵	Canada	Cohort	yes	319	1	28.8	Exposure	PE-HELLP BW<P10	BPD28 (39) BPD36 (25)	4	1	3	8
Somaschini, 2012 ¹⁷⁶	Italy	Cohort	no	366	12	28.5	Outcome	Any HDP BW<P10	BPD 28 (39) BPD 36 (21) sBPD (10)	3	1	3	7
Soudee, 2014 ¹⁷⁷	France	Ca-Co	no	293	1	28.5	Exposure	BW<P3	BPD28 (57) BPD36 (24) BPD/Death (37)	4	2	3	9
Spiegler, 2013 ¹⁷⁸	Germany	Cohort	no	1577	28	29.2	Exposure	PE	BPD36 (12)	3	1	3	7
Starr, 2020 ¹⁷⁹	USA	Cohort	no	546	24	27.9	Outcome	Any HDP BW<P10	BPD/Death (45)	4	2	3	9
Streubel, 2008 ¹⁸⁰	USA	Cohort	yes	133	1	26.6	Outcome	BW<P10	BPD36 (29)	4	1	3	8
Sun, 2019 ¹⁸¹	China	Cohort	no	296	1	30.0	Outcome	Any HDP BW<P10	BPD28 (49)	4	1	3	8

First author, year	Country	Design	Prospective?	Total infants	Centers	GA of cohort (weeks)	Independent variable	Exposures	Outcomes (% of incidence in total group)	NOS Selection	NOS Comparability	NOS Outcome/Exposure	NOS Total
Tammela, 1992 ¹⁸²	Finland	Cohort	yes	46	1	31.7	Outcome	BW<P3	BPD36 (50)	4	1	3	8
Teberg, 1991 ¹⁸³	USA	Cohort	yes	236	1	30.4	Outcome	Any HDP BW<P10	BPD28 (25)	4	2	3	9
Todd, 1997 ¹⁸⁴	Australia	Ca-Co	yes	296	1	28.0	Outcome	Any HDP	BPD36 (50)	3	1	3	7
Tokumasu, 2016 ¹⁸⁵	Japan	Cohort	no	4518	79	26.1	Exposure	PE-HELLP	BPD36 (25)	4	1	3	8
Torchin, 2016 ¹⁸⁶	France	Cohort	yes	2638	?	29.7	Exposure	Any HDP IUGR	BPD36 (10)	4	1	3	8
Torrance, 2007 ¹⁸⁷	Netherlands	Cohort	no	187	1	30.4	Exposure	Any HDP	BPD28 (23)	4	2	3	9
Tsai, 2015 ¹⁸⁸	Taiwan	Cohort	yes	1680	21	28.3	Exposure	BW<P10	BPD36 (32) BPD/Death (47)	4	2	3	9
Turunen, 2011 ¹⁸⁹	Finland	Ca-Co	yes	46	1	26.9	Exposure	PE	BPD36 (70)	4	2	3	9
Uberos, 2020 ¹⁹⁰	Spain	Cohort	no	389	1	29.0	Outcome	Any HDP BW<P10	BPD36 (41)	4	2	3	9
Van Mastright , 2018 ¹⁹¹	The Netherlands	Cohort	yes	111	1	28.0	Outcome	PE	BPD/Death (40)	4	2	3	9
Vayaltrikkovil, 2019 ¹⁹²	Canada	Cohort	Yes	126	1	26.2	Outcome	Any HDP BW<P10	BPD-PH (19)	3	1	3	7
Viscardi, 2004 ¹⁹³	USA	Cohort	yes	262	2	28.0	Outcome	PE BW<P10	BPD28 (58)	4	2	3	9
Vrlenich, 1995 ¹⁹⁴	USA	Cohort	yes	406	10	30.3	Outcome	BW<P3	BPD36 (23)	4	1	2	7
Wadhawan, 2007 ¹⁹⁵	USA	Cohort	no	9461	19	26.0	Exposure	BW<P10	BPD36 (42) BPD/Death (55)	4	1	3	8
Wang, 2014 ¹⁹⁶	China	Cohort	yes	73	1	30.5	Outcome	Any HDP	BPD28 (33)	3	1	3	7
Wang, 2018 ¹⁹⁷	China	Cohort	no	191	1	28.2	Outcome	Any HDP BW<P3	BPD-PH (19)	3	1	3	7
Wang, 2013 ¹⁹⁸	USA	Ca-Co	no	1649	Network	27.0	Outcome	IUGR	BPD36 (52)	4	2	3	9

First author, year	Country	Design	Prospective?	Total infants	Centers	GA of cohort (weeks)	Independent variable	Exposures	Outcomes (% of incidence in total group)	NOS Selection	NOS Comparability	NOS Outcome/Exposure	NOS Total
Wang, 2010 ¹⁹⁹	Taiwan	Cohort	yes	72	2	28.3	Outcome	BW<P10	BPD28 (78) BPD36 (52) sBPD (36)	4	2	3	9
Weismann, 2017 ²⁰⁰	USA	Cohort	yes	159	1	25.6	Outcome	Any HDP BW<P10	BPD-PH (28)	4	2	3	9
Wemhöner, 2011 ²⁰¹	Austria	Cohort	no	95	1	27.7	Outcome	BW<P10	BPD36	4	1	3	8
Westby Wold, 2009 ²⁰²	Norway	Cohort	no	365	15	26.0	Exposure	BW<P5	BPD36 (40) BPD/Death (60)	4	1	3	8
Wilmink 2019 ²⁰³	The Netherlands	Cohort	no	273	1	29.3	Exposure	PE	BPD28 (24) BPD36 (11) BPD/Death (19)	4	2	3	9
Wirbelauer, 2010 ²⁰⁴	Germany	Cohort	yes	49	1	27.7	Exposure	BW<P10	BPD28 (29)	4	1	3	8
Withagen, 2001 ²⁰⁵	Netherlands	Ca-Co	no	666	1	30.7	Exposure	PE-HELLP	BPD28 (13)	4	2	3	9
Wocadlo, 1994 ²⁰⁶	Australia	Cohort	yes	36	1	27.6	Exposure	BW<P10	BPD28 (50)	4	1	2	7
Won Choi, 2006 ²⁰⁷	Korea	Cohort	yes	75	1	28.5	Outcome	Any HDP	BPD36 (35)	3	1	3	7
Xu, 2016 ²⁰⁸	China	Cohort	no	42	1	28.0	Outcome	BW<P10	BPD36 (76) sBPD (36)	3	1	3	7
Xydis, 2013 ²⁰⁹	Greece	Cohort	yes	205	1	32.9	Exposure	BW<P3	BPD36 (59)	4	2	3	9
Yallapragada, 2016 ²¹⁰	USA	Ca-Co	yes	14	1	26.2	Outcome	PE BW<P10	BPD-PH (50)	4	1	3	8
Yen, 2013 ²¹¹	Taiwan	Cohort	no	5753	21	30.0	Exposure	PE	BPD36 (35)	3	1	3	7
Yilmaz, 2017 ²¹²	Turkey	Cohort	yes	40	1	30.2	Outcome	PE BW<P10	BPD28 (35)	4	1	3	8
Yoon, 1999 ²¹³	Korea	Cohort	yes	203	1	31.0	Outcome	BW<P10	BPD28 (17)	4	2	3	9
Yu, 2011 ²¹⁴	Korea	Cohort	no	415	1	26.2	Exposure	BW<P3	BPD36 (37)	4	2	3	9
Zanardo, 2002 ²¹⁵	Italy	Ca-Co	no	100	1	27.7	Outcome	BW<P10	BPD28 (50)	3	1	3	7

First author, year	Country	Design	Prospective?	Total infants	Centers	GA of cohort (weeks)	Independent variable	Exposures	Outcomes (% of incidence in total group)	NOS Selection	NOS Comparability	NOS Outcome/Exposure	NOS Total
Zhang, 2011 ²¹⁶	China	cohort	no	116	1	30.2	Outcome	Any HDP BW<P10	BPD28 (48)	3	1	3	7

BPD28: BPD defined as oxygen requirement on postnatal day 28; BPD36: defined as oxygen requirement at the postmenstrual age of 36 weeks; BPD-PH: BPD-associated pulmonary hypertension; BW: birth weight; HDP: hypertensive disorders of pregnancy; IUGR: intrauterine growth restriction (defined on the basis of fetal growth assessment); NOS: Newcastle Ottawa Scale; P3: 3rd percentile; P5: 5th percentile; P10: 10th percentile; PE: preeclampsia; sBPD: severe BPD (\geq 30% oxygen or mechanical ventilation at the postmenstrual age of 36 weeks); SGA: small for gestational age. The number in parentheses after the outcome indicates the outcome rate in the study.

Supplementary Table 2. Meta-regression analysis (continuous covariates)

Association	Covariate	Covariate unit	K	Coefficient	95% CI		P	R ² analog
					Lower limit	Upper limit		
HDP and BPD36	MD in GA (exposed minus unexposed)	week	24	-0.500	-0.651	-0.349	<0.001	0.95
	GA total cohort (weeks)	week	55	-0.077	-0.197	0.043	0.112	0.29
	Male sex (logOR)	logOR	20	0.190	-0.670	1.050	0.665	0.00
	ACS (logOR)	logOR	17	-0.148	-0.578	0.282	0.501	0.00
	RDS (logOR)	logOR	12	0.266	-0.185	0.716	0.248	0.00
	Median year of cohort	year	60	0.008	-0.018	0.035	0.426	0.00
SGA/IUGR and BPD36	MD in GA (exposed minus unexposed)	week	35	-0.311	-0.571	-0.051	0.002	0.08
	GA total cohort	week	91	-0.039	-0.133	0.056	0.295	0.00
	Male sex (logOR)	logOR	21	-0.222	-1.374	0.930	0.705	0.02
	ACS (logOR)	logOR	18	-0.371	-0.742	0.000	0.050	0.08
	RDS (logOR)	logOR	16	0.354	-0.288	0.997	0.280	0.00
	Median year of cohort	year	89	0.021	-0.001	0.042	0.056	0.01

Random effects (method of moments), univariate meta-regression. Coefficient indicates the change in the log of the OR of the association between BPD36 and the corresponding exposure for a unit change in the predictor covariate. R² analog: total between-study variance explained by the moderator. ACS: antenatal corticosteroids; BPD36: bronchopulmonary dysplasia defined as oxygen requirement at the postmenstrual age of 36 weeks; BW: birth weight; CI: confidence interval; GA: gestational age; HDP: hypertensive disorders of pregnancy; IUGR: intrauterine growth restriction (defined on the basis of fetal growth assessment); K: number of studies; OR: odds ratio; SGA: small for gestational age.

Supplementary table 3. Meta-analyses on the association between SGA/IUGR and HDP

Meta-analysis	Subgroup	K	OR	95% CI		P	Heterogeneity	
				Lower limit	Upper limit		I ² (%)	P
Exposure: SGA/IUGR Outcome: HDP	BW<P10	12	5.00	3.96	6.31	<0.001	87.1	<0.001
	BW<P5	2	8.31	4.53	15.23	<0.001	0.0	0.645
	BW<P3 or -2SD	3	6.78	4.29	10.70	<0.001	0.0	0.670
	IUGR	4	5.71	4.23	7.71	<0.001	73.4	0.010
	SGA/IUGR overall	21	5.62	4.77	6.62	<0.001	80.0	<0.001
Exposure: HDP Outcome: SGA/IUGR	Any HDP	3	6.22	2.32	13.79	<0.001	80.3	0.006
	Preeclampsia	16	4.06	2.65	6.22	<0.001	98.0	<0.001
	Preeclampsia -HELLP	3	11.07	3.84	31.94	<0.001	91.8	<0.001
	HDP overall	22	4.86	3.36	7.01	<0.001	97.6	<0.001

Mixed effects analysis. A random effects model is used to combine studies within each subgroup. A fixed effect model is used to combine subgroups and yield the overall effect. OR>1 indicates association with increased risk of the outcome and OR<1 indicates association with decreased risk of the outcome.

CI: confidence interval; GA: gestational age; HDP: hypertensive disorders of pregnancy; IUGR: intrauterine growth restriction (defined on the basis of fetal growth assessment); K: number of studies; OR: odds ratio; SGA: small for gestational age.

Supplementary table 4. Raw count data of the meta-analysis on the association between HDP and BPD28.

Study Name	BPD28-yes/total		Study Name	BPD28-yes/total	
	HDP-yes	HDP-no		HDP-yes	HDP-no
Abramovici 1999 ⁶	21/47	277/419	Naveda 2016 ¹³⁹	29/76	7/68
Bursal 2017 ²⁹	13/140	16/144	Newman 2011 ¹⁴⁰	18/35	62/121
Cheng 2004 ³⁸	10/28	22/61	Ozkan 2012 ¹⁴⁵	45/117	42/215
Cunha 2005 ⁴³	11/29	34/57	Palta 1996 ¹⁴⁶	18/121	134/511
Demirel 2009 ⁴⁸	1/2	55/104	Park 2012 ¹⁴⁷	11/38	58/153
Duan 2016 ⁵¹	16/48	55/195	Schiff 1993 ¹⁶⁶	2/69	0/69
Gray 1997 ⁷¹	17/26	61/132	Shima 2013 ¹⁷²	21/48	109/258
Gursoy 2015 ⁷⁶	18/108	132/544	Soliman 2017 ¹⁷⁵	39/102	86/217
Hentges 2015 ⁸⁰	6/37	17/51	Sun 2019 ¹⁸¹	20/48	124/248
Hernandez 2004 ⁸¹	5/11	17/33	Teberg 1991 ¹⁸³	3/44	57/189
Hiett 2001 ⁸²	40/58	30/58	Torrance 2007 ¹⁸⁷	33/120	10/67
Kim 1996 ¹⁰⁰	16/35	34/70	Viscardi 2004 ¹⁹³	15/36	136/226
Koksal 2012 ¹⁰⁴	12/36	19/66	Wang 2014 ¹⁹⁶	7/24	17/49
Koroglu 2014 ¹⁰⁶	10/20	37/89	Wilmink 2019 ²⁰³	23/90	36/157
Lardon-F 2017 ¹¹¹	21/47	39/82	Withagen 2001 ²⁰⁵	33/222	51/444
Madoglio 2016- ¹²⁰	10/32	16/41	Yilmaz 2017 ²¹²	4/12	10/28
			Zhang 2011 ²¹⁶	14/33	42/83

Supplementary table 5. Raw count data of the meta-analysis on the association between SGA/IUGR and BPD28.

Study Name	BPD28-yes/total		Study Name	BPD28-yes/total	
	SGA/IUGR yes	SGA/IUGR no		SGA/IUGR yes	SGA/IUGR no
Amin 1997 ⁹	26/62	74/124	Newman 2011 ¹⁴⁰	8/16	72/140
Bardin 1997 ¹⁶	20/20	79/95	Ruiz-P 2014 ¹⁶³	OR 1.09 (95% CI 0.42-3.16)	
Bose 2011 ²⁰	39/52	447/880	Sharma 2004 ¹⁷⁰	OR 1.09 (95% CI 0.64-1.86)	
Bozzetti 2013 ²³	4/32	25/278	Shima 2013 ¹⁷²	23 /48	107/258
Burns 1997 ²⁸	5/12	59/105	Somaschini 2012 ¹⁷⁶	13/46	128/320
Charafeddine 1999 ³⁴	7/36	81/141	Soudee 2014 ¹⁷⁷	22/39	120/208
Chye 1995 ⁴¹	5/14	73/142	Sun 2019 ¹⁸¹	21/41	123/255
Enginner 2010 ⁵⁵	15/44	20/77	Teberg 1991 ¹⁸³	16/60	44/176
Garite 2004 ⁶⁵	287/1451	4068/22978	Viscardi 2004 ¹⁹³	6/15	145/247
Gortner 2003 ⁷⁰	7/74	2/74	Wang L.Y. 2010 ¹⁹⁹	15/25	41/47
Greenough 2005 ⁷²	1/2	16/32	Wirbelauer 2010 ²⁰⁴	7/18	7/31
Gursoy 2015 ⁷⁶	12/55	138/597	Wocadlo 1994 ²⁰⁶	9/18	9/18
Kim 2020 ¹⁰²	17/19	90/98	Xydis 2013 ²⁰⁹	3/33	13/172
Lal 2003 ¹⁰⁹	96/333	497/2212	Yilmaz 2017 ²¹²	2/8	12/32
Malavolti 2018 ¹²²	32/53	325/557	Yoon 1999 ²¹³	7/42	27/161
May 2009 ¹²⁶	9/14	37/66	Zanardo 2002 ²¹⁵	13/18	37/82
Mello 2017 ¹²⁹	8/49	41/63	Zhang 2011 ²¹⁶	11/36	45/80

Supplementary table 6. Raw count data of the meta-analysis on the association between HDP and BPD36.

Study Name	BPD36-yes/total		Study Name	BPD36-yes/total	
	HDP-yes	HDP-no		HDP-yes	HDP-no
Ameenudeen 2007 ⁸	11/74	23/159	Mao 2018 ¹²³	8/13	11/26
Baker 2012 ¹⁴	3/13	10/49	Marshall 1999 ¹²⁴	31/178	193/687
Bose 2009 ¹⁹	108/167	538/1074	Matic 2017 ¹²⁵	137/379	624/2170
Bossung ²¹	368/2562	2030/13383	McGowan 2009 ¹²⁷	6/22	41/74
Cederqvist 2003 ³²	9/11	9/21	Mohamed 2015 ¹³¹	12/16	51/80
Çetinkaya 2010 ³³	13/51	6/33	Morrow 2017 ¹³³	56/154	186/433
Da Silva 2018 ⁴⁴	7/25	15/42	Naeem 2020 ¹³⁶	17/24	9/12
Dasgupta 2018 ⁴⁵	20/35	16/17	Novitsky 2015 ¹⁴²	42/252	140/654
De Souza 2015 ⁴⁷	3/30	3/30	O'Shea 2012 ¹⁴³	52/138	280/613
Dogan 2020 ⁵⁰	11/18	21/60	Ozkan 2012 ¹⁴⁵	26/117	19/215
Duan 2016b ⁵¹	10/27	51/120	Park 2012 ¹⁴⁷	9/38	46/153
Eriksson 2015 ⁵⁶	139/534	395/1721	Qi 2013 ¹⁴⁹	3/11	22/49
Falciglia 2003 ⁵⁹	2/7	21/39	Razak 2020 ¹⁵⁴	21/79	27/158
Fujioka 2014 ⁶¹	13/22	42/75	Redline 2002 ¹⁵⁵	22/71	90/300
Gagliardi 2013 ⁶³	63/441	204/1644	Regev 2015 ¹⁵⁶	146/929	105/864
Gagliardi 2014 ⁶⁴	459/2096	428/1510	Rocha 2018 ¹⁶⁰	27/75	92/419
Gemmell 2016 ⁶⁷	1523/3625	8235/24221	Schlapbach 2010 ¹⁶⁷	5/33	9/66
Hansen 2010 ⁷⁸	12/29	15/78	Shima 2013 ¹⁷²	8/48	45/258
Iwatani 2013 ⁸⁶	7/10	19/41	Shin 2020 ¹⁷³	91/203	673/1624
Jeon 2020 ⁸⁹	31/55	221/466	Silveira 2007 ¹⁷⁴	4/40	8/46
Jo 2015 ⁹⁰	106/446	583/1690	Soliman 2017 ¹⁷⁵	23/102	56/217
Khan 2006 ⁹⁶	16/34	121/272	Spiegler 2013 ¹⁷⁸	48/353	134/922
Khan 2020 ⁹⁷	11/16	33/52	Todd 1997 ¹⁸⁴	35/57	113/239
Kim 1996 ¹⁰⁰	5/35	11/70	Tokumasu 2016 ¹⁸⁵	136/331	1003/2887
Kim 2018 ¹⁰¹	24/59	37/140	Torchin 2016 ¹⁸⁶	79/605	180/1506
Klinger 2013 ¹⁰³	329/2470	1334/9669	Turunen 2011 ¹⁸⁹	11/21	21/25
Lardon-F 2017 ¹¹¹	10/47	15/82	Uberos 2020 ¹⁹⁰	9/28	150/361
Löfqvist 2012 ¹¹⁷	9/23	49/85	Wilmink 2019 ²⁰³	12/90	15/157
Lohmann 2014 ¹¹⁸	5/7	5/15	Won Choi 2006 ²⁰⁷	6/19	20/56
Madoglio 2016 ¹²⁰	4/32	9/41	Yen 2013 ²¹¹	204/847	1802/4906
Mahlman 2017 ¹²¹	23/47	37/127			

Supplementary table 7. Raw count data of the meta-analysis on the association between SGA/IUGR and BPD36.

Study Name	BPD36-yes/total		Study Name	BPD36-yes/total	
	SGA/IUGR yes	SGA/IUGR no		SGA/IUGR yes	SGA/IUGR no
Altman 2011 ⁷	24/840	44/5834	Mahlman 2017 ¹²¹	23/43	37/131
Ameenudeen 2007 ⁸	3/40	31/193	Mao 2018 ¹²³	7/12	12/27
Aucott 2004 ¹²	9/31	11/64	May 2009 ¹²⁶	9/14	22/66
Baer 2016 ¹³	179/8418	1004/70233	McGowan 2009 ¹²⁷	5/5	44/93
Baker 2012 ¹⁴	3/6	10/56	Melamed 2016 ¹²⁸	164/918	652/5649
Bardin 1997 ¹⁶	13/20	30/95	Mendez 2020 ¹³⁰	4/13	11/88
Bose 2009 ¹⁹	60/77	579/1181	Mohamed 2015 ¹³¹	13/17	50/79
Brener 2017 ²⁴	9/45	36/158	Monier 2017 ¹³²	362/720	1028/2199
Brumbaugh 2018 ²⁵	10/13	115/138	Morrow 2017 ¹³³	66/158	176/429
Brumbaugh 2020 ²⁶	92/131	1010/2179	Nascimento 2020 ¹³⁷	2/8	19/32
Candel 2016 ³⁰	17/72	0/57	Natarajan 2012 ¹³⁸	71/151	80/1008
Chen 2016 ³⁷	OR 2.15 (95% CI 1.75-2.64)		Nobile 2017 ¹⁴¹	31/98	83/417
Claas 2011 ⁴²	9/21	48/80	O'Shea 2012 ¹⁴³	43/127	289/624
De Jesus 2013 ⁴⁶	145/385	928/2586	Patel 2017 ¹⁴⁸	18/55	59/199
Duan 2016b ⁵¹	5/11	56/136	Qi 2013 ¹⁴⁹	3/7	22/53
Durrmeyer 2012 ⁵³	12/23	42/328	Qiu 2012 ¹⁵⁰	322/1249	2201/10660
El Ayoubi 2016 ⁵⁴	110/446	511/4139	Rakza 2007 ¹⁵²	4/14	6/30
Eriksson 2015 ⁵⁶	185/482	476/1757	Ranke 2007 ¹⁵³	20/51	18/46
Estevez 2017 ⁵⁷	4/34	1/76	Regev 2003 ¹⁵⁷	OR 3.42 (95% CI 2.28-5.12)	
Fernandez-Rodriguez, 2021 ⁶⁰	3/23	5/71	Reiss 2003 ¹⁵⁸	40/183	145/1182
Fujioka 2014 ⁶¹	31/43	24/54	Rijken 2007 ¹⁵⁹	5/23	30/135
Gagliardi 2009 ⁶²	33/100	159/1109	Rocha 2010 ¹⁶¹	9/53	36/152
Garofoli 2014 ⁶⁶	3/35	4/41	Rojas 2012 ¹⁶⁰	3/12	61/200
Giapros 2012 ⁶⁸	20/35	39/133	Ruiz-P 2014 ¹⁶³	OR 2.88 (95% CI 1.20-6.91)	
Gortner 1999 ⁶⁹	9/59	11/258	Sasi 2015 ¹⁶⁴	69/153	54/306
Grisaru-G 2012 ⁷³	106/408	865/7505	Serenius 2004 ¹⁶⁹	OR 1.97 (95% CI 0.81-4.79)	
Guimaraes 2010 ⁷⁴	7/47	40/209	Sharma 2004 ¹⁷⁰	OR 2.20 (95% CI 1.20-4.03)	
Hartung 2005 ⁷⁹	9/44	0/44	Shima 2013 ¹⁷²	15/48	38/258
Hikino 2012 ⁸³	0/5	13/21	Soliman 2017 ¹⁷⁵	13/24	66/295
Huusko 2015 ⁸⁴ (Canada)	11/14	47/112	Somaschini 2012 ¹⁷⁶	8/46	68/318
Huusko 2015 ⁸⁴ (Finland)	56/156	103/493	Soudee 2014 ¹⁷⁷	16/35	40/194
Jakuskiene 2011 ⁸⁷	OR 4.10 (95% CI 1.19-14.15)		Streubel 2008 ¹⁸⁰	5/19	33/114
Jensen 2019 ⁸⁸	288/631	1494/5633	Tammela 1992 ¹⁸²	2/17	21/29
Kalra 2014 ⁹¹	5/7	28/53	Torchin 2016 ¹⁸⁶	90/481	169/1630
Karagianni 2011 ⁹⁴	13/52	48/167	Tsai 2015 ¹⁸⁸	230/451	306/992
Karagianni 2013 ⁹⁵	6/9	22/52	Uberos 2020 ¹⁹⁰	25/79	134/310
Khan 2020 ⁹⁷	3/4	41/64	Vrilenich 1995 ¹⁹⁴	20/125	75/281
Klinger 2013 ¹⁰³	384/2515	1279/9624	Wadhawan 2007 ¹⁹⁵	517/1248	3452/8213
Korhonen 2004 ¹⁰⁵	6/20	28/48	Wang 2013 ¹⁹⁸	72/122	783/1527
Lal 2003 ¹⁰⁹	74/348	274/2195	Wang L.Y. 2010 ¹⁹⁹	10/25	41/47
Lardon-Fdz 2017 ¹¹¹	18/40	47/89	Wemhoner 2011 ²⁰¹	3/8	23/87
Lehtinen 2017 ¹¹²	12/25	9/28	Westby W 2009 ²⁰²	15/31	130/334

Leroy 2018 ¹¹³	7/9	25/53	Xu 2016 ²⁰⁸	2/4	30/38
Lin 2005 ¹¹⁵	28/62	84/162	Xydis 2013 ²⁰⁹	14/33	106/172
Lodha 2014 ¹¹⁶	78/183	500/847	Yu 2011 ²¹⁴	13/49	142/366
Lofqvist 2012 ¹¹⁷	16/26	42/82			

Supplementary table 8. Raw count data of the meta-analysis on the association between HDP and severe BPD.

Study Name	Severe BPD yes/total		Study Name	Severe BPD yes/total	
	HDP-yes	HDP-no		HDP-yes	HDP-no
EXPRESS 2010 ⁵⁸	OR 2.00 (95% CI 1.00-4.00)		Newman 2011 ¹⁴⁰	8/35	40/111
Guo 2015 ⁷⁵	4/10	44/65	Ozkan 2012 ¹⁴⁵	14/117	16/215
Kim 2014 ⁹⁹	3/5	34/51	Park 2012 ¹⁴⁷	5/31	33/160
Lardon-F 2017 ¹¹¹	4/47	5/82	Somaschini 2017 ¹⁷⁶	4/62	32/304

Supplementary table 9. Raw count data of the meta-analysis on the association between SGA/IUGR and severe BPD.

Study Name	Severe BPD yes/total		Study Name	Severe BPD yes/total	
	SGA/IUGR yes	SGA/IUGR no		SGA/IUGR yes	SGA/IUGR no
Bose 2011 ²⁰	11/52	75/880	Lardon-F 2017 ¹¹¹	4/40	5/89
Brumbaugh 2018 ²⁵	8/13	75/138	May 2009 ¹²⁶	5/14	9/71
Brumbaugh 2020 ²⁶	16/131	103/2179	Monier 2017 ¹³²	72/720	131/2199
EXPRESS 2010 ⁵⁸	OR 1.90 (95% CI 1.10-3.28)		Morsing ¹³⁴	21/30	10/32
Guo 2015 ⁷⁵	5/11	43/67	Newman 2011 ¹⁴⁰	5/16	43/140
Hadchouel 2020 ⁷⁷	3/3	8/16	Somaschini 2017 ¹⁷⁶	4/46	32/320
Imamura 2017 ⁸⁵	16/32	51/137	Wang 2010 ¹⁹⁹	3/15	17/41
Kim 2014 ⁹⁹	6/7	31/49	Xu 2016 ²⁰⁸	2/4	13/38
Kim 2020 ¹⁰²	5/21	16/98			

Supplementary table 10. Raw count data of the meta-analysis on the association between HDP and BPD36 or death.

Study Name	BPD36 or death yes/total		Study Name	BPD36 or death yes/total	
	HDP-yes	HDP-no		HDP-yes	HDP-no
Balena 2017 ¹⁵	25/66	16/47	Oh 2005 ¹⁴⁴	166/354	631/1028
Lapcharoensap 2015 ¹¹⁰	1172/2864	5909/12188	Starr 2020 ¹⁷⁹	28/63	218/483
Li 2013 ¹¹⁴	7/21	51/139	Van Mastrigt 2018 ¹⁹¹	11/25	33/86
Madoglio 2016 ¹²⁰	13/32	20/40	Wilmink 2019 ²⁰³	21/99	32/174

Supplementary table 11. Raw count data of the meta-analysis on the association between SGA/IUGR and BPD36 or death.

Study Name	BPD36 or death-yes/total		Study Name	BPD36 or death-yes/total	
	SGA/IUGR yes	SGA/IUGR no		SGA/IUGR yes	SGA/IUGR no
Botet 2012 ²²	37/149	79/266	Rakza 2007 ¹⁵²	7/17	7/31
Durrmeyer 2012 ⁵³	13/24	86/372	Regev 2003 ¹⁵⁶	235/406	1019/2538
Jensen 2019 ⁸⁸	400/743	1826/5695	Sasi 2015 ¹⁶⁴	70/153	66/306
Kandasamy 2015 ⁹³	16/35	19/117	Schena 2015 ¹⁶⁵	28/69	77/173
Lapcharoensap 2015 ¹¹⁰	483/919	6598/14133	Soudee 2014 ¹⁷⁷	32/51	77/242
Li 2013 ¹¹⁴	9/21	49/139	Starr 2020 ¹⁷⁹	20/45	226/501
Lal 2003 ¹⁰⁹	147/401	512/2433	Tsai 2015 ¹⁸⁸	349/560	434/1120
Nobile 2017 ¹⁴¹	60/98	134/418	Wadhawan 2007 ¹⁹⁵	743/1248	4496/8213
Qiu 2012 ¹⁵⁰	471/1249	3068/10660	Westby 2009 ²⁰²	25/31	195/334

Supplementary table 12. Raw count data of the meta-analysis on the association between HDP and BPD-associated pulmonary hypertension.

Study Name	BPD-associated PH yes/total		Study Name	BPD-associated PH no/total	
	HDP-yes	HDP-no		HDP-yes	HDP-no
Blanca 2018 ¹⁸	2/10	5/56	MacKenzie 2019 ¹¹⁹	2/21	22/63
Bruno 2015 ²⁷	6/67	31/236	Mourani 2015 ¹³⁵	12/74	27/193
Carton 2016 ³¹	4/19	18/65	Naeem 2020 ¹³⁶	3/17	3/9
Check 2013 ³⁵	11/32	28/106	Ra 2013 ¹⁵¹	7/19	11/66
Choi 2019 ³⁹	3/5	14/35	Seo 2020 ¹⁶⁸	6/13	18/68
Dasgupta 2018 ⁴⁵	3/20	5/16	Sheth 2020 ¹⁷¹	15/49	44/171
Kanaan 2018 ⁹²	19/106	144/1234	Vayaltrikkovil 2019 ¹⁹²	4/18	19/103
Kim 2014 ⁹⁹	2/3	13/34	Wang 2018 ¹⁹⁷	6/13	31/141
Kunjunju 2017 ¹⁰⁷	10/17	12/39	Weismann 2017 ²⁰⁰	12/36	32/124
Lagata 2018 ¹⁰⁸	123/437	247/1240	Yallapragada 2016 ²¹⁰	2/2	5/12

Supplementary table 13. Raw count data of the meta-analysis on the association between SGA/IUGR and BPD-associated pulmonary hypertension.

Study Name	BPD-associated PH yes/total		Study Name	BPD-associated PH no/total	
	SGA/IUGR yes	SGA/IUGR no		SGA/IUGR yes	SGA/IUGR no
An 2010 ¹⁰	6/17	23/99	Khemani 2007 ⁹⁸	8/11	10/31
Aswani 2016 ¹¹	9/39	10/191	Kim 2014 ⁹⁹	3/6	12/31
Bhat 2012 ¹⁷	12/26	24/119	Lagata 2018 ¹⁰⁸	141/403	229/1274
Bruno 2015 ²⁷	10/49	27/254	Mourani 2015 ¹³⁵	9/47	30/230
Carton 2016 ³¹	2/11	20/73	Seo 2020 ¹⁶⁸	4/13	20/68
Check 2013 ³⁵	15/23	24/115	Sheth 2020 ¹⁷¹	16/28	43/192
Chen 2020 ³⁶	12/21	49/169	Vayaltrikkovil 2019 ¹⁹²	5/14	19/97
Choi 2015 ³⁹	15/52	35/122	Wang 2018 ¹⁹⁷	6/9	31/145
De Vries 2017 ⁴⁹	6/100	12/477	Weismann 2017 ²⁰⁰	9/30	35/129
Kanaan 2018 ⁹²	55/302	108/1038	Yallapragada 2016 ²¹⁰	4/5	3/9

References

1. Bramer WM, Milic J, Mast F. Reviewing retrieved references for inclusion in systematic reviews using EndNote. *JMLA*. 2017;**105**:84.
2. Wan X, Wang W, Liu J, Tong T. Estimating the sample mean and standard deviation from the sample size, median, range and/or interquartile range. *BMC Med Res Methodol* 2014;**14**:135.
3. Borenstein M, Hedges LV, Higgins J, Rothstein HR. Subgroup analyses. In: *Introduction to Meta-analysis* 2009:149-186.
4. Borenstein M, Hedges LV, Higgins J, Rothstein HR. Identifying and quantifying heterogeneity. In: *Introduction to Meta-Analysis* 2009:107-126.
5. Borenstein M, Hedges LV, Higgins J, Rothstein HR. Meta-Regression. In: *Introduction to Meta-Analysis* 2009:187-203.
6. Abramovici D, Friedman SA, Mercer BM, Audibert F, Kao L, Sibai BM. Neonatal outcome in severe preeclampsia at 24 to 36 weeks' gestation: does the HELLP (hemolysis, elevated liver enzymes, and low platelet count) syndrome matter? *Am J Obstet Gynecol*. 1999;**180**:221-225.
7. Altman M, Vanpée M, Cnattingius S, Norman M. Neonatal morbidity in moderately preterm infants: a Swedish national population-based study. *J Pediatr* 2011;**158**:239-244. e231.
8. Ameenudeen S, Boo N, Chan L. Risk factors associated with chronic lung disease in Malaysian very low birthweight infants. *Med J Malaysia* 2007;**62**:40-45.
9. Amin H, Singhal N, Sauve R. Impact of intrauterine growth restriction on neurodevelopmental and growth outcomes in very low birthweight infants. *Acta Paediatr* 1997;**86**:306-314.
10. An HS, Bae EJ, Kim GB, et al. Pulmonary hypertension in preterm infants with bronchopulmonary dysplasia. *Korean Circ J* 2010;**40**:131-136.
11. Aswani R, Hayman L, Nichols G, et al. Oxygen requirement as a screening tool for the detection of late pulmonary hypertension in extremely low birth weight infants. *Cardiol Young* 2016;**26**:521.
12. Aucott SW, Donohue PK, Northington FJ. Increased morbidity in severe early intrauterine growth restriction. *J Perinatol* 2004;**24**:435-440.
13. Baer R, Rogers E, Partridge J, et al. Population-based risks of mortality and preterm morbidity by gestational age and birth weight. *J Perinatol* 2016;**36**:1008-1013.
14. Baker CD, Balasubramanian V, Mourani PM, et al. Cord blood angiogenic progenitor cells are decreased in bronchopulmonary dysplasia. *Eur Respir J* 2012;**40**:1516-1522.
15. Balena-Borneman J, Ambalavanan N, Tiwari HK, Griffin RL, Halloran B, Askenazi D. Biomarkers associated with bronchopulmonary dysplasia/mortality in premature infants. *Pediatr Res* 2017;**81**:519-525.
16. Bardin C, Zalkowitz P, Papageorgiou A. Outcome of small-for-gestational age and appropriate-for-gestational age infants born before 27 weeks of gestation. *Pediatrics* 1997;**100**:e4-e4.
17. Bhat R, Salas AA, Foster C, Carlo WA, Ambalavanan N. Prospective analysis of pulmonary hypertension in extremely low birth weight infants. *Pediatrics* 2012;**129**:e682-e689.
18. Blanca AJ, Duijts L, van Mastrigt E, et al. Right ventricular function in infants with bronchopulmonary dysplasia and pulmonary hypertension: a pilot study. *Pulm Circ* 2018;**9**:2045894018816063.
19. Bose C, Van Marter LJ, Laughon M, et al. Fetal growth restriction and chronic lung disease among infants born before the 28th week of gestation. *Pediatrics* 2009;**124**:e450-e458.
20. Bose C, Laughon M, Allred EN, et al. Blood protein concentrations in the first two postnatal weeks that predict bronchopulmonary dysplasia among infants born before the 28th week of gestation. *Pediatr Res* 2011;**69**:347-353.
21. Bossung V, Fortmann ML, Fusch C, Rausch T, Herting E, Swoboda I, et al. Neonatal outcome after preeclampsia and HELLP syndrome: a population-based cohort study in Germany. *Front Pediatr* 2020;**8**.
22. Botet F, Figueras-Aloy J, Miracle-Echegoyen X, Rodríguez-Miguélez JM, Salvia-Roiges MD, Carbonell-Estrany X. Trends in survival among extremely-low-birth-weight infants (less than 1000 g) without significant bronchopulmonary dysplasia. *BMC pediatr* 2012;**12**:63.
23. Bozzetti V, Paterlini G, DeLorenzo P, et al. Feeding tolerance of preterm infants appropriate for gestational age (AGA) as compared to those small for gestational age (SGA). *J Matern Fetal Neonatal Med* 2013;**26**:1610-1615.
24. Brener-Dik PH, Gualdrón YMN, Galletti MF, Criolioli CM, Mariani GL. Bronchopulmonary dysplasia: incidence and risk factors. *Arch Argent Pediatr* 2017;**115**:476-482.
25. Brumbaugh JE, Colaizy TT, Patel NM, Klein JM. The changing relationship between bronchopulmonary dysplasia and cognition in very preterm infants. *Acta Paediatr* 2018;**107**:1339-1344.
26. Brumbaugh JE, Bell EF, Grey SF, et al. Behavior Profiles at 2 Years for Children Born Extremely Preterm with Bronchopulmonary Dysplasia. *J Pediatr* 2020.
27. Bruno CJ, Meerkov M, Capone C, et al. CRIB scores as a tool for assessing risk for the development of pulmonary hypertension in extremely preterm infants with bronchopulmonary dysplasia. *Am J Perinatol* 2015;**32**:1031-1037.
28. Burns Y, Gray P, O'Callaghan M. Bronchopulmonary dysplasia: a comparative study of motor development to two years of age. *Aust J Physiother*. 1997;**43**:19-25.
29. BURSAL DURAMAZ B BL, SALİHOĞLU O, ERTAŞ K, HATİPOĞLU S Neonatal outcomes of preterm infants born to preeclamptic mothers. *Marmara Med J* 2017;**30**:8-13.
30. Candel-Pau JC, Salinas FC, López JP, Lezcano AC, García OS, Olivé EL. Perinatal outcome and cardiac dysfunction in preterm growth-restricted neonates in relation to placental impairment severity. *An Pediatr (English Edition)* 2016;**85**:170-180.
31. Cartón Sánchez AJ. Hipertensión pulmonar estimada por ecocardiografía en prematuros con displasia broncopulmonar: frecuencia, evolución y factores de riesgo. *Tesis Doctoral* 2016. <https://repositorio.uam.es/handle/10486/674863>.
32. Cedergvist K, Haglund C, Heikkilä P, et al. Pulmonary trypsin-2 in the development of bronchopulmonary dysplasia in preterm infants. *Pediatrics* 2003;**112**:570-577.
33. Cetinkaya M, Ozkan H, Koksall N, Karali Z, Ozgur T. Neonatal outcomes of premature infants born to preeclamptic mothers. *J Matern Fetal Neonatal Med* 2010;**23**:425-430.
34. Charafeddine L, D'Angio CT, Phelps DL. Atypical chronic lung disease patterns in neonates. *Pediatrics* 1999;**103**:759-765.
35. Check J, Gotteiner N, Liu X, et al. Fetal growth restriction and pulmonary hypertension in premature infants with bronchopulmonary dysplasia. *J Perinatol* 2013;**33**:553-557.
36. Chen LL, Zmuda EJ, Talavera MM, et al. Dual-specificity phosphatase (DUSP) genetic variants predict pulmonary hypertension in patients with bronchopulmonary dysplasia. *Pediatr Res* 2020;**87**:81-87.

37. Chen F, Bajwa NM, Rimensberger PC, Posfay-Barbe KM, Pfister RE. Thirteen-year mortality and morbidity in preterm infants in Switzerland. *Arch Dis Child Fetal Neonatal Ed* 2016;**101**:F377-F383.
38. Cheng SW, Chou HC, Tsou KI, Fang LJ, Tsao PN. Delivery before 32 weeks of gestation for maternal pre-eclampsia: neonatal outcome and 2-year developmental outcome. *Early Hum Dev* 2004;**76**:39-46.
39. Choi EK, Jung YH, Kim H-S, et al. The impact of atrial left-to-right shunt on pulmonary hypertension in preterm infants with moderate or severe bronchopulmonary dysplasia. *Pediatr Neonatol* 2015;**56**:317-323.
40. Choi EK, Shin SH, Kim E-K, Kim H-S. Developmental outcomes of preterm infants with bronchopulmonary dysplasia-associated pulmonary hypertension at 18–24 months of corrected age. *BMC pediatr* 2019;**19**:26.
41. Chye J, Gray P. Rehospitalization and growth of infants with bronchopulmonary dysplasia: a matched control study. *J Paediatr Child Health* 1995;**31**:105-111.
42. Claas M, de Vries L, Koopman C, et al. Postnatal growth of preterm born children \leq 750 g at birth. *Early Hum Dev* 2011;**87**:495-507.
43. Cunha GS, Mezzacappa F, Ribeiro JD. Risk factors for bronchopulmonary dysplasia in very low birth weight newborns treated with mechanical ventilation in the first week of life. *J Trop Pediatr* 2005;**51**:334-340.
44. da Silva Martins A, Méio MDBB, Saint Clair SG, Lima PAT, Milanese BG, Moreira MEL. Growth and body composition in preterm newborns with bronchopulmonary dysplasia: a cohort study. *J Perinat Med* 2018;**46**:913-918.
45. Dasgupta S, Aly AM, Malloy MH, Okorodudu AO, Jain SK. NTproBNP as a surrogate biomarker for early screening of pulmonary hypertension in preterm infants with bronchopulmonary dysplasia *J Perinatol* 2018;**38**:1252-1257.
46. De Jesus LC, Pappas A, Shankaran S, et al. Outcomes of small for gestational age infants born at < 27 weeks' gestation. *J Pediatr* 2013;**163**:55-60. e53.
47. De Souza-Rugolo LM, de Sa MP, Kurokawa CS, et al. There is no difference in nitric oxide metabolites and neonatal outcome between premature infants born to pre-eclamptic and those born to normotensive women. *Paediatr Int Child Health* 2015;**35**:47-52.
48. Demirel N, Bas AY, Zenciroglu A. Bronchopulmonary dysplasia in very low birth weight infants. *Indian J Pediatr* 2009;**76**:695-698.
49. DeVries L, Heyne R, Ramaciotti C, et al. Mortality among infants with evolving bronchopulmonary dysplasia increases with major surgery and with pulmonary hypertension. *J Perinatol* 2017;**37**:1043-1046.
50. Dogan P. The role of red cell distribution width as a predictor of bronchopulmonary dysplasia in preterm infants Haydarpasa Numune. *Med J* 2019;**60**:357-361.
51. Duan J, Kong X, Li Q, et al. Association between anemia and bronchopulmonary dysplasia in preterm infants. *Sci Rep* 2016;**6**:22717.
52. Duan J, Kong X, Li Q, et al. Association between Hemoglobin Levels in the First 3 Days of Life and Bronchopulmonary Dysplasia in Preterm Infants. *Am J Perinat* 2016;**33**:998-1002.
53. Durrmeyer X, Kayem G, Sinico M, Dassieu G, Danan C, Decobert F. Perinatal risk factors for bronchopulmonary dysplasia in extremely low gestational age infants: a pregnancy disorder-based approach. *J Pediatr* 2012;**160**:578-583. e572.
54. El Ayoubi M, Jarreau P-H, Van Reempts P, et al. Does the antenatal detection of fetal growth restriction (FGR) have a prognostic value for mortality and short-term morbidity for very preterm infants? Results from the MOSAIC cohort. *J Matern Fetal Neonatal Med* 2016;**29**:596-601.
55. Engineer N, Kumar S. Perinatal variables and neonatal outcomes in severely growth restricted preterm fetuses. *Acta Obstet Gynecol Scand* 2010;**89**:1174-1181.
56. Eriksson L, Haglund B, Odland V, Altman M, Ewald U, Kieler H. Perinatal conditions related to growth restriction and inflammation are associated with an increased risk of bronchopulmonary dysplasia. *Acta Paediatr* 2015;**104**:259-263.
57. Estévez Domingo M. Análisis de la morbimortalidad a corto y medio plazo en los recién nacidos prematuros pequeños para la edad gestacional. *Tesis Doctoral* 2017. <http://hdl.handle.net/10810/31149>
58. Express group. Incidence of and risk factors for neonatal morbidity after active perinatal care: extremely preterm infants study in Sweden (EXPRESS). *Acta Paediatr* 2010;**99**:978-992.
59. Falciglia HS, Johnson JR, Sullivan JA, et al. Role of antioxidant nutrients and lipid peroxidation in premature infants with respiratory distress syndrome and bronchopulmonary dysplasia. *Am J Perinatol* 2003;**20**:097-108.
60. Fernandez-Rodriguez B, de Alba C, Villalain C, Pallás CR, Galindo A, Herraiz I. Obstetric and pediatric growth charts for the detection of fetal growth restriction and neonatal adverse outcomes in preterm newborns before 34 weeks of gestation. *J Matern Fetal Neonatal Med* 2021;**34**(7):1112-9.
61. Fujioka K, Shibata A, Yokota T, et al. Association of a vascular endothelial growth factor polymorphism with the development of bronchopulmonary dysplasia in Japanese premature newborns. *Sci Rep* 2014;**4**:4459.
62. Gagliardi L, Bellù R, Zanini R, Dammann O. Bronchopulmonary dysplasia and brain white matter damage in the preterm infant: a complex relationship. *Paediatr Perinat Epidemiol* 2009;**23**:582-590.
63. Gagliardi L, Rusconi F, Da Fre M, et al. Pregnancy disorders leading to very preterm birth influence neonatal outcomes: results of the population-based ACTION cohort study. *Pediatr Res* 2013;**73**:794-801.
64. Gagliardi L, Rusconi F, Bellu R, Zanini R. Association of maternal hypertension and chorioamnionitis with preterm outcomes. *Pediatrics* 2014;**134**:e154-161.
65. Garite TJ, Clark R, Thorp JA. Intrauterine growth restriction increases morbidity and mortality among premature neonates. *Am J Obstet Gynecol* 2004;**191**:481-487.
66. Garofoli F, Ciardelli L, Mazzucchelli I, et al. The red cell distribution width (RDW): value and role in preterm, IUGR (intrauterine growth restricted), full-term infants. *Hematology* 2014;**19**:365-369.
67. Gemmell L, Martin L, Murphy KE, et al. Hypertensive disorders of pregnancy and outcomes of preterm infants of 24 to 28 weeks' gestation. *J Perinatol* 2016;**36**:1067-1072.
68. Giapros V, Drougia A, Krallis N, Theocharis P, Andronikou S. Morbidity and mortality patterns in small-for-gestational age infants born preterm. *J Matern Fetal Neonatal Med* 2012;**25**:153-157.
69. Gortner L, Wauer RR, Stock GJ, et al. Neonatal outcome in small for gestational age infants: do they really better? *J Perinat Med* 1999;**27**:484-489.
70. Gortner L, van Husen M, Thyen U, Gembruch U, Friedrich H-J, Landmann E. Outcome in preterm small for gestational age infants compared to appropriate for gestational age preterms at the age of 2 years: a prospective study. *Eur J Obstet Gynecol Reprod Biol* 2003;**110**:S93-S97.
71. Gray PH, Hurley TM, Rogers YM, et al. Survival and neonatal and neurodevelopmental outcome of 24-29 week gestation infants according to primary cause of preterm delivery. *Aust N Z J Obstet Gynaecol* 1997;**37**:161-168.
72. Greenough A, Dimitriou G, Bhat RY, et al. Lung volumes in infants who had mild to moderate bronchopulmonary dysplasia. *Eur J Pediatr* 2005;**164**:583-586.

73. Grisar-Granovsky S, Reichman B, Lerner-Geva L, et al. Mortality and morbidity in preterm small-for-gestational-age infants: a population-based study. *Am J Obstet Gynecol* 2012;**206**:150. e151-150. e157.
74. Guimarães H, Rocha G, Vasconcellos G, et al. Factores de risco de displasia broncopulmonar em cinco unidades portuguesas de cuidados intensivos neonatais. *Rev Port Pneumol (English Edition)* 2010;**16**:419-430.
75. Guo MM-H, Chung C-H, Chen F-S, Chen C-C, Huang H-C, Chung M-Y. Severe bronchopulmonary dysplasia is associated with higher fluid intake in very low-birth-weight infants: a retrospective study. *Am J Perinatol* 2015;**32**:155-162.
76. Gursoy T, Hayran M, Derin H, Ovali F. A clinical scoring system to predict the development of bronchopulmonary dysplasia. *Am J Perinatol* 2015;**32**:659-666.
77. Hadchouel A, Decobert F, Besmond C, Delacourt C. Exome sequencing of extreme phenotypes in bronchopulmonary dysplasia. *Eur J Pediatr* 2020;**179**:579-586.
78. Hansen AR, Barnés CM, Folkman J, McElrath TF. Maternal preeclampsia predicts the development of bronchopulmonary dysplasia. *J Pediatr* 2010;**156**:532-536.
79. Hartung J, Kalache K, Heyna C, et al. Outcome of 60 neonates who had ARED flow prenatally compared with a matched control group of appropriate-for-gestational age preterm neonates. *Ultrasound Obstet Gynecol* 2005;**25**:566-572.
80. Hentges CR, Silveira RC, Procianny RS. Angiogenic and Antiangiogenic Factors in Preterm Neonates Born to Mothers with and without Preeclampsia. *Am J Perinatol* 2015;**32**:1185-1190.
81. Hernandez-Ronquillo L, Tellez-Zenteno JF, Weder-Cisneros N, Salinas-Ramirez V, Zapata-Pallagi JA, da Silva O. Risk factors for the development of bronchopulmonary dysplasia: a case-control study. *Arch Med Res* 2004;**35**:549-553.
82. Hiett AK, Brown HL, Britton KA. Outcome of infants delivered between 24 and 28 weeks' gestation in women with severe preeclampsia. *J Matern Fetal Med* 2001;**10**:301-304.
83. Hikino S, Ohga S, Kinjo T, et al. Tracheal aspirate gene expression in preterm newborns and development of bronchopulmonary dysplasia. *Pediatr Int* 2012;**54**:208-214.
84. Huusko JM, Mahlman M, Karjalainen MK, et al. Polymorphisms of the gene encoding Kit ligand are associated with bronchopulmonary dysplasia. *Pediatr Pulmonol* 2015;**50**:260-270.
85. Imamura T, Sato M, Go H, et al. The microbiome of the lower respiratory tract in premature infants with and without severe bronchopulmonary dysplasia. *Am J Perinatol* 2017;**34**:80-87.
86. Iwatani S, Mizobuchi M, Tanaka S, et al. Increased volume of tracheal aspirate fluid predicts the development of bronchopulmonary dysplasia. *Early Hum Dev* 2013;**89**:113-117.
87. Jakuskiene R, Vollmer B, Saferis V, Daugeliene D. Neonatal outcomes of very preterm infants admitted to a tertiary center in Lithuania between the years 2003 and 2005. *Eur J Pediatr* 2011;**170**:1293-1303.
88. Jensen EA, Foglia EE, Dysart KC, et al. Adverse effects of small for gestational age differ by gestational week among very preterm infants. *Arch Dis Child Fetal Neonatal Ed* 2019;**104**:F192-F198.
89. Jeon GW. Changes in the Incidence of Bronchopulmonary Dysplasia among Preterm Infants in a Single Center over 10 Years. *Neonatal Med* 2020;**27**:1-7.
90. Jo HS, Cho KH, Cho SI, Song ES, Kim BI. Recent Changes in the Incidence of Bronchopulmonary Dysplasia among Very-Low-Birth-Weight Infants in Korea. *J Korean Med Sci*. 2015;**30** Suppl 1:S81-87.
91. Kalra VK, Aggarwal S, Arora P, Natarajan G. B-type natriuretic peptide levels in preterm neonates with bronchopulmonary dysplasia: A marker of severity? *Pediatr Pulmonol* 2014;**49**:1106-1111.
92. Kanaan U, Srivatsa B, Huckaby J, Kelleman M. Association of unit-wide oxygen saturation target on incidence of pulmonary hypertension in very low birthweight premature infants. *J Perinatol* 2018;**38**:148-153.
93. Kandasamy J, Roane C, Szalai A, Ambalavanan N. Serum eotaxin-1 is increased in extremely-low-birth-weight infants with bronchopulmonary dysplasia or death. *Pediatr Res* 2015;**78**:498-504.
94. Karagianni P, Tsakalidis C, Kyriakidou M, et al. Neuromotor outcomes in infants with bronchopulmonary dysplasia. *Pediatr Neurol* 2011;**44**:40-46.
95. Karagianni P, Rallis D, Fidani L, et al. Glutathione-S-Transferase P1 polymorphisms association with bronchopulmonary dysplasia in preterm infants. *Hippokratia* 2013;**17**:363.
96. Khan MA, Kuzma-O'Reilly B, Brodsky N, Bhandari V. Site-specific characteristics of infants developing bronchopulmonary dysplasia. *J Perinatol* 2006;**26**:428-435.
97. Khan S, Concina VA, Schneider D, Westgate P, Arriagada S, Bada H. Role of NT-proBNP in the prediction of moderate to severe Bronchopulmonary Dysplasia in preterm infants. *Pediatr Pulmonol* 2020;**55**:376-382.
98. Khemani E, McElhinney DB, Rhein L, et al. Pulmonary artery hypertension in formerly premature infants with bronchopulmonary dysplasia: clinical features and outcomes in the surfactant era. *Pediatrics* 2007;**120**:1260-1269.
99. Kim D-H, Kim H-S. Serial changes of serum endostatin and angiotensin-1 levels in preterm infants with severe bronchopulmonary dysplasia and subsequent pulmonary artery hypertension. *Neonatology* 2014;**106**:55-61.
100. Kim CR, Vohr BR, Oh W. Effects of maternal hypertension in very-low-birth-weight infants. *Arch Pediatr Adolesc Med* 1996;**150**:686-691.
101. Kim DH, Shin SH, Kim EK, Kim HS. Association of increased cord blood soluble endoglin with the development of bronchopulmonary dysplasia in preterm infants with maternal preeclampsia. *Pregnancy Hypertens* 2018;**13**:148-153.
102. Kim YH, Kim KW, Eun HS, et al. Small for gestational age birth may increase airflow limitation in bronchopulmonary dysplasia. *Pediatr Pulmonol* 2020;**55**:346-353.
103. Klinger G, Sokolover N, Boyko V, et al. Perinatal risk factors for bronchopulmonary dysplasia in a national cohort of very-low-birthweight infants. *Am J Obstet Gynecol* 2013;**208**:115. e111-115. e119.
104. Koksai N, Kayik B, Çetinkaya M, et al. Value of serum and bronchoalveolar fluid lavage pro-and anti-inflammatory cytokine levels for predicting bronchopulmonary dysplasia in premature infants. *Eur Cytokine Netw* 2012;**23**:29-35.
105. Korhonen P, Laitinen J, Hyödynmaa E, Tammela O. Respiratory outcome in school-aged, very-low-birth-weight children in the surfactant era. *Acta Paediatr* 2004;**93**:316-321.
106. Koroglu OA, Yalaz M, Levent E, Akisu M, Kültürsay N. Cardiovascular consequences of bronchopulmonary dysplasia in prematurely born preschool children. *Neonatology* 2013;**104**:283-289.
107. Kunjunju A, Gopagondanahalli K, Chan Y, Sehgal A. Bronchopulmonary dysplasia-associated pulmonary hypertension: clues from placental pathology. *J Perinatol* 2017;**37**:1310-1314.
108. Lagatta JM, Hysinger EB, Zaniletti I, et al. The impact of pulmonary hypertension in preterm infants with severe bronchopulmonary dysplasia through 1 year. *The J Pediatr* 2018;**203**:218-224. e213.
109. Lal MK, Manktelow BN, Draper ES, Field DJ. Chronic lung disease of prematurity and intrauterine growth retardation: a population-based study. *Pediatrics* 2003;**111**:483-487.
110. Lapcharoensap W, Gage SC, Kan P, et al. Hospital variation and risk factors for bronchopulmonary dysplasia in a population-based cohort. *JAMA Pediatr* 2015;**169**:e143676-e143676.

111. Lardon-Fernandez M, Uberos J, Molina-Oya M, Narbona-Lopez E. Epidemiological factors involved in the development of bronchopulmonary dysplasia in very low birth-weight preterm infants. *Minerva Pediatr* 2017;**69**:42-49.
112. Lehtinen A, Korhonen P, Hyödynmaa E, et al. Adipokines played a limited role in predicting temporary growth differences between very low birthweight infants with and without bronchopulmonary dysplasia. *Acta Paediatr* 2017;**106**:1583-1588.
113. Leroy S, Caumette E, Waddington C, Hébert A, Brant R, Lavoie PM. A time-based analysis of inflammation in infants at risk of bronchopulmonary dysplasia. *J Pediatr* 2018;**192**:60-65. e61.
114. Li Y, Yan J, Li M, et al. Addition of SNAP to perinatal risk factors improves the prediction of bronchopulmonary dysplasia or death in critically ill preterm infants. *BMC pediatr* 2013;**13**:138.
115. Lin H-C, Su B-H, Chang J-S, Hsu C-M, Tsai C-H, Tsai F-J. Nonassociation of interleukin 4 intron 3 and 590 promoter polymorphisms with bronchopulmonary dysplasia for ventilated preterm infants. *Neonatology* 2005;**87**:181-186.
116. Lodha A, Sauve R, Bhandari V, et al. Need for supplemental oxygen at discharge in infants with bronchopulmonary dysplasia is not associated with worse neurodevelopmental outcomes at 3 years corrected age. *PLoS One* 2014;**9**.
117. Löfqvist C, Hellgren G, Niklasson A, et al. Low postnatal serum IGF-I levels are associated with bronchopulmonary dysplasia (BPD). *Acta Paediatr* 2012;**101**:1211-1216.
118. Lohmann P, Luna RA, Hollister EB, et al. The airway microbiome of intubated premature infants: characteristics and changes that predict the development of bronchopulmonary dysplasia. *Pediatr Res* 2014;**76**:294-301.
119. MacKenzie K, Cunningham K, Thomas S, et al. Incidence, risk factors, and outcomes of pulmonary hypertension in preterm infants with bronchopulmonary dysplasia. *J Paediatr Child Health* 2020;**25**:222-227.
120. Madoglio RJ, Rugolo LM, Kurokawa CS, Sa MP, Lyra JC, Antunes LC. Inflammatory and oxidative stress airway markers in premature newborns of hypertensive mothers. *Braz J Med Biol Res* 2016;**49**:e5160.
121. Mahlman M, Karjalainen MK, Huusko JM, et al. Genome-wide association study of broncho pulmonary dysplasia: a potential role for variants near the CRP gene. *Sci Rep* 2017;**7**:9271.
122. Malavolti AM, Bassler D, Arlettaz-Mieth R, Faldella G, Latal B, Natalucci G. Bronchopulmonary dysplasia—impact of severity and timing of diagnosis on neurodevelopment of preterm infants: a retrospective cohort study. *BMJ Paediatr Open* 2018;**2**.
123. Mao X, Qiu J, Zhao L, et al. Vitamin D and IL-10 deficiency in preterm neonates with bronchopulmonary dysplasia. *Front Pediatr* 2018;**6**:246.
124. Marshall DD, Kotelchuck M, Young TE, Bose CL, Kruyer L, O'Shea TM. Risk factors for chronic lung disease in the surfactant era: a North Carolina population-based study of very low birth weight infants. North Carolina Neonatologists Association. *Pediatrics* 1999;**104**:1345-1350.
125. Matic M, Inati V, Abdel-Latif ME, Kent AL. Maternal hypertensive disorders are associated with increased use of respiratory support but not chronic lung disease or poorer neurodevelopmental outcomes in preterm neonates at <29 weeks of gestation. *J Paediatr Child Health* 2017;**53**:391-398.
126. May C, Williams O, Milner A, et al. Relation of exhaled nitric oxide levels to development of bronchopulmonary dysplasia. *Arch Dis Child Fetal Neonatal Ed* 2009;**94**:F205-F209.
127. McGowan EC, Kostadinov S, McLean K, et al. Placental IL-10 dysregulation and association with bronchopulmonary dysplasia risk. *Pediatr Res* 2009;**66**:455-460.
128. Melamed N, Pittini A, Barrett J, et al. Antenatal corticosteroids and outcomes of small-for-gestational-age neonates. *Obstet Gynecol* 2016;**128**:1001-1008.
129. de Mello RR, Rodrigues Reis AB, da Silva KS. Cognitive performance of premature infants: association between bronchopulmonary dysplasia and cognitive skills. Cross-sectional study. *Sao Paulo Med J* 2017;**135**:383-390.
130. Méndez-Abad P, Zafra-Rodríguez P, Lubián-López S, Benavente-Fernández I. Myocardial function maturation in very-low-birth-weight infants and development of bronchopulmonary dysplasia. *Front Pediatr* 2020;**7**:556.
131. Mohamed I, Elremaly W, Rouleau T, Lavoie J-C. Oxygen and parenteral nutrition two main oxidants for extremely preterm infants: 'it all adds up'. *J Neonatal Perinatal Med* 2015;**8**:189-197.
132. Monier I, Ancel P-Y, Ego A, et al. Fetal and neonatal outcomes of preterm infants born before 32 weeks of gestation according to antenatal vs postnatal assessments of restricted growth. *Am J Obstet Gynecol* 2017;**216**:516. e511-516. e510.
133. Morrow LA, Wagner BD, Ingram DA, et al. Antenatal determinants of bronchopulmonary dysplasia and late respiratory disease in preterm infants. *Am J Respir Crit Care Med* 2017;**196**:364-374.
134. Morsing E, Gustafsson P, Brodzski J. Lung function in children born after foetal growth restriction and very preterm birth. *Acta Paediatr* 2012;**101**:48-54.
135. Mourani PM, Sontag MK, Younoszai A, et al. Early pulmonary vascular disease in preterm infants at risk for bronchopulmonary dysplasia. *Am J Respir Crit Care Med* 2015;**191**:87-95.
136. Naeem B, Ayub A, Aly AM, Malloy MH, Okorodudu AO, Jain SK. Urinary NT-proBNP as a potential noninvasive biomarker for screening of pulmonary hypertension in preterm infants: a pilot study. *J Perinatol* 2020;**40**:628-632.
137. Nascimento CP, Maia LP, Alves PT, et al. Invasive mechanical ventilation and biomarkers as predictors of bronchopulmonary dysplasia in preterm infants. *J Pediatr* 2020.
138. Natarajan G, Pappas A, Shankaran S, et al. Outcomes of extremely low birth weight infants with bronchopulmonary dysplasia: impact of the physiologic definition. *Early Hum Dev* 2012;**88**:509-515.
139. Naveda Romero OE. Risk factors for the development of bronchopulmonary dysplasia: A case-control study. [Spanish]. *Pediatrics* 2016;**49**:1-7.
140. Newman JB, DeBastos AG, Batton D, Raz S. Neonatal respiratory dysfunction and neuropsychological performance at the preschool age: a study of very preterm infants with bronchopulmonary dysplasia. *Neuropsychology* 2011;**25**:666-678.
141. Nobile S, Marchionni P, Carnielli VP. Neonatal outcome of small for gestational age preterm infants. *European J Pediatr* 2017;**176**:1083-1088.
142. Novitsky A, Tuttle D, Locke RG, Saiman L, Mackley A, Paul DA. Prolonged Early Antibiotic Use and Bronchopulmonary Dysplasia in Very Low Birth Weight Infants. *Am J Perinatol* 2015;**32**:043-048.
143. O'Shea JE, Davis PG, Doyle LW, Victorian Infant C. Maternal preeclampsia and risk of bronchopulmonary dysplasia in preterm infants. *Pediatr Res* 2012;**71**:210-214.
144. Oh W, Poindexter BB, Perritt R, et al. Association between fluid intake and weight loss during the first ten days of life and risk of bronchopulmonary dysplasia in extremely low birth weight infants. *J Pediatr* 2005;**147**:786-790.
145. Ozkan H, Cetinkaya M, Koksall N. Increased incidence of bronchopulmonary dysplasia in preterm infants exposed to preeclampsia. *J Matern Fetal Neona* 2012;**25**:2681-2685.
146. Palta M, Sadek M, Gabbert D, et al. The relation of maternal complications to outcomes in very low birthweight infants in an era of changing neonatal care. *American J Perinatol* 1996;**13**:109-114.
147. Park YH, Lee GM, Yoon JM, et al. Effect of early postnatal neutropenia in very low birth weight infants born to mothers with pregnancy-induced hypertension. *Korean J Pediatr* 2012;**55**:462-469.

148. Patel AL, Johnson TJ, Robin B, et al. Influence of own mother's milk on bronchopulmonary dysplasia and costs. *Arch Dis Child Fetal Neonatal Ed* 2017;**102**:F256-F261.
149. Qi Y, Jiang Q, Chen C, Cao Y, Qian L. Circulating endothelial progenitor cells decrease in infants with bronchopulmonary dysplasia and increase after inhaled nitric oxide. *PLoS One* 2013;**8**:e79060.
150. Qiu X, Lodha A, Shah PS, et al. Neonatal outcomes of small for gestational age preterm infants in Canada. *Am J Perinatol* 2012;**29**:87-94.
151. Ra JJ, Lee SM, Eun HS, et al. Risk factors of pulmonary hypertension in preterm infants with chronic lung disease. 2013.
152. Rakza T, Magnenant E, Klosowski S, Tourneux P, Bachiri A, Storme L. Early hemodynamic consequences of patent ductus arteriosus in preterm infants with intrauterine growth restriction. *J Pediatr* 2007;**151**:624-628.
153. Ranke MB, Vollmer B, Traunecker R, et al. Growth and development are similar in VLBW children born appropriate and small for gestational age: an interim report on 97 preschool children. *J Pediatr Endocrinol Metab* 2007;**20**:1017-1026.
154. Razak A, Patel W, Durrani N, et al. Neonatal respiratory outcomes in pregnancy induced hypertension: introducing a novel index. *J Matern Fetal Neonatal Med* 2020;**33**:625-632.
155. Redline RW, Wilson-Costello D, Hack M. Placental and other perinatal risk factors for chronic lung disease in very low birth weight infants. *Pediatr Res* 2002;**52**:713-719.
156. Regev RH, Arnon S, Litmanovitz I, et al. Outcome of singleton preterm small for gestational age infants born to mothers with pregnancy-induced hypertension. A population-based study. *J Matern Fetal Neonatal Med* 2015;**28**:666-673.
157. Regev RH, Lusky A, Dolfin T, et al. Excess mortality and morbidity among small-for-gestational-age premature infants: a population-based study. *J Pediatr* 2003;**143**:186-191.
158. Reiss I, Landmann E, Heckmann M, Misselwitz B, Gortner L. Increased risk of bronchopulmonary dysplasia and increased mortality in very preterm infants being small for gestational age. *Arch Gynecol Obstet* 2003;**269**:40-44.
159. Rijken M, Wit JM, Veen S. Similar growth in preterm infants with intra- or extrauterine growth restriction. *A Regional Follow-Up Study at Two Years of Age in Extremely Preterm and Very Preterm Infants* 2007;**73**.
160. Rocha G, de Lima FF, Machado AP, Guimarães H. Preeclampsia predicts higher incidence of bronchopulmonary dysplasia. *J Perinatol* 2018;**38**:1165-1173.
161. Rocha G, Ribeiro O, Guimarães H. Fluid and electrolyte balance during the first week of life and risk of bronchopulmonary dysplasia in the preterm neonate. *Clinics* 2010;**65**:663-674.
162. Rojas MX, Rojas MA, Lozano JM, Rondón MA, Charry LP. Regional variation on rates of bronchopulmonary dysplasia and associated risk factors. *ISRN Pediatrics* 2012;**2012**: 685151.
163. Ruiz-Peláez JG, Charpak N. Epidemia de displasia broncopulmonar: incidencia y factores asociados en una cohorte de niños prematuros en Bogotá, Colombia. *Biomédica* 2014;**34**:29-39.
164. Sasi A, Abraham V, Davies-Tuck M, et al. Impact of intrauterine growth restriction on preterm lung disease. *Acta Paediatr* 2015;**104**:e552-e556.
165. Schena F, Francescato G, Cappelleri A, et al. Association between hemodynamically significant patent ductus arteriosus and bronchopulmonary dysplasia. *J Pediatr* 2015;**166**:1488-1492.
166. Schiff E, Friedman SA, Mercer BM, Sibai BM. Fetal lung maturity is not accelerated in preeclamptic pregnancies. *Am J Obstet Gynecol* 1993;**169**:1096-1101.
167. Schlapbach LJ, Ersch J, Adams M, Bernet V, Bucher HU, Latal B. Impact of chorioamnionitis and preeclampsia on neurodevelopmental outcome in preterm infants below 32 weeks gestational age. *Acta Paediatr* 2010;**99**:1504-1509.
168. Seo YM, Yum SK, Sung IK. Respiratory Severity Score with Regard to Birthweight during the Early Days of Life for Predicting Pulmonary Hypertension in Preterm Infants. *J Trop Pediatr* 2020.
169. Serenius F, Ewald U, Farooqi A, Holmgren PÅ, Håkansson S, Sedin G. Short-term outcome after active perinatal management at 23–25 weeks of gestation. A study from two Swedish tertiary care centres. Part 2: infant survival. *Acta Paediatr* 2004;**93**:1081-1089.
170. Sharma P, McKay K, Rosenkrantz TS, Hussain N. Comparisons of mortality and pre-discharge respiratory outcomes in small-for-gestational-age and appropriate-for-gestational-age premature infants. *BMC pediatr* 2004;**4**:9.
171. Sheth S, Goto L, Bhandari V, Abraham B, Mowes A. Factors associated with development of early and late pulmonary hypertension in preterm infants with bronchopulmonary dysplasia. *J Perinatol* 2020;**40**:138-148.
172. Shima Y, Kumasaka S, Migita M. Perinatal risk factors for adverse long-term pulmonary outcome in premature infants: Comparison of different definitions of bronchopulmonary dysplasia/chronic lung disease. *Pediatr Int* 2013;**55**:578-581.
173. Shin SH, Shin SH, Kim SH, et al. The Association of Pregnancy-induced Hypertension with Bronchopulmonary Dysplasia—A Retrospective Study Based on the Korean Neonatal Network database. *Sci Rep* 2020;**10**:1-6.
174. Silveira RC, Procianny RS. Growth and neurodevelopment outcome of very low birth weight infants delivered by preeclamptic mothers. *Acta Paediatr* 2007;**96**:1738-1742.
175. Soliman N, Chaput K, Alshaikh B, Yusuf K. Preeclampsia and the risk of bronchopulmonary dysplasia in preterm infants less than 32 weeks' gestation. *Am J Perinatol* 2017;**34**:585-592.
176. Somaschini M, Castiglioni E, Volonteri C, Cursi M, Ferrari M, Carrera P. Genetic predisposing factors to bronchopulmonary dysplasia: preliminary data from a multicentre study. *J Matern Fetal Neonatal Med* 2012;**25**:119-122.
177. Soudée S, Vuillemin L, Alberti C, et al. Fetal growth restriction is worse than extreme prematurity for the developing lung. *Neonatology* 2014;**106**:304-310.
178. Spiegler J, Stichtenoth G, Weichert J, et al. Pregnancy risk factors for very premature delivery: what role do hypertension, obesity and diabetes play? *Arch Gynecol Obstet* 2013;**288**:57-64.
179. Starr MC, Boohaker L, Eldredge LC, et al. Acute kidney injury and bronchopulmonary dysplasia in premature neonates born less than 32 weeks' gestation. *Am J Perinatol* 2020;**37**:341-348.
180. Streubel A, Donohue PK, Aucott SW. The epidemiology of atypical chronic lung disease in extremely low birth weight infants. *J Perinatol* 2008;**28**:141-148.
181. Sun Y, Chen C, Zhang X, et al. High neutrophil-to-lymphocyte ratio is an early predictor of bronchopulmonary dysplasia. *Front Pediatr* 2019;**7**:464.
182. Tammela OK, Koivisto ME. A 1-year follow-up of low birth weight infants with and without bronchopulmonary dysplasia: health, growth, clinical lung disease, cardiovascular and neurological sequelae. *Early Hum Dev* 1992;**30**:109-120.
183. Teberg AJ, Pena I, Finello K, Aguilar T, Hodgman JE. Prediction of neurodevelopmental outcome in infants with and without bronchopulmonary dysplasia. *Am J Med Sci* 1991;**301**:369-374.
184. Todd DA, Jana A, John E. Chronic oxygen dependency in infants born at 24-32 weeks' gestation: The role of antenatal and neonatal factors. *J Paediatr Child Health* 1997;**33**:402-407.
185. Tokumasu H, Tokumasu S, Kawakami K. Impact of pre-eclampsia in extremely premature infants: Population-based study. *Pediatr Int* 2016;**58**:578-583.

186. Torchin H, Ancel PY, Goffinet F, et al. Placental Complications and Bronchopulmonary Dysplasia: EPIPAGE-2 Cohort Study. *Pediatrics* 2016;**137**:e20152163.
187. Torrance HL, Mulder EJ, Brouwers HA, van Bel F, Visser GH. Respiratory outcome in preterm small for gestational age fetuses with or without abnormal umbilical artery Doppler and/or maternal hypertension. *J Matern Fetal Neonat* 2007;**20**:613-621.
188. Tsai L-Y, Chen Y-L, Tsou K-I, Mu S-C, Group TPIDCS. The impact of small-for-gestational-age on neonatal outcome among very-low-birth-weight infants. *Pediatr Neonatol* 2015;**56**:101-107.
189. Turunen R, Andersson S, Laivuori H, et al. Increased postnatal inflammation in mechanically ventilated preterm infants born to mothers with early-onset preeclampsia. *Neonatology* 2011;**100**:241-247.
190. Uberos J, Jimenez-Montilla S, Molina-Oya M, Garcia-Serrano JL. Early energy restriction in premature infants and bronchopulmonary dysplasia: A cohort study. *Br J Nutr* 2020;**123**:1024-1031.
191. van Mastrigt E, Zweekhorst S, Bol B, et al. Ceramides in tracheal aspirates of preterm infants: Marker for bronchopulmonary dysplasia. *PLoS one* 2018;**13**:e0185969.
192. Vayaltrikkovil S, Vorhies E, Stritzke A, et al. Prospective study of pulmonary hypertension in preterm infants with bronchopulmonary dysplasia. *Pediatr Pulmonol* 2019;**54**:171-178.
193. Viscardi RM, Muhumuza CK, Rodriguez A, et al. Inflammatory markers in intrauterine and fetal blood and cerebrospinal fluid compartments are associated with adverse pulmonary and neurologic outcomes in preterm infants. *Pediatr Res* 2004;**55**:1009-1017.
194. Vrlench LA, Bozynski ME, Shyr Y, Schork MA, Roloff DW, McCormick MC. The effect of bronchopulmonary dysplasia on growth at school age. *Pediatrics* 1995;**95**:855-859.
195. Wadhawan R, Oh W, Perritt R, et al. Association between early postnatal weight loss and death or BPD in small and appropriate for gestational age extremely low-birth-weight infants. *J Perinatol* 2007;**27**:359-364.
196. Wang K, Huang X, Lu H, Zhang Z. A comparison of KL-6 and Clara cell protein as markers for predicting bronchopulmonary dysplasia in preterm infants. *Dis Markers*. 2014;**2014**:736536.
197. Wang C-H, Shi L-P, Ma X-L, DU L. Clinical features and prognosis of bronchopulmonary dysplasia complicated by pulmonary hypertension in preterm infants. *Chin J Contemp Pediatr* 2018;**20**:893-896.
198. Wang H, Julien KRS, Stevenson DK, et al. A genome-wide association study (GWAS) for bronchopulmonary dysplasia. *Pediatrics* 2013;**132**:290-297.
199. Wang LY, Luo HJ, Hsieh WS, et al. Severity of bronchopulmonary dysplasia and increased risk of feeding desaturation and growth delay in very low birth weight preterm infants. *Pediatr Pulmonol* 2010;**45**:165-173.
200. Weismann C, Asnes J, Bazy-Asaad A, Tolomeo C, Ehrenkranz R, Bizzarro M. Pulmonary hypertension in preterm infants: results of a prospective screening program. *J Perinatol* 2017;**37**:572-577.
201. Wemhöner A, Ortner D, Tschirch E, Strasak A, Rüdiger M. Nutrition of preterm infants in relation to bronchopulmonary dysplasia. *BMC Pulm Med* 2011;**11**:1-6.
202. Westby-Wold S, Sommerfelt K, Reigstad H, et al. Neonatal mortality and morbidity in extremely preterm small for gestational age infants: a population based study. *Arch Dis Child Fetal Neonatal Ed* 2009;**94**:F363-F367.
203. Wilmsink FA, Reijnierse J, Reiss IK, Steegers EA, de Jonge RC. Preeclampsia and risk of developing bronchopulmonary dysplasia in very preterm neonates. *Pregnancy Hypertens* 2019;**15**:57-63.
204. Wirbelauer J, Thomas W, Rieger L, Speer CP. Intrauterine growth retardation in preterm infants \leq 32 weeks of gestation is associated with low white blood cell counts. *Am J Perinatol* 2010;**27**:819-824.
205. Withagen ML, Visser W, Wallenburg HC. Neonatal outcome of temporizing treatment in early-onset preeclampsia. *Eur J Obstet Gynecol Reprod Biol* 2001;**94**:211-215.
206. Wocadlo C, Rieger I. Developmental outcome at 12 months corrected age for infants born less than 30 weeks gestation: influence of reduced intrauterine and postnatal growth. *Early Hum Dev* 1994;**39**:127-137.
207. Won Choi C, Il Kim B, Kim HS, Dong Park J, Choi JH, Woo Son D. Increase of interleukin-6 in tracheal aspirate at birth: A predictor of subsequent bronchopulmonary dysplasia in preterm infants. *Acta Paediatr*. 2006;**95**:38-43.
208. Xu Y-P. Bronchopulmonary dysplasia in preterm infants born at less than 32 weeks gestation. *Global Pediatric Health* 2016;**3**:2333794X16668773.
209. Xydis V, Drougia A, Giapros V, Argyropoulou M, Andronikou S. Brain growth in preterm infants is affected by the degree of growth restriction at birth. *J Matern Fetal Neonatal Med* 2013;**26**:673-679.
210. Yallapragada SG, Mestan KK, Palac H, et al. Placental villous vascularity is decreased in premature infants with bronchopulmonary dysplasia-associated pulmonary hypertension. *Pediatr Dev Pathol* 2016;**19**:101-107.
211. Yen TA, Yang HI, Hsieh WS, et al. Preeclampsia and the risk of bronchopulmonary dysplasia in VLBW infants: a population based study. *PLoS one* 2013;**8**:e75168.
212. Yılmaz C, Köksal N, Özkan H, Dorum BA, Bağcı O. Low serum IGF-1 and increased cytokine levels in tracheal aspirate samples are associated with bronchopulmonary dysplasia. *Turk J Pediatr* 2017;**59**:122-129.
213. Yoon BH, Romero R, Kim KS, et al. A systemic fetal inflammatory response and the development of bronchopulmonary dysplasia-An elevated. *Am J Obstet Gynecol* 1999;**181**:773-779.
214. Yu HJ, Kim ES, Kim JK, et al. Outcomes of small for gestational age micropremies depending on how young or how small they are. *Korean J. Pediatr* 2011;**54**:246.
215. Zanardo V, Savio V, Giacomini C, Rinaldi A, Marzari F, Chiarelli S. Relationship between neonatal leukemoid reaction and bronchopulmonary dysplasia in low-birth-weight infants: a cross-sectional study. *Am J Perinatol* 2002;**19**:379-386.
216. Zhang H, Fang J, Su H, Chen M. Risk factors for bronchopulmonary dysplasia in neonates born at \leq 1500 g (1999–2009). *Pediatr Int* 2011;**53**:915-920.