

**LETTER****Television viewing and asthma: spurious relationship?**

In the April 2009 issue of *Thorax*, Sherriff and co-authors report on data taken from the ALSPAC study, addressing the association between television viewing in early childhood and the development of asthma.<sup>1</sup> They found that, after adjustment for body mass index, there was a relationship between the two, showing a significant trend.

I was surprised to see that television viewing was viewed solely as a proxy for a sedentary lifestyle, but not as being asso-

ciated with other risk factors for developing asthma. For example, although the authors corrected for smoking during pregnancy, they did not include parental smoking at home in their model. It is not unlikely that among parents of children that were reported to have been watching television for longer, many of them were smoking in the presence of their child.

Adjustment for such additional factors is warranted before discussing the consequences of the study findings.

**Johannes C van der Wouden**

**Correspondence to** Johannes C van der Wouden, Department of General Practice, Erasmus MC, Room

Ff325, PO Box 2040, Rotterdam, CA 3000, The Netherlands; j.vanderwouden@erasmusmc.nl

**Competing interests** None.

**Provenance and peer review** Not commissioned; externally peer reviewed.

Accepted 21 August 2009

*Thorax* 2010;■:1. doi:10.1136/thx.2009.121111

**REFERENCE**

1. **Sherriff A**, Maitra A, Ness AR, *et al*. Association of duration of television viewing in early childhood with the subsequent development of asthma. *Thorax* 2009;64:321–5.