

**Lung cancer survival and stage at diagnosis in Australia, Canada, Denmark, Norway, Sweden and the United Kingdom: a population-based study, 2004-2007**

**WEB APPENDIX**

**Web appendix table 1** Stage-specific sets of weights used for age standardisation of non-small cell lung cancer estimates

**Web appendix table 2** Stage-specific sets of weights used for age standardisation of small-cell lung cancer estimates

**Web appendix figure 1** Proportions of non-small cell (top panel) and small-cell (bottom panel) lung cancer patients with missing data on stage and observed cumulative stage distribution by age at diagnosis and country, TNM and SEER Summary Stage 2000

**Web appendix table 1. Stage-specific sets of weights used for age standardisation of non-small cell lung cancer estimates**


<b>TNM stage</b>	<b>Age</b>	<b>Weights</b>	<b>SEER Summary Stage 2000</b>	<b>Age</b>	<b>Weights</b>
All patients	15-44	0.01	All patients	15-44	0.02
	45-54	0.07		45-54	0.08
	55-64	0.22		55-64	0.23
	65-74	0.32		65-74	0.32
	75-84	0.31		75-84	0.29
	85-99	0.07		85-99	0.07
TNM stage I	15-44	0.02	Localised	15-44	0.02
	45-54	0.06		45-54	0.06
	55-64	0.22		55-64	0.22
	65-74	0.34		65-74	0.34
	75-84	0.30		75-84	0.30
	85-99	0.06		85-99	0.06
TNM stage II	15-44	0.01	Regional	15-44	0.02
	45-54	0.07		45-54	0.07
	55-64	0.23		55-64	0.24
	65-74	0.34		65-74	0.34
	75-84	0.29		75-84	0.28
	85-99	0.05		85-99	0.06
TNM stage III	15-44	0.01	Distant	15-44	0.02
	45-54	0.07		45-54	0.10
	55-64	0.22		55-64	0.25
	65-74	0.33		65-74	0.32
	75-84	0.30		75-84	0.26
	85-99	0.07		85-99	0.05
TNM stage IV	15-44	0.02	Missing stage	15-44	0.01
	45-54	0.09		45-54	0.05
	55-64	0.24		55-64	0.15
	65-74	0.33		65-74	0.27
	75-84	0.27		75-84	0.38
	85-99	0.05		85-99	0.14
Missing stage	15-44	0.01			
	45-54	0.04			
	55-64	0.15			
	65-74	0.27			
	75-84	0.39			
	85-99	0.14			

**Web appendix table 2. Stage-specific sets of weights used for age standardisation of small-cell lung cancer estimates**

<b>TNM stage</b>	<b>Age</b>	<b>Weights</b>	<b>SEER Summary Stage 2000</b>	<b>Age</b>	<b>Weights</b>
All patients	15-59	0.21	All patients	15-59	0.21
	60-74	0.53		60-74	0.53
	75-99	0.26		75-99	0.26
TNM stage I & II	15-59	0.18	Localised	15-59	0.19
	60-74	0.53		60-74	0.52
	75-99	0.29		75-99	0.29
TNM stage III	15-59	0.21	Regional	15-59	0.21
	60-74	0.55		60-74	0.54
	75-99	0.24		75-99	0.25
TNM stage IV	15-59	0.22	Distant	15-59	0.22
	60-74	0.55		60-74	0.53
	75-99	0.24		75-99	0.25
Missing stage	15-59	0.18	Missing stage	15-59	0.18
	60-74	0.49		60-74	0.48
	75-99	0.32		75-99	0.33

Web appendix figure 1. Proportions of non-small cell (top panel) and small-cell (bottom panel) lung cancer patients with missing data on stage and observed cumulative stage distribution by age at diagnosis and country, TNM and SEER Summary Stage 2000

**NON-SMALL CELL LUNG CANCER**



**SMALL-CELL LUNG CANCER**

