

Supplementary Table 2: Patient-reported outcome (PRO) results [mean (standard deviation)]

	50 mg once daily			100 mg twice daily		
		Change from Baseline			Change from Baseline	
		Baseline	24 weeks		Baseline	24 weeks
Patients, n	46	40	36	42	35	29
Leicester Cough Questionnaire						
Total	15.74 (4.11)	0.22 (3.61)	-1.28 (3.51)	16.67 (3.44)	-0.44 (3.11)	-0.53 (2.84)
Physical	5.07 (1.20)	0.15 (1.06)	-0.28 (1.08)	5.32 (1.03)	0.02 (0.94)	-0.05 (0.91)
Psychological	5.23 (1.54)	0.06 (1.34)	-0.46 (1.22)	5.63 (1.31)	-0.22 (1.27)	-0.20 (0.97)
Social	5.43 (1.50)	0.01 (1.36)	-0.54 (1.38)	5.72 (1.26)	-0.24 (1.11)	-0.29 (1.15)
King's Brief ILD Questionnaire						
Total	53.5 (14.0)	-0.2 (8.1)	-0.3 (11.4)	57.2 (10.9)	3.1 (13.2)	2.1 (15.5)
Breathlessness and activities	40.0 (18.1)	-2.4 (13.3)	-2.3 (15.9)	45.0 (17.7)	6.3 (16.7)	3.8 (20.1)
Chest symptoms	64.8 (23.6)	1.2 (17.6)	-1.0 (18.4)	70.1 (19.5)	0.7 (16.5)	-1.0 (19.1)
Psychological	53.7 (20.0)	2.2 (13.3)	-0.5 (15.1)	58.1 (15.3)	3.4 (19.5)	3.2 (22.5)

ILD: interstitial lung disease.